

COMPARATIVE STUDIES OF THE EFFECT OF BSA VS FCS AS A SUPPLEMENT IN TCM-199 ON THE *IN VITRO* MATURATION RATE OF BUFFALO OOCYTES COLLECTED FROM SLAUGHTERHOUSE OVARIES

Yosef Deneke^{1,*}, Prem Singh Yadav², Rajib Deb³ and Trilok Nanda⁴

ABSTRACT

Availability of developmentally competent buffalo oocytes is critical for *in vitro* embryo production and application of related biotechniques. The objective of the present study was to assess the effect of BSA in place of FCS as maturation medium supplement on *in vitro* maturation buffalo oocytes. Oocytes were aspirated from abattoir ovarian follicles of 2-8 mm diameter followed by maturation in TCM-199 supplemented with hCG, PMSG and containing either 0.4% BSA (group-I) or 10% FCS (group-II). Based on cumulus expansion maturation rate, it was assessed that among the two groups, Group 2 showed significantly higher percentage values (89.1±3.5%) as compared to Group 1 (73.9±4.2%).

Keywords: buffalo oocytes, *in vitro* maturation, BSA, FCS

INTRODUCTION

In buffaloes, embryo transfer has had limited success as compared to other livestock species. The buffalo has low productive capacity, evidenced by a lower number of follicles in the

ovary (Agrawal and Tomar, 1998), a high percentage of atretic follicles, changes in acrosomal protein and membrane damage during freezing, because freezing buffalo semen results in acrosomal damage mediated leakage of enzymes, alteration of pH, complete withdrawal of the hydration shell of protein in solution and loss of sperm motility (Nandi *et al.*, 2003).

In vitro maturation (IVM) of unfertilized oocytes has great potential for cattle breeding especially when combined with *in vitro* fertilization (IVF) and *in vitro* culture (IVC) and cryopreservation techniques. The culture medium employed for IVM is important in view of its effect on the maturation rate of follicular oocytes and also on embryonic development following IVF (Bavister *et al.*, 1992). The commonly used media are complex, buffered with bicarbonate or HEPES and supplemented with various sera and /or gonadotrophin (FSH/LH) and/or steroid (estradiol 17 β) hormones. It is known that the culture conditions employed for IVM of mammalian oocytes can significantly influence IVF rates and subsequent embryonic development (Brackett *et al.*, 1989; Abdoon *et al.*, 2001). Various types of media viz., TCM-199 (Singh *et al.*, 1989; Totey *et al.*, 1993; Madan *et al.*, 1994; Nandi *et al.*, 2000), Ham's F-10 (Singh *et al.*, 1989; Totey *et al.*, 1993), MEM (Abdoon *et al.*, 2001) have been

¹Department of Animal Biotechnology, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India, E-mail: yosefdeneke@yahoo.com

²Central Institute for Research on Buffaloes, Hisar, Haryana, India

³Division of Animal Biotechnology, Indian Veterinary Research Institute, Izatnagar, Uttar Pradesh, India

commonly used for IVM-IVF studies in buffaloes. Among them, TCM-199 is the most commonly used medium. These complex media cannot support oocyte maturation on their own and are usually supplemented with hormones (Totey *et al.*, 1993; Nandi *et al.*, 2000), sera (Bacci *et al.*, 1991; Lu and Hsu, 1991; Madan *et al.*, 1994; Totey *et al.*, 1996) or follicular fluid (Chauhan *et al.*, 1997) which introduce many known and unknown substances to the IVM medium for proper maturation. In the present studies we have evaluated the effect of bovine serum albumin (BSA) and fetal calf serum (FCS) as supplements in TCM-199 as a basic *in vitro* maturation medium for buffalo oocytes.

MATERIALS AND METHODS

Collection of ovaries

Buffalo ovaries were collected from a Delhi abattoir in sterile normal saline solution (NSS) supplemented with antibiotics (penicillin 100 IU/ml, streptomycin 50 µg/ml, Hi-Media, India) at 30-35°C in a thermos flask and transported to the laboratory within 4 h of slaughter.

Retrieval of oocytes

In the laboratory, the surrounding tissues were trimmed off and the ovaries were washed four to five times with sterile and warm (30-35°C) NSS. The ovaries were then exposed to 70% ethyl alcohol for 2 seconds and finally washed with phosphate buffered saline and immediately blotted with paper towel (Hurt *et al.*, 2000). Oocytes from ovarian follicles of 2 to 8 mm in diameter were aspirated using 18G needle attached to 5 ml sterile disposable syringe (Dispovan, India) containing 0.5 ml aspiration medium.

The aspiration medium consisted of phosphate buffered saline (Gibco, USA) supplemented with 0.4% fatty acid free embryo tested bovine serum albumin (BSA) (Sigma, USA) and antibiotics. The aspirated suspension was poured into a sterile 90 mm Petri dish (Griener, Germany) having 4 to 6 ml aspiration medium. The dishes were searched under a stereo zoom microscope (Olympus, Japan). Cumulus-oocytes complexes (COCs) were located and picked up from the Petri dish. These oocytes were then washed serially three times with TCM-199 (Gibco, USA), supplemented with 0.4% BSA or 10% fetal calf serum (FCS) (Gibco, Mexico) as per requirement in 35 mm petridish (Griener, Germany) containing 2 ml of the maturation media and finally transferred into washing drops of 100 µl in a 60 mm Petri dish (Griesner, Germany) for a serial wash in 3 drops horizontally. The COCs were graded as described by Nandi *et al.* (1998):

Grade-I: Compact cumulus oocyte complexes with unexpanded cumulus mass having ≥ 5 layers of cumulus cells and homogenous evenly granular ooplasm.

Grade-II: COCs similar to Grade-I but with 2-4 layers of cumulus cells.

Grade-III: Oocytes with partially denuded or completely devoid of cumulus cells and having an irregular dark ooplasm.

Grade-IV: Oocytes with highly expanded or scattered cumulus cells and an irregular dark ooplasm.

The oocytes of Grades I and II were used for *in vitro* maturation within 2 h of their removal from the follicles. Oocytes were matured in two groups. Group1 consisted of oocytes matured in TCM-199 supplemented with 0.4% BSA and Group 2 those matured with 10% FCS supplementation.

***In vitro* maturation of oocytes**

The maturation media included TCM-199 that was supplemented with either (a) 0.4% BSA or (b) 10% FCS (heat inactivated in water bath at 56°C for 30 minutes) and streptomycin (100 µg/ml, penicillin 100 IU/ml, Sigma, USA), l-glutamine (200 mM, Sigma, USA) and hormones PMSG (10 IU/ml) and hCG (10 IU/ml) (Sigma, USA). After washing twice with 2 ml maturation medium (without hormone supplement) in a 35 mm sterile Petri dish (Griener, Germany), 15-20 COCs were randomly placed in 100 µl of maturation drops (medium + hormone) (Shamsuddin *et al.*, 1993) in a 35 mm sterile Petri dish. The maturation drops were covered with warm (35-37°C) light weight mineral oil (Sigma, USA) and kept for 24 h in a CO₂ incubator at 38.5°C under a condition of 5% CO₂ in air with a relative humidity of 90 to 95%.

Maturation of the oocytes was evaluated after 24 h of culture to access the degree of cumulus cell expansion under a stereo zoom microscope and also the appearance of the polar body using the methods described by Kobayashi *et al.* (1994) and Nandi *et al.* (1998).

Degree-0 (slight or no expansion of cumulus cells): oocytes having cumulus cells tightly adherent to the zona pellucida,

Degree-1 (moderate cumulus cell expansion): oocytes having expansion of the cumulus cell mass to 2-diameter away from zona pellucida, cells were homogenously spread and

cluster cells were still observed.

Degree-2 (full cumulus cell expansion): oocytes showed enlargement of cumulus cell mass to at least 3-diameters away from the zona pellucida, cells were homogenously spread and clustered cells were no longer present.

RESULTS

A total number of 1954 oocytes were cultured for 24 h in a CO₂ incubator with 5% CO₂ in air under humidified conditions at 38.5°C and their maturation was assessed by the expansion of their cumulus cells and formation of polar bodies (Microphotographs 1, 2, Plate-2). Based on cumulus expansion maturation rate was assessed in two groups which resulted in 819 matured out of 1108 oocytes cultured in TCM-199 supplemented with 0.4% BSA (group-I) and 754 oocytes matured out of 846 cultured in TCM-199 with 10% FCS (group-II), showing $73.9 \pm 4.2\%$ and $89.1 \pm 3.5\%$ maturation rates in the two, respective groups (Table 1).

DISCUSSION

It has been accepted that the expansion of cumulus cells is important to achieve complete oocyte maturation since it is correlated to the

Table 1. Oocytes maturation rates in TCM-199 with supplements 0.4% BSA vs 10% FCS.

TCM-199 Supplements	No. of oocytes used	No. of oocytes matured	% Maturation rate
0.4% BSA	1108	819±6.6	73.9±4.2
10% FCS	846	754±4.2	89.1±3.5

fertilization rate and developmental potential in ovine and bovine oocytes (Saigmillier and Moor, 1984; Cox *et al.*, 1993). Based on cumulus-cell expansion, in the present study, about 73.9±4.2% in Group 1 and 89.1±3.5% in Group 2 were considered to be mature after 24 h of *in vitro* maturation.

In buffalo oocytes, 80 percent maturation has been reported by Totey *et al.* (1993) in TCM-199 supplemented with FSH, LH and estradiol. Thus, the results in the present study are comparable with those of Totey *et al.* (1992; 1993a), Singh *et al.* (1989), Chauhan *et al.* (1991), Madan *et al.* (1994 b) and Das *et al.* (1996b). Brackett *et al.* (1989) reported 95-100 percent maturation of cow oocytes in the above medium. When the medium is supplemented with other supplements like hormones and follicular fluid, the maturation rate becomes higher: 95-100% (Chauhan *et al.*, 1997). The differences in maturation rates may be due to a number of factors like health of oocytes at the time of collection, physiological and nutritional status of slaughtered animals, time elapsed during transportation of ovaries and composition of the medium. Das *et al.* (1996b) and Chauhan *et al.* (1997) did not have desired maturation rates from buffalo oocytes in TCM-199 with FCS and FSH.

Fukuda *et al.* (1990) reported 74 percent maturation rate for bovine oocytes cultured in TCM-199 supplemented with 10% bovine estrus serum (BES). Totey *et al.* (1993) also reported a maturation rate of 76% when oocytes were matured in TCM-199 supplemented with hormones and BES.

Cumulus cells are important during oocyte maturation and contribute to the production of cytoplasmic maturation factors (Vanderhyden and Armstorng, 1989) and prevention of hardening of the zona pellucida (De Felici and Siracusa, 1982). Additionally, the cumulus cells secrete non-

sulfated glycosaminoglycan and hyaluronic acid (Ball *et al.*, 1983), when stimulated by FSH, which causes their dispersion, forming a mucous matrix between and around cumulus cells. Hyaluronic acid promotes the acrosome reaction of epididymal bovine spermatozoa (Handrow *et al.*, 1982) and the dispersed cumulus cells makes it easier for sperm cells to reach the oocytes, although oocytes without expanded cumulus cells have been fertilized *in vitro* with high frequency (Schroeder and Eppig, 1994; Sirard *et al.*, 1988).

ACKNOWLEDGEMENT

The first author is thankful to the CCS HAU and to the Director, Central Institute for Research on Buffaloes, Hisar, Haryana, India for providing necessary facilities to conduct the present research work.

REFERENCES

- Agrawal, S.K. and O.S. Tomer. 1998. *Reproductive Technologies in Buffalo*. Indian Veterinary Institute, Izatnagar, Bareilly, Uttar Pradesh, India.
- Bacci, M.L., G. Galeati, M. Mattoli, R. Boni and E. Seren. 1991. *In vitro* maturation and *in vitro* fertilization of buffalo oocytes, p. 97. *In Proceedings of 3rd World Buffalo Congress*, Varna, Bulgaria.
- Ball, G.D., M.L. Feibfried, R.W. Lenz, R.L. Ax, B.D. Bavister and N.L. First. 1983. Factors affecting successful *in vitro* fertilization of bovine follicular oocytes. *Biol. Reprod.*, **28**(3): 717-725.
- Bavister, B.D., T.A. Rose-Hellenkant and T. Pinyo-

- pomint. 1992. Development of *in vitro* matured, *in vitro* fertilized bovine embryos into morulae and blastocysts in defined medium. *Theriogenology*, **37**: 124-146.
- Brackett, B.G., A.I. Younis and R.A.F. Hosken. 1989. Enhanced viability after *in vitro* fertilization of bovine oocytes matured *in vitro* with high concentration of lutenizing hormone. *Fertil. Steril.*, **51**: 319-4324.
- Chauhan, M.S., P. Palta, S. Das, P.K. Katoyar and M.L. Madan. 1997a. Replacement of serum and hormone additives with follicular fluid in the IVM medium: effects on maturation, fertilization and subsequent development of buffalo oocytes *in vitro*. *Theriogenology*, **48**: 461-469.
- Cox, J.F., J. Hormazabal and A. Santa Maria. 1993. Effect of cumulus on *in vitro* fertilization of bovine matured oocytes. *Theriogenology*, **40**: 1259-1267.
- De Felici, M. and G. Siracusa. 1982. "Spontaneous" hardening of the zona pellucida of mouse oocytes during *in vitro* culture. *Gamete Res.*, **6**: 107-113.
- Fukuda, Y., M. Ichikawa, K. Naito and Y. Toyada. 1990. Birth of normal calves resulting from bovine oocytes matured, fertilized and cultured with cumulus cells *in vitro* up to blastocyst stage. *Biol. Reprod.*, **22**: 669.
- Handrow, R.R., R.W. Lenz and R.L. Ax. 1982. Structural comparisons among glycosaminoglycans reaction to promote an acrosome in bovine spermatozoa. *Biochem. Biophys. Res. Co.*, **107**: 1326-1332.
- Kobayashi, K., S. Yamashita and H. Hoshi. 1994. Influence of epidermal growth factor and transforming growth factor- α on *in vitro* maturation of cumulus cell-enclosed bovine oocytes in a defined medium. *J. Reprod. Fertil.*, **100**: 349-446.
- Lu, G. and T.T. Hsu. 1991. *In vitro* fertilization of water buffalo. p. 100. *In Proceedings of 3rd World Buffalo Congress*, Varna, Bulgaria.
- Madan, M.L., M.S. Chauhan, S.K. Singla and R.S. Manik. 1994. Pregnancies established from water buffalo (*Bubalus bubalis*) blastocysts derived from *in vitro* matured *in vitro* fertilized oocytes and co-cultured with cumulus cells. *Theriogenology*, **57**(3): 1095-1103.
- Nandi, S., M. Ravindranatha, P.S.P. Gupta, H.M. Raghu and P.V. Sarma. 2003. Development competence and post-thaw survivability of buffalo embryos produced *in vitro*: effect of growth factors in oocytes maturation medium and embryo culture system. *Theriogenology*, **60**(9): 1621-1631.
- Schroeder, A.C., A.K. Champlin, L.E. Morbraaten and J.J. Eppig. 1990. Development capacity of mouse oocytes cryopreserved before and after maturation *in vitro*. *J. Reprod. Fertil.*, **89**: 43-50.
- Sirard, M.A. and N.L. First. 1988. *In vitro* inhibition of oocytes nuclear maturation in bovine. *Biol. Reprod.*, **39**: 229-234.
- Singh, G., S.M. Totey and G.P. Talwar. 1989. *In vitro* fertilization of oocytes matured *in vitro*. *Theriogenology*, **31**: 255.
- Totey, S.M., C.H. Pawshe and G.P. Singh. 1993. *In vitro* maturation and fertilization of buffalo oocytes (*Bubalus bubalis*): Effect of media, hormone and sera. *Theriogenology*, **39**: 1153-1171.
- Vanderhyden, B.C. and D.T. Armstrong. 1989. Role of cumulus cells and serum on the *in vitro* maturation, fertilization, and subsequent development of rat oocytes. *Biol. Reprod.*, **40**: 720-728.