

FACTORS INFLUENCING THE DRY PERIOD AND CALVING INTERVAL IN DIFFERENT GRADES OF BUFFALOES

Shashi Sanker, Dharendra Kumar*, K.G. Mandal, R.K. Taggar and A.K. Das

ABSTRACT

Data on 920 buffaloes were analyzed for dry period and calving interval. The buffaloes comprised three genetic groups, viz., Graded Murrah, *Diara* buffaloes and non-descript buffaloes. Buffaloes were enumerated from 145 dairy units located in and around Patna Bihar. Genetic Groups had shown highly significant ($p < 0.01$) influence on both the traits. The dry period varied as 130 (days) in Graded Murrah, 151 (days) in *Diara* buffalo and 150 (days) in non-descript ones. Farming system and location did not significantly influence these traits. Lactation order had highly significant effect on these traits. The dry period varied from 157 days in first lactation to 140 days in fourth lactation while calving interval varied from 466 days in the first lactation to 441 days in the fourth lactation.

Keywords: dry period, calving interval, buffaloes, genetic effect, non genetic effect

INTRODUCTION

The dry period is a directly observed economic trait of very high practical significance in dairy farming. Calving interval is the indicator of sound reproductive status of milch animals. A period of 12-13 months has been recommended as

an ideal calving interval in cows and buffaloes. A milch animal is supposed to be economical if she has shorter dry period and lower calving interval. Thus the dry period and the calving interval are the important economic traits that determine the milk production efficiency of buffaloes. Considering the role of these two traits, the present investigation was carried out to study the effect of various genetic and non-genetic factors on these two traits.

MATERIALS AND METHODS

The present study was conducted on altogether 920 buffaloes consisting of 331 Graded Murrah, 221 *Diara* and 368 non-descript buffaloes which were enumerated from 145 dairy units located in and around Patna district of Bihar, India. The different genetic groups (i) *Punjabiya* (Graded Murrah) having Murrah germplasm in their ancestry, spiral orientation of the horn, relatively less clearance of the body from the ground, small and typical face cut and jet black skin colour. (ii) *Deshila* (*Diara*) true breeding population evolved in the Tal and Diara areas of the North and South Gangatic plain of Bihar around Patna and (iii) non-descript (other than above two types). The whole area under study was divided into three distinct zones considering geographical attributes, viz., Zone I: North West Patna, Zone II: South West

Division of Animal Breeding and Genetics, Bihar Veterinary College, Patna, Bihar 800-014, India, *E-mail: dr.dhirendrakumar@yahoo.com

Patna, Zone III: East Patna. The enumerated dairy units were grouped according to the farming system adopted by the farmers which are mixed farming (animal husbandry integrated with agriculture) and only animal husbandry practices. Records on dry period (days) and calving interval(days) of the buffaloes were classified into four groups on the basis of sequence of lactation up to fourth parity. The data were subjected to least squares analysis (Harvey, 1975). The Duncan Multiple Range Test as modified by Kramer (1957) was used to examine the pair wise comparison among least squares means.

RESULT AND DISCUSSION

Dry period

The overall least squares mean was found to be 144.34 ± 0.77 days (Table 1) for the dry period in the three genetic groups of buffaloes which was higher than the optimum range. The reason for this could be the milking of their animals by the private *khatal* [cattle shed] owners in late gestation and the indiscriminate use of oxytocin for letting down of milk, which interfere with the subsequent pregnancy of the animals. Similarly longer dry periods were reported by Rao *et al.* (1995), Sethi (1996-97) and Yadav *et al.* (2003) in Murrah, by Pathodiya *et al.* (1998) and Yadav (1995-96) in Surti, and by Paliwal (1994) in Mehsana and Chawla (1996-97) in Nili-Ravi buffaloes. Shorter dry periods have been reported by Dev Raj and Gupta (1994) in local buffaloes of Rajasthan and by Sinha (2006) in buffaloes consisting of three different genetic groups in and around Barh (Bihar). Genetic group and order of lactation had a significant ($P < 0.01$) while location and farming system had a non-significant influence on dry period (Table 1).

Genetic group: The least squares means of Graded Murrah had the shortest dry period (130.48 ± 1.32 days) which was significantly ($P < 0.05$) lower than the Diara and non-descript buffaloes by 21.12 and 20.45 days, respectively (Table2). Diara buffaloes though had the longest dry period (151.60 ± 1.30 days) but did not differ significantly from the non-descript type. Although the dry period is supposed to be influenced by non-genetic causes, in the present study, the genetic divergence between Graded Murrah, Diara and non-descript buffaloes might have resulted in expression of a genetic effect on dry period that was significant. Similarly, Sinha (2006) observed the effect of genetic constitution on dry period in buffaloes to be significant. Much variation in the dry period reported by various workers (Siddiquee *et al.*, 1984, Singh, 1992, Dev Raj and Gupta, 1994, Pathodiya *et al.*, 1998 and Kumar, 2004) for different breeds of buffaloes may be indicative that genetic constitution of the animals might be a considerable factor for variation in dry period. Contrary to the findings of the present study, Priya Raj (2002) did not record a significant effect of genetic group on the dry period in cows.

Location: The animals located in different zones did not differ significantly among themselves with respect to their dry periods and the effect of location contributed only 0.56% to the total variation in this trait (Table 1). However, the least squares mean for average dry period was the longest (145.09 ± 1.40 days) for the animals located in the *khatal*s in East Patna followed by those located in North West (144.85 ± 1.25 days) and South West (143.07 ± 1.29 days) zones (Table2). Srivastava *et al.* (1998), Rao *et al.* (2000), Priya Raj (2002) and Kumar (2004) also reported this effect to be non-significant in cows and buffaloes under the private sector.

Farming System: The farming system did not have significant effect on dry period (Table 1). As evident from Table 2, the animals managed in both the units had almost the same duration of dry periods. The average estimates of dry period in the units involved in dairy farming alone and those maintained in the units integrated with agriculture farming were 144.58 ± 1.11 and 144.10 ± 1.05 days, respectively.

Parity: Parity had a highly significant ($P < 0.01$) effect on dry period (Table 1). The average dry period was found to be the longest (157.85 ± 1.39 days) in first calvers followed by second (143.57 ± 1.33 days) and third (135.65 ± 1.50 days) calvers (Table 2). The average dry periods in second, third and fourth calvers were found decrease significantly ($P < 0.05$) by 14.28, 22.20 and 17.58 days, respectively, compared to the first calvers. The average dry period of fourth calvers was reckoned to be increased by 4.62 days over the third calvers but did not differ significantly. Singh (1992) also observed a similar trend to that of the present study. Contrary to the report of Singh (1992), a significant effect of parity was reported by Kumar (2004) in cows and buffaloes in private dairy units, but he did not find a definite trend as observed in the present investigation.

Calving interval

The overall least squares mean for calving interval in buffaloes of three genetic groups *viz.* Graded Murrah, Diara and non-descript types, in and around Patna, was observed to be 450.24 ± 1.53 days (Table 2), which was longer than the optimum range desirable for profitable milk production but close to the estimates reported by Dev Raj and Gupta (1994), Kumar (2004) and Sinha (2006). Higher estimates of calving interval in comparison to the findings of the present study have been

reported by Johari and Bhat (1979), Rao *et al.* (1995) and Yadav *et al.* (2003) in Murrah buffaloes and Siddiquee *et al.* (1984) and Singh (1992) in Mehsana buffalo. Least squares analysis of variance (Table 1) revealed that genetic constitution of the animals and order of lactation had significant ($P < 0.01$) effects on calving interval. The effects of zone and farming system were not significant statistically.

Genetic group: Genetic group had a highly significant ($P < 0.01$) influence on calving interval, and its contribution to the total variation in calving interval was the highest: 77.61% (Table 1). As evident from Table 2, the Graded Murrah had the shortest calving interval (424.32 ± 2.60 days), which was significantly ($P < 0.05$) less than the estimates of Diara and non-descript buffaloes by 39.89 and 37.87 days, respectively. Diara buffaloes had the longest inter calving period (464.21 ± 2.57 days) but did not differ significantly from the non-descript types. Calving interval is supposed to be influenced by non-genetic causes, but in the present investigation, the genetic divergence between Graded Murrah, Diara and non-descript buffaloes might have resulted into expression of a genetic effect that was significant. Singh *et al.* (2000) and Kumar (2004) also reported the effect of genetic group on calving interval to be statistically significant in the case of crossbred cows and buffaloes in private dairy units. A significant effect of genetic group on calving interval in buffaloes maintained in private dairy units was also reported by Sinha (2006), but the longest and shorter calving intervals, respectively, in non-descript and Graded Murrah reported by him were contrary to the findings of the present study.

Location: The animals in different locations did not differ significantly with respect to their calving interval, and the contribution of

Table 1. Least squares means±SE and CV% of dry period (days) and calving interval (days) of buffaloes in and around Patna.

| Particulars | Dry period (days) Mean ± S.E. | Calving interval (days) Mean ± S.E. |
|---------------------------|--|--|
| Overall Mean (μ) | 144.34±0.77 (9.34) | 450.24±1.53 (5.95) |
| Factors | | |
| Genetic group | ** | ** |
| Graded Murrah | 130.48 ^a ±1.32 (10.11) | 424.32 ^a ±2.60 (6.12) |
| Diara | 151.60 ^b ±1.30 (8.66) | 464.21 ^b ±2.57 (5.59) |
| Non descript | 150.93 ^b ±1.30 (8.82) | 462.19 ^b ±2.55 (5.65) |
| Location | NS | NS |
| 1. North West Patna | 144.85±1.25 (9.13) | 452.17±2.46 (5.75) |
| 2. South West Patna | 143.07±1.29 (9.28) | 448.42±2.55 (5.85) |
| 3. East Patna | 145.09±1.40 (9.10) | 450.13±2.75 (5.76) |
| Farming System | NS | NS |
| 1. Animal husbandry alone | 144.10±1.05 (9.21) | 448.34±2.07 (5.84) |
| 2. Mixed farming | 144.58±1.11 (9.31) | 452.14±2.18 (5.84) |
| Lactation order | ** | ** |
| 1 st | 157.85 ^a ±1.39 (8.30) | 466.70 ^a ±2.74 (5.53) |
| 2 nd | 143.57 ^b ±1.33 (9.07) | 453.98 ^b ±2.61 (5.63) |
| 3 rd | 135.65 ^c ±1.50 (9.64) | 438.56 ^c ±2.96 (5.88) |
| 4 th | 140.27 ^{bc} ±1.93 (9.33) | 441.72 ^c ±3.80 (5.83) |

Means with different superscripts (column-wise) differed significantly ($P < 0.05$).

Values in parentheses are CV%.




Figure 1. Changes in dry periods (days) due to different factors.




Figure 2. Changes in calving intervals (days) due to different factors.

location to the total variation for this trait was only 0.57% (Table2). The animals located in North-West zone of Patna had the longest inter calving period (452.17 ± 2.46 days) followed by those animals located in the East (450.13 ± 2.75 days) and the South West (448.50 ± 2.55 days) zones. Srivastava *et al.* (1988), Rao *et al.* (2000), Priya Raj (2002) and Kumar (2004) also reported this effect to be non-significant in cows and buffaloes maintained in private dairy units.

Farming system: The farming system did not have significant influence on calving interval; the contribution of the farming system to the total variation was reckoned to be 1.66% (Table2). As evident from Table19, the animals maintained in the units involved dairy farming alone exhibited a lower calving interval (448.34 ± 2.07 day) than the animals managed in the units integrated with agriculture farming (452.14 ± 2.18 days). However, the animals maintained under the two different farming systems did not differ significantly with respect to this trait. Johari and Bhatt (1979) reported highly significant effects of farms and periods on calving interval in buffaloes.

Parity: As evident from Table 1, the influence of order of lactation on calving interval was highly significant ($P < 0.01$); its contribution to the total variation for this trait was next to the effect of genetic group, which was accounted to be 19.16%. A definite trend was observed in the variation of calving interval from parity to parity. The average first calving interval was found to be longest (466.70 ± 2.74 days) followed by second (453.98 ± 2.61 days) and third (438.56 ± 2.96 days) calving intervals (Table-2). The average estimates of calving interval in second and third parities were found to decrease significantly ($P < 0.05$) by 12.72 and 28.14 days, respectively, compared to the first calving interval. The fourth calving interval was

found to be increased by 13.16 days over that of the third parity, but the two did not differ significantly. A significant effect of parity on calving interval was also reported by Singh (1992) and Kumar (2004) in cows and buffaloes maintained in private dairy units. The trend of decreasing calving interval from first to third parity as observed in the present study was also reported by Singh (1992). However, Siddiquie *et al.* (1984), Tailor and Jain (1986) and Rahejha (1992) reported non-significant effects of parity on calving interval.

Diara: Buffaloes though had significantly ($P < 0.05$) higher dry period and calving interval in comparison to Graded Murrah but non significantly higher fry period and calving interval than non-descript types. There were non-significant effects of location and farming system observed in this study. Therefore, work of a similar type may be repeated in the entire Tal and Diara areas of the river Ganges, Gandak and Sone pertaining to Bihar(India) to identify and enumerate the number and performance status of *Diara* buffaloes so that a suitable breeding plan can be chalked out for improvement of *Diara* buffaloes.

REFERENCES

- Dev, Raj and J.N. Gupta. 1994. An economic analysis of milk production in Churu district in Rajasthan. *Indian J. Dairy Sci.*, **47**(4): 294-301.
- Gupta, B.D., S.N. Kaushik and R.R. Mishra. 1994. Study on reproduction efficiency parameter of Murrah buffalo. *Indian J. Dairy Sci.*, **47**: 4.
- Harvey, W.R. 1975. Least Squares Analysis of Data with Unequal Subclass Number. United State Department of Agriculture (USDA).

- Johari, D.C. and P.N. Bhat. 1979b. Effect of genetic and non-genetic factors on reproductive traits in Indian buffaloes. *Indian J. Anim. Sci.*, **49**(1): 1-6.
- Kramer, C.Y. 1957. Extension of Multiple Range Test to group correlated adjusted means. *Biometrics*, **13**: 13-17.
- Kumar, N. 2004. *Genetic analysis of milk production efficiency in cattle and buffaloes in and around Dharbhanga (Bihar)*. M.V.Sc. Thesis, RAU, Pusa (Samastipur), Bihar, India.
- Paliwal, P.C. 1994. *Genetic and economic investigations on the productivity of medium sized (Surti) buffaloes*. Ph.D. Thesis, R.B.S. College, Bichpuri, Agra (U.P.), India.
- Pathodiya, O.P., L.S. Jain, S.P. Tailor and Bechcha Singh. 1998. Effect of period of calving on different traits in Surti buffalo. *Indian J. Dairy Sci.*, **51**(5): 280-284.
- Priya, Raj. 2002. *Studies on milk production and its economics in crossbred cows under farmer's managerial conditions of Patna (Bihar)*. M.V.Sc. Thesis, Rajendra Agricultural University, Pusa, Bihar, India.
- Raheja, K.L. 1992. Selection free estimates of genetic parameters of production and reproduction traits of first three lactation in Murrah buffaloes. *Indian J. Anim. Sci.*, **62**(2): 149-154.
- Rao, S.J., B.V.R. Rao and G.N. Rao. 2000. Performance of crossbred cows and buffaloes under village conditions of Vishakhapatnam district of Andhra Pradesh. *Indian J. Dairy Sc.*, **53**(3): 222-226.
- Rao, B.D. and C.B. Singh. 1995. Impact of Operation Flood on economics of the buffalo milk production in Guntur district. Anadhra Pradesh. *Indian Dairyman*, **7**(4): 47-50.
- Sethi, R.K. 1996-1997. Production performance of Murrah buffaloes. Annual Report, 1996-1997, CIRB, Hisar, India.
- Shrivastava, A.K., C.S.P. Singh and S.K. Verma. 1998. A study on the effects of various factors on production traits in Zebu x Friesian crossbred cows maintained under un-organized farming system. *Indian Vet. Med. J.*, **22**(1): 19-22.
- Siddiquee, G.M., R.K. Tajane and M.B. Pande. 1984. Effect of year and parity on some of the reproductive traits in Mehsana buffaloes. *Livestock Advisor*, **9**(10): 23-25.
- Singh, D.V. 1992. *Breed characterization of Mehsana buffaloes and strategies of their genetic improvement*. Ph.D. Thesis, NDRI, Deemed University, Karnal, India.
- Sinha, R.K. 2006. *Characterization of buffalo genetic resources in Tal and Diara areas in and around Barh (Patna)*. M.V. Sc. Thesis, RAU, Pusa (Samastipur), Bihar, India.
- Yadav, B.S., M.C. Yadav, A. Singh and F.H. Khan. 2003b. Factors affecting calving interval and dry period in Murrah buffalo. *Indian Vet. Med. J.*, **27**: 145-146.
- Yadav, S.B.S. 1995-1996. Network Project on Surti buffaloes, Vallabhnagar Unit. Annual Report, Livestock Research Station, Vallabhnagar, RAU, Bikaner, India.