

เอกสารอ้างอิง

- ประศาสตร์ เกื้อมณี. 2538. เทคนิคการเพาะเลี้ยงเนื้อเยื่อพืช, 158 หน้า. กรุงเทพฯ :
ภาควิชาชีววิทยา คณะวิทยาศาสตร์ มหาวิทยาลัยเกษตรศาสตร์.
- สถาบันวิจัยยาง กรมวิชาการเกษตร กระทรวงเกษตรและสหกรณ์ (2544). คำแนะนำ
พันธุ์ยางปี 2544. หน้า 1-30.
- สมปอง เตชะโต และอรุณี ม่วงแก้วงาม. (2535). การเพาะเลี้ยงเนื้อเยื่อยางพารา I การ
ขยายพันธุ์ยางในหลอดทดลอง. ว.สงขลานครินทร์ 14 : 123-132.
- Bradford, M. M. (1976). A rapid and sensitive method for quantitation of
microgram quantities of protein utilizing the principal of protein dye-
binding. Anal. Biochem., 72, 246-254.
- Breton, F., Sanier, C. and d'Auzac, J. (1997). Scopoletin degradation in relation
to resistance of *Hevea brasiliensis* to *Corynespora cassiicola*. J. Plant
Physiol., 151, 595-602.
- Burner R. L. (1964). Determination of reducing sugar value 3,5-dinitrosalicylic
acid method. Methods Carbohydr. Chem., 4, 67-71.
- Chirapongsatonkul, N., Nakudom, P. and Churngchow, N. (2008). Defense
responses of calli and seeds of *Hevea brasiliensis* to zoospores and the
elicitor of *Phytophthora palmivora*. J. of Phytopathol., 156, 732-741.
- Churngchow, N., Suntaro, A. and Wititsuwannakul, R. (1995). β -1,3-glucanase
isozymes from the latex of *Hevea brasiliensis*. Phytochemistry, 39, 505-
509.
- Churngchow, N. and Rattarasarn, M. (2000). The elicitor secreted by
Phytophthora palmivora, a rubber tree pathogen. Phytochemistry, 54,
33-38.

- Churngchow, N. and Rattarasarn M. (2001). Biosynthesis of scopoletin in *Hevea brasiliensis* leaves inoculated with *Phytophthora palmivora*. J. Plant Physiol., 158, 875- 882.
- Darvill, G. A. and Albersheim P. (1984). Phytoalexins and their elicitors - a defense against microbial infection in plants. Ann. Rev. Plant Physiol., 35, 243-275.
- De Leo, F., Volpicella, M., Licciulli, F., Liuni, S. Gallerani, R. and Ceci L. R. (2002). PLANT-Pis : a data base for plant protease inhibitors and their genes. Nucl. Acids Res., 30 (1), 347-348.
- Edwards, R., Stones, T. M., Gutierrez-Mellado, M. C. and Jorin, J. (1997). Characterization and inducibility of a scopoletin degrading enzyme from sunflower. Phytochemistry, 45, 1109-1114.
- Friend, J., Reynolds, S. B., Aveyard, M. A. (1973). Phenylalanine ammonia lyase, chlorogenic acid and lignin in potato tuber tissue inoculated with *Phytophthora infestans*. Physiol. Plant Pathol., 3, 495-507.
- Giesemann, A., Biehl, B. and Lieberei, R. (1986). Identification of scopoletin as a phytoalexin of the rubber-tree *Hevea brasiliensis*. J. Phytopathol., 117, 373-376.
- Guest, D. and Brown, J. (1997). Infection process, epidemiology and crop-loss assessment. Plant pathogens and Diseases, 246-286. Brown, J.F. and Ogle, H.J. Australia. Rockvale Publications.
- Hermann, S. (2006). Protocol Tricine-SDS-PAGE. Nature Protoc., 1(1), 16-23.
- Hermosa, M. R., Turra, D., Fogliano, V., Monte, E. and Lorito, M. (2006). Identification and characterization of potato protease inhibitors able to inhibit pathogenicity and growth of *Botrytis cinerea*. Physiol. and Mol. Plant Pathol., 68, 138-148.
- Henninger, H. Zur Kultur von. 1963. *Phytophthora infestans* auf vollsynthetischen Nahrs substraten. Z Allg Mikrobiol., 3, 126-135.

- Higuchi, T. (1985). Biosynthesis of lignin. P. 141-160, In : Biosynthesis and biodegradation of wood components. T. Higuchi, ed. Academic Press. New York.
- Jouili, H., El Ferjani, E. (2003). Changes in antioxidant and lignifying enzyme activities in sunflower roots (*Helianthus annuus* L.) stressed with copper excess. Plant Biol. and Pathol., 326 : 639–644.
- Kuc', J. (1995). Phytoalexin, stress metabolism, and disease resistant in plants. Ann. Rev. Phytopathol., 33, 275-291.
- Mallikarjuna Rao, N., Nayana Rao, H. and Pattabiraman, T. N. (1983). Enzyme inhibitors from plants. Isolation and characterization of a protease inhibitor from arrow root (*Maranta arundinaceae*) tuber. J. Biosci., 5, 21-33.
- Martina, R., Joelle, F., and Marie ,T. T. (1989). Induction of protease Inhibitors in tobacco cell suspension culture by elicitors of *Phytophthora parasitica* var. *nicotianael*. Plant Physiol., 90 :1065-1070.
- Milat, M-L, Ricci, P.,Bonnet, P., Blein J-P. (1990). Capsidiol and ethylene production by tobacco cells in response to cryptogein, an elicitor from *Phytophthora cryptogea*. Phytochemistry, 30, 2171-2173
- Mónica P., Emilia L-S., Rosa, S-M., Gabriel, S. and Pablo, R-P. (1999). Antifungal activity of a plant cystatin. Molec. Plant-Microb. Interact., 12, 624-627.
- Nespoulous, C., Huet J. C. and Pernollet, J. C. (1992). Structure-function relationships of α and β elicitin, signal proteins involved in the plant-*Phytophthora* interaction. Planta, 186, 551-557.
- Perez, F.J., Villegas, D. and Mejia, N. (2002). Ascorbic acid and flavonoid-peroxidase reaction as a detoxifying system of H_2O_2 in grapevine leavea. Phytochemistry, 60, 573-580.



- Tan, A. M. and Low, F. C. (1975). Phytoalexin production by *Hevea brasiliensis* in response to infection by *Colletotrichum gloeosporioides* and its effect on other fungi. In : Proceeding International Rubber Conference, 1975. Kuala Lumpur, Malaysia, 217-227.
- Thomas, E. C. and Lowell L. B. (1982). Partial purification of proteinase inhibitors from wounded tomato plants. *Plant Physiol.*, 69 (2), 537-542.
- Rattarasarn, M. (2003). Defense responses of *Hevea brasiliensis* against zoospores and elicitor from *Phytophthora palmivora*. Doctor of Philosophy thesis in Biochemistry. Prince of Songkla University. Thailand.
- Ricci, P., Bonnet, P., Huet J. C., Sallantin, M., Beauvais-Cante, F., Brnetaeu, M., Billard, V., Michel, G. and Pernollet, J. C. (1989). Structure and activity of proteins from pathogenic fungi *Phytophthora* eliciting necrosis and acquired resistance in tobacco. *Eur. J. Biochem.*, 183, 555-563.
- Seidl, D.S., Hugo, A. and Jaffe, W.G. (1978) Purification of a subtilisin inhibitor from black bean seeds. *FEBS Lett.*, 92 (2) : 245-250.
- Silva, M. C., Nicole , M. Guerra-Guimaraes, L. and Rodrigues Jr, C. J. (2002 a). Hypersensitive cell death and post-haustorial defense response arrest the orange rust (*Hemileia vastatrix*) growth in resistant coffee leaves. *Physiol. Mol. Plant Pathol.*, 60, 169-183.
- Sritanyarat, W., Pearce, G., Siems, W. F., Ryan, A. C., Wititsuwannakul, R. and Wititsuwannakul, D. (2006). Isolation and characterization of iso inhibitors of the potato protease inhibitor I family from the latex of the rubber tree, *Hevea brasiliensis*. *Phytochemistry*, 67, 1644-1650.
- Valueva, T.A. and Mosolov, V.V. (2004). Role of inhibitors of proteolytic enzymes in plant defense against Phytopathogenic microorganisms. *Biochemistry (Moscow)*, 69 (11) : 1305-1309.

- Van Loon, L. C. and Van Strien, E. A. (1999). The families of pathogenesis-related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiol. and Mol. Plant Pathol.*, 55, 85-97.
- Walsh, T. A. and Twitchell, W. P. (1991). Two Kunitz-type proteinase inhibitors from potato tubers. *Plant Physiol.*, 97, 15-18.
- Wang, S., Lin, J., Ye, M., Bun, N.T., Rao, P., Ye, X. (2006). Isolation and characterization of a novel mung bean protease inhibitor with antipathogenic and anti-proliferative activities. *Peptides*, 27 : 3129 – 3136.
- Wang, Y.C., Hu, D.W., Zhang, Z.G., Ma, Z.C., Zheng, X.B. and Li, D.B.. (2003). Purification and immunocytolocalization of a novel *Phytophthora boehmeriae* protein inducing the hypersensitive response and systemic acquired resistance in tobacco and Chinese cabbage. *Physiol. and Mol. Plant Pathol.*, 63, 223-232.



