

เอกสารอ้างอิง

- Alderet, J.E., Karl, D.W., and Park, C.H. (1993) Production of poly(hydroxybutyrate), homopolymer and copolymer from ethanol and propanol in a fed-batch culture. *Biotechnol. Prog.* 9: 520-525.
- Anderson, A.J. and Dawes, A.E. (1990) Occurrence, Metabolism, Metabolic Role, and Industrial Uses of Bacterial Polyhydroxyalkanoates. *Microbiology Reviews.* 54:450-470.
- Chen, Guo-Qiang and Page, J.W. (1994) The effect of substrate on the molecular weight of poly-hydroxybutyrate produced by *Azotobacter vinelandii* UWD. *Biotechnology Letters.* 16: 155-160.
- Curtin, D. and Smillie, W.G. 1983. Soil Solution Composition as Affected by Liming and Incubation. *Soil Science Society of America Journal.* 47: 701-707.
- Filipovic, S.S., Ristic, D.M. and Sakac, B.M. 2002. Technology of corn steep application in animal mashes and their quality. *Roum. Biotechnol. Lett.* 7: 705-710.
- Gogolewski, S., Jovanovic, M., Perren, S.M., Dillon, J.G., Hughes, M.K., 1993. Tissue Response and in vivo Degradation of Select Polyhydroxyacid: Polylactides (PLA), Poly(3-hydroxybutyrate) (PHB), and Poly(3-hydroxybutyrate-co-3-hydroxyvalerate) (PHB/VA). *J. Biomed. Mater. Res.* 27, 1135-1148.
- Hahn, K.S., Chang, K.Y. and Lee, Y.S. 1995. Recovery and characterization of poly(3-Hydroxybutyric acid) synthesized in *Alcaligenes eutrophus* and recombinant *Escherichia coli*. *Applied and Environmental Microbiology.* 61: 34-39.
- Jianchun Li, Bo Zhu, Yong He, Yoshio Inoue, 2003. Thermal and Infrared Spectroscopic Studies on Hydrogen-Bonding Interaction between Poly(3-hydroxybutyrate) and Catechin. *Polym. J.* 35, 384.
- Lee, S.Y., Chang, H.N., 1995. Production of poly(3-hydroxyalkanoates). *Adv. Biochem. Eng. Biotechnol.* 52, 27-58.
- Mona, K. Gouda, Azza, E. Swellam, Sanaa, H. Omar, 2001. Production of PHB by a *Bacillus megaterium* strain using sugarcane molasses and corn steep liquor as sole carbon and nitrogen source, *Microbiological research.* 156, 201-207.
- Page, W.J., 1992 a. Production of polyhydroxyalkanoates by *Azotobacter vinelandii* UWD in beet molasses culture. *FEMS Microbiol. Rev.* 103, 149-158.

- Page, W.J., 1992 b. Suitability of commercial beet molasses fractions as substrates for polyhydroxyalkanoate production by *Azotobacter vinelandii* UWD. *Biotechnol. Lett.* 14, 385-390.
- Ramsay, B.A., Lomaliza, K., Chavarie, C., Dubé, B., Bataille, P., Ramsay, J.A. (1990) Production of poly-(β -hydroxybutyric-Co- β -Hydroxyvaleric) acids. *Appl. Environ. Microbiol.* 56: 2093-2098.
- Rehm, A.H. Bernd. 2003. Polyester synthases: natural catalysts for plastics. *Biochem J.* 376: 15-33.
- Steinbüchel, A., Hustede, E., Liebergesell, M., Pieper, U., Timm, A., and Valentin, H. (1992) Molecular basis for biosynthesis and accumulation of polyhydroxyalkanoic acids in bacteria. *FEMS Microbiol. Rev.* 103: 217-230.
- Williams, F.S. and Martin, P.D. 1996. Application of PHAs in Medicine and Pharmacy. Tephra, Inc. 303 Third Street, Cambridge.
- Sei Kwang Hahn et al., Yong Keun Chang,* and Sang Yup Lee, 1994)