

เอกสารอ้างอิง

จิระเดช แจ่มสว่าง จินตนา แซนนະ วรร矜วีໄລ เกษณรา เฉลิมลาก ช่วยประสิทธิ์ สุพรณี ชีวาริยะกุล ธีรยุทธ ตู้จินดา ศรปราษฐ์ รainer ศวรรยางกุล วุฒิชัย ญาณอรรถ กัทลีวัลย์ สุขช่วย และ สำนักกายาพาด. 2535. การควบคุมโรคต้นแห้งของข้าวบาร์เลย์โดยวิธีคลุกเมล็ดด้วยมวลปีวภาพของเชื้อรา *Trichoderma hazianum*. สารสารข่าวศูนย์ปฏิบัติการวิจัยและเรือนปลูกพืชทดลอง มหาวิทยาลัยเกษตรศาสตร์. 6: 4-8.

จิระเดช แจ่มสว่าง จินตนา ฉะนະ ปราโมทย์ ศิริโรจน์ กณิษฐา สังกกะหะ วิชชุพร ว่องสุวรรณเลิศ และวรรณวีไล เกษนรา. 2536. ประสิทธิภาพของเชื้อราไตรโคเดอร์มา ชาเขียวน้ำ พันธุ์กล้ายที่ต้านทานต่อบนโน้มิลในการยับยั้งการเจริญของเชื้อราสเคลอโรเรียม รอล์ฟส์ไอ. n.76 ในบทคัดย่อของการประชุมวิชาการครั้งที่ 5 ของสมาคมเทคโนโลยีชีวภาพแห่งประเทศไทย เรื่องเทคโนโลยีชีวภาพเพื่อพัฒนาคุณภาพชีวิต. 25-27 พฤษภาคม 2536. โรงแรมไฮลตัน อินเตอร์เนชันแนล กรุงเทพฯ.

พรพรรณ พรหมเสถียร. 2543. ผลของสาร secondary metabolite ของเชื้อรา *Trichoderma* spp. ต่อเชื้อรา *Sclerotium rolfsii* Sacc. สาเหตุโรคโคนเน่าของมะเขือเทศ. วิทยานิพนธ์วิทยาศาสตรมหาบัณฑิต มหาวิทยาลัยขอนแก่น.

วีระศักดิ์ ศักดิ์ศิริรัตน์ และระวีวรรณ ศรีละเอียด. 2529. การศึกษาเชื้อ *Trichoderma* spp. เพื่อใช้ป้องกันโรคโคนเน่าของผักและถั่วลิสงโดยชีววิธี. กำนงเกษตร. 13: 278-282.

สุวิตา แสพศาล. 2549. เอนไซม์ย่อยสลาย ความสัมพันธ์ทางพันธุกรรมของกลีบนิวคลีโอไทด์ ในส่วน ITS1-5.8S-ITS2 ของ rDNA และการโคลนยืนไกดีเนสของเชื้อรา *Trichoderma* spp. วิทยานิพนรปวิจัยสาขาวิชาศาสตรมหาบัณฑิต สาขาวิชาโรคพืชวิทยา บัณฑิตวิทยาลัยมหาวิทยาลัยขอนแก่น.

Atkinson, M.M. 1993. Molecular mechanism of pathogen recognition by plant. Advance Pl. Pathol. 42: 83-95.

Bunyatratchata, W., W. Saksirirat, P. Sirithorn, and P. Teerakupisut. 2005. Race identification of fusarium with pathogen of tomato, *Fusarium oxysporum* f. sp. *lycopersici* by pathogenic reaction on standard differential host and development of Thai differential host. Khon Kaen Agriculture Journal 33, 95-107.

Barker, C., 2000. Systemic acquired resistance. Pp. 198-217 in: Dickinson M. and Beynon J. (ed). Molecular plant pathology. Sheffield Academic Press, Sheffield, U.K.

Bigirimana, J., G. De Meyer, J. Poppe, Y. Elad, and M. Hofte. 1997. Induction of systemic resistance on bean (*Phaseolus vulgaris*) by *Trichoderma harzianum*. Med. Fac. Landbouw, Univ. Gent. 62: 1001-1007.

- Bol, J.F., H.J. M. Linthort, and B.J.C. Cornelissen. 1990. Plant pathogenesis-related protein induce by virus infection. Ann. Rev. Phytopathol. 28:113-138.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem. 72: 248-254.
- Charirak, P., W. Saksirirat, and W. Bunyatratchata. 2008. Effect of antagonistic fungi, *Trichoderma* spp. on enzyme activity induction of β -1,3-glucanase and chitinase derived from tomato leaf. 9th National Grad Research Conference. 14-15 March 2008.
- Darvill, A.G. and P. Albersheim. 1984. Phytoalexin and their elicitors – a defense against microbial infection in plants. Ann. Rev. Pl. Physiol. 35: 243-275.
- Dixon, R. A. and M. J. Harrison. 1992. Activation, structure and organization of genes involved in microbial defense in plants. Advc. Genetic. 28: 165-234.
- Edreva, A., D. Blancard, R. Delon, P. Bonnet and P. Ricci. 2002. Biochemical changes in β -cryptogein-elicited tobacco: a possible of acquire resistance .Beitraege zur Tabakforschung Inter. 20: 53-59.
- Engelberth, J., E.A. Schmelz, H.T. Alborn, Y.J. Cardoza, J. Huang, and J.H. Tumlinson. 2003. Simultaneous quantification of jasmonic acid and salicylic acid in plants by vapor-phase extraction and gas chromatography–chemical ionization–mass spectrometry. Anal. Biochem. 312: 242–250.
- Gaffney, T., L. Friedrich, B. Vernoij, D. Negrotto, G. Nye, S. Uknnes, E. Ward, H. Kessmann, and J. Ryals. 1993. Requirement of salicylic acid for the induction of systemic acquired resistance. Science 261: 754-756.
- Gailite, A., I. Samsone, and G. Ievinsh. 2005. Ethylene is involved in *Trichoderma*-induced resistance of bean plants against *Pseudomonas syringae*. Acta Universitatis Latviensis 691: 59-70.
- Hahlbrock, K., D. Sheel, E. Logemann, T. Nurnberger, M. Parniske, S. Reinold, W. R. Sacks, and E. Schmelzer. 1995. Oligopeptide elicitor-mediated defense gene activation in cultured parsley cells. Anal. Biochem. 122: 252–258.
- Hahn, M. G., P. Buchell, F. Cervone, S. H. Doares, R. A. O'Neil, A. G. Darvill, and P. Albersheim. 1989. Roles of cell wall constituents in plant–pathogen interactions. In: Plant-Microbe Interactions-Molecular and Genetic Perspectives (eds. T Kosung and E.W. Nester), vol. 3, pp. 131-181, McGraw-hill, NY.

- Harman, G.E., C.R. Howell, A. Viterbo, I. Chet, and M. Lorito. 2004. *Trichoderma* species-opportunistic, avirulent plant symbions. *Microbiology* 2: 43-56.
- Heil, M. and R.M. Bostock. 2002. Induce systemic resistance (ISR) against pathogens in the context of induce resistance. *Annal. Botany* 89: 503-512.
- Hennin, C., E. Diederichsen and M. Hofte. 2001. Local and systemic resistance to fungal pathogens triggered by an AVR9-mediated hypersensitive response in tomato and oilseed rape carrying the *cf-9* resistance gene. *Physiol. Molec. Pl. Pathol.* 59: 287-259.
- Hommer, D.W., D.K. Bell, and C.A. Jawoeki. 1972. Efficacy of *Trichoderma harzianum* as a biocontrol for *Sclerotium rolfsii*. *Phytopathology* 73:1043-1046.
- Horst, L.E., J. Locke, C.R. Krause, R.W. McMahon, L.V. Madden, and H.A.J. Hoitink. 2005. Suppression of Botrytis blight of begonia by *Trichoderma hamatum* 382 in peat and compost-amended potting mixes. *Plant Disease* 89 (11): 1195-1200.
- Hurtado, O. 2004. Study and manipulation of the salicylic acid-dependent defense pathway in plants parasitized by *Orobanche aegyptiaca* Pers. Master of Sciences Thesis, Plant Physiology, Virginia Polytechnic Institute and State University, USA.
- Jones, D.A. and D. Takemoto, 2004. Plant innate immunity-direct and indirect recognition of general and specific pathogen-associated molecules. *Curr. Opin. Inmu.* 16:48-62.
- Kaewplung, P. 2001. Seed export, a billion Baht business. *Seed Newsletter* 8: 2-3.
- Khan, J., J.J. Ooka, L.V. Madden, S.A. Miller, and H.A.J. Hoitink, 2004. Systemic resistance induced by *Trichoderma hamatum* 382 in cucumber against Phytophthora crown rot and leaf blight. *Pl. Dis.* 88: 280-286.
- Klement, Z. 1982. Hypersensitivity. In: *Plantpathogenic Prokaryotes*. (eds. M. S. Mount and G. H. Lacy), Volume 2, pp. 149-177, Academic press, NY, USA.
- Kombrink, E. and I. E. Somssich. 1995. Defense response of plant to pathogens. *Ad. Botani. Res.* 21: 1-34.
- Mauch-Mani, B. and J.P. Metraux. 1998. Salicylic acid and systemic acquired resistance to pathogen attack. *Annal. Botany* 82: 535-540.
- Neuenschwander, U., B. Vernooij, L. Friedrich, S. Uknes, H. Kessmann, and J. Ryals. 1995. Is hydrogen peroxide a second messenger of salicylic acid in systemic acquire resistance ? . *Plant J.* 8: 227-233.

- Parker, E.J. 2000. Signalling in plant disease resistance. Pp. 198-217 in: Dickinson M. and Beynon J. (eds). Molecular plant pathology. Sheffield Academic Press, Sheffield, U.K.
- Rifai, M.A. 1969. A revision of the genus *Trichoderma*. Mycol. Papers. C. M. I. 116: 1-56.
- Saksirirat, W. and H.H. Hoppe. 1991. Secretion of extracellular enzyme by *Verticillium psalliotae* Treschow and *Verticillium lecanii* (Zimm,) Viegas during growth on uredospore of the soybean rust fungus (*Phakopsora pachyrhizi* Syd). Inliquid cultures. J. Phytopathol. 131: 161-173.
- Saksirirat, W., N. Boonsakdaporn, P. Sirithorn, and A. Pachinburavan. 1996. An application of the mycoparasite, *Trichoderma harzianum* Raifai. in combination with mancozeb for control tomato stem rot in northeast Thailand. Proceeding of the International Workshop on Biological Control of Plant Diseases. May 22-27, Beijing, China. Pp.327-329.
- Saksirirat, W., M. Chuebandit, P. Sirithorn and N. Sanoamuang. 2005. Species diversity of antagonistic fungus, *Trichoderma* spp. From seed production fields and its potential for control Fusarium wilt of tomato and cucurbits. The 4th International Conference on Biopesticides. Feb.13-18, 2005. Imperial Maeping Hotel, Chiangmai, Thailand.
- Saksirirat, W., P. Chareerak and W. Bunyatrachata. 2009. Induce systemic resistance of biocontrol fungus, *Trichoderma* spp. against bacterial and gray leaf spot in tomatoes. The International Symposium Go..Organic 2009, The Approach of Organic Agriculture: New market, food security and a clean environment, 19-21 August 2009, Pullman Bangkok King Power Hotel, Bangkok, Thailand
- Samuels, G.J. 1996. *Trichoderma* a review of biology and systematics of the genus. Myco. Res. 100: 923-935.
- Shulaev, V., J. Leon, and I. Rakkin. 1995. Is salicylic acid a translocated signal of systemic acquire resistance in tobacco ? Plant Cell 7: 1691-1701.
- Sivan, A.,Y. Elad and I. Chet., 1984. Biological control effect a new isolate of *Trichoderma hazianum* on *Pythium aphanidermatum*. Phytopathology 74: 498-501
- Silverman, F.P., P.D. Pertracek, D.F. Heiman, C.M. Fledderman, and P. Warrior. 2005. Salicylate activity. 3. Structure relationship to systemic acquired resistance. J. Agric. Food Chem. 53: 9775-9780.

- Sirithorn, P. 2001. Seed disease and seed export: case study of bacterial spot of tomato and fruit blotch of watermelon. KKU plant pathology Bulletin. Siripan off-set Khon Kean. p.69-81.
- Sommich, I. E., and K. Hahlbrock. 1998. Pathogen defense in plants-a paradigm of biological complexity. *Plant science* 3: 86-90.
- Somogyi, M. 1952. Notes on sugar determination. *J. Bio. Chem.* 195: 19-23.
- Staskawicz, B.J., F.M. Ausubel, B.J. Baker, J.G. Ellis and J.D.G. Jones. 1995. Molecular-genetics of plant disease resistance. *Science* 268: 661-667.
- Stasz, T.E., G.E. Harman, and N.F. Weeden. 1988. Protoplast preparation and fusion in two biocontrol strains of *Trichoderma harzianum*. *Mycologia* 80:141-150.
- Yang, S.S. and Y. Wang. 1999. Protease and amylase production of *Streptomyces rimosus* in submerged and solid state cultivations. *Bot. Bull. Acad. Sin.* 40: 259-265.
- Yedidia, I., N. Benhamou, and I. Chet. 1999. Induction of defense response in cucumber plants (*Cucumis sativus* L.) by the biocontrol agent *Trichoderma harzianum*. *Applied and Environmental Microbiology*. 65 (3): 1061-1070.
- Yedidia, I., N. Benhamou, Y. Kapulnik, and I. Chet. 2000. Induction and accumulation of PR proteins activity during early stage of root colonization by the mycoparasite *Trichoderma harzianum* strain T-203. *Plant Physiol. Biochem.* 38: 863-873.
- Yedidia, I., M. Shores, Z. Kerem, N. Benhamou, Y. Kapulnik, and I. Chet. 2003. Concomitant induction of systemic resistance to *Pseudomonas syringae* pv. *lycrysans* in cucumber by *Trichoderma asperellum* (T-203) and accumulation of Phytoalexins. *Applied and Environmental Microbiology* 69 (12): 7343-7353.
- Van Loon, L.C. and E.A. Van Strien. 1999. The families of pathogenesis-related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiol. Molec. Pl. Pathol.* 55: 85-97.
- Van Loon, L.C. and A. Van Kammen. 1970. Polyacrylamide disc electrophoresis of the soluble leaf proteins from *Nicotina tabacum* var. "Samaun" and "Samsum NN" II. Changes in protein constitution after infection with tobacco mosaic virus. *Virology* 40: 199-211.
- Van Loon, L.C. and E.A. Van Strien, 1999. The families of pathogenesis-related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiol. Molec. Pl. Pathol.* 55: 85-97.

- Wang, L.J. and H.L. Shao. 2006. Salicylic acid-induced heat or cold tolerance in relation to Ca^{2+} homeostasis and antioxidant systems in young grape plant. Plant Sci. 170(4): 685-694.
- Wasternack, C., I. Stenzel, B. Hause, G. Hause, C. Kutter, H. Maucher, J. Neumerkel, I. Feussner, and O. Miersch. 2006. The wound response in tomato—role of jasmonic acid. J. Pl. Physiol. 163: 297-306.
- Zhou, N., T.L. Tootle, F. Tsui, D.F. Klessig, and J. Glazebrook. 1998. *PAD4* functions upstream from salicylic acid to control defense responses in *Arabidopsis*. Plant Cell 10: 1021-1030.



