

**Full Paper**

## **An analogue of Eulerian polynomials related to $L$ -type function**

**Serkan Araci<sup>1,\*</sup>, Mehmet Acikgoz<sup>2</sup>, Roberto B. Corcino<sup>3</sup> and Cenap Özel<sup>4</sup>**

<sup>1</sup>Department of Economics, Faculty of Economics, Administrative and Social Science, Hasan Kalyoncu University, TR-27410 Gaziantep, Turkey

<sup>2</sup>Department of Mathematics, Faculty of Arts and Science, University of Gaziantep, TR-27310 Gaziantep, Turkey

<sup>3</sup>Department of Mathematics, Cebu Normal University, Cebu City 6000, Philippines

<sup>4</sup>Department of Mathematics, Dokuz Eylül University, TR-35160 Izmir, Turkey

\* Corresponding author, e-mail: [mtsrkn@hotmail.com](mailto:mtsrkn@hotmail.com)

Received: 24 April 2014 / Accepted: 27 July 2015 / Published: 25 August 2015

---

**Abstract:** We introduce Dirichlet's type of twisted Eulerian polynomials by using  $p$ -adic fermionic  $q$ -invariant integral in the  $p$ -adic integer ring and obtain some new interesting identities. Using a complex contour integral representation based on the generating function of Dirichlet's type of twisted Eulerian polynomials, we get  $L$ -type function which interpolates for Dirichlet's type of Eulerian polynomials at negative integers.

**Keywords:**  $p$ -adic fermionic  $q$ -integral on  $Z_p$ , Eulerian polynomials, contour integral,  $L$ -function, Dirichlet character

---

### **INTRODUCTION**

The Eulerian polynomials  $A_n(x)$  are given by means of the following exponential generating series:

$$\sum_{n=0}^{\infty} A_n(x) \frac{t^n}{n!} = e^{A(x)t} = \frac{1-x}{e^{t(1-x)} - x}. \quad (1)$$

In fact, the Eulerian polynomials can be generated via the following recurrence relation:

$$(A(t) + (t-1))^n - tA_n(t) = \begin{cases} 1-t & \text{if } n=0 \\ 0 & \text{if } n \neq 0, \end{cases} \quad (2)$$

where we have used  $A^n(x) := A_n(x)$  symbolically [7, 22].

Let  $p$  be a fixed odd prime number. Throughout this paper we make use of the following

notations:  $Z_p$  denotes the ring of  $p$ -adic rational integers,  $Q$  denotes the field of rational numbers,  $Q_p$  denotes the field of  $p$ -adic rational numbers and  $C_p$  denotes the completion of algebraic closure of  $Q_p$ . Let  $N$  be a set of natural numbers and  $N^* = N \cup \{0\}$ . The normalised  $p$ -adic absolute value is defined by

$$|p|_p = \frac{1}{p}.$$

When one talks of a  $q$ -extension,  $q$  can be variously considered as an indeterminate, a complex number  $q \in C$ , or a  $p$ -adic number  $q \in C_p$ . If  $q \in C$ , one normally assumes  $|q| < 1$ . If  $q \in C_p$ , one normally assumes  $|q-1| < p^{-\frac{1}{p-1}}$ . The  $q$ -integer  $[x]_q$  with  $x$  in  $C$  is also defined by

$$[x]_q = \frac{1-q^x}{1-q} \quad \text{and} \quad [x]_{-q} = \frac{1-(-q)^x}{1+q}.$$

It is easy to see that  $\lim_{q \rightarrow 1} [x]_q = x$  [1-26].

For a positive integer  $d$  with  $(d, p) = 1$ , set

$$X = X_d = \varprojlim_n Z / dp^n Z = \bigcup_{a=0}^{dp-1} (a + dpZ_p)$$

with

$$a + dp^n Z_p = \{x \in X \mid x \equiv a \pmod{dp^n}\},$$

where  $a \in Z$  satisfies the condition  $0 \leq a < dp^n$ , and let  $\sigma : X \rightarrow Z_p$  be the transformation introduced by the inverse limit of the natural transformation:

$$Z / dp^n Z \text{ a } Z / p^n Z.$$

If  $f$  is a function of  $Z_p$ , then we will utilise the same notation to indicate the function  $f \circ \sigma$ .

Let  $UD(Z_p)$  be the space of uniformly differentiable functions on  $Z_p$ . That is, for  $f \in UD(Z_p)$ , the  $p$ -adic  $q$ -integral on  $Z_p$  was originally defined by Kim [11-14] as follows:

$$I_q(f) = \int_X f(v) d\mu_q(v) = \int_{Z_p} f(v) d\mu_q(v) = \lim_{n \rightarrow \infty} \frac{1}{[p^n]_q} \sum_{v=0}^{p^n-1} q^v f(v) \quad (3)$$

The bosonic integral is considered as the bosonic limit  $q \rightarrow 1$ ,  $I_1(f) = \lim_{q \rightarrow 1} I_q(f)$ .

Similarly, the  $p$ -adic fermionic integration on  $Z_p$  was firstly defined by Kim [16, 18] as follows:

$$I_{-q}(f) = \lim_{q \rightarrow -q} I_q(f) = \int_{Z_p} f(v) d\mu_{-q}(v). \quad (4)$$

The above integrals are powerful tools in the studying of  $p$ -adic analogue of some special polynomials such as Bernoulli polynomials, Euler polynomials, Genocchi polynomials, Frobenius-Euler polynomials, Eulerian polynomials, Boole polynomials and their various generalisations [1-33].

By (4), we have the following integral equation:

$$q^n I_{-q}(f_n) + (-1)^{n-1} I_{-q}(f) = [2]_q \sum_{l=0}^{n-1} (-1)^{n-1-l} q^l f(l) \quad (5)$$

where  $f_n(x)$  is a translation given by  $f_n(x) := f(x+n)$ . It follows from (5) that

$$q I_{-q}(f_1) + I_{-q}(f) = [2]_q f(0). \quad (6)$$

If we replace  $q$  by  $q^{-1}$  in (6), it becomes

$$I_{-q^{-1}}(f_1) + q I_{-q^{-1}}(f) = [2]_q f(0). \quad (7)$$

Recently, Kim et al. [22] considered the case  $f(x) = e^{-x(1+q)t}$  in (7) and derived the following Witt's formula of the Eulerian polynomials for  $n \in \mathbb{N}^*$ :

$$I_{-q^{-1}}(x^n) = \frac{(-1)^n}{(1+q)^n} A_n(-q). \quad (8)$$

Note that Kim and Kim [23, 24] also introduced the  $q$ -analogue of Eulerian polynomials and the  $p$ -adic integral representation of the Boole polynomials.

Based on the work of Kim et al. [22], Araci et al. [7] defined the generating function of Dirichlet's type of the Eulerian polynomials as follows:

$$\sum_{n=0}^{\infty} A_{n,\chi}(-q) \frac{t^n}{n!} = [2]_q \sum_{l=0}^{d-1} (-1)^l q^{d-l+1} \chi(l) \frac{e^{-l(1+q)t}}{e^{-d(1+q)t} + q^d}. \quad (9)$$

Also, they gave the following Witt's formula for Dirichlet's type of the Eulerian polynomials:

$$I_{-q^{-1}}(\chi(x)x^n) = \frac{(-1)^n}{(1+q)^n} A_{n,\chi}(-q).$$

In this paper we construct Dirichlet's type of twisted Eulerian polynomials. By using a complex contour integral representation (or known as Mellin transformation) based on the generating function of Dirichlet type of twisted Eulerian polynomials, we define an  $L$ -type function which interpolates Dirichlet's type of twisted Eulerian polynomials at negative integers.

## ON DIRICHLET'S TYPE OF TWISTED EULERIAN POLYNOMIALS

By using (5), we have

$$I_{-q^{-1}}(f_d) + q^d I_{-q^{-1}}(f) = [2]_q \sum_{l=0}^{d-1} (-1)^l q^{d-l+1} f(l). \quad (10)$$

Let  $C_{p^n} = \{\zeta \mid \zeta^{p^n} = 1\}$  be the cyclic group of order  $p^n$ , and let

$$T_p = \lim_{n \rightarrow \infty} C_{p^n} = C_{p^\infty} = \bigcup_{n \geq 0} C_{p^n},$$

where  $T_p$  is locally constant space. Let  $\chi$  be a Dirichlet character of conductor  $d$  ( $=$  odd) and  $\zeta \in T_p$ . Taking  $f(x) = \zeta^x \chi(x) e^{-x(1+q)t}$  in (10) gives

$$\begin{aligned} I_{-q^{-1}} \left( \zeta^{x+d} \chi(x+d) e^{-(x+d)(1+q)t} \right) + q^d I_{-q^{-1}} \left( \zeta^x \chi(x) e^{-x(1+q)t} \right) \\ = [2]_q \sum_{l=0}^{d-1} (-1)^l q^{d-l+1} \chi(l) e^{-l(1+q)t} \zeta^l. \end{aligned}$$

From this, we derive

$$I_{-q^{-1}} \left( \zeta^x \chi(x) e^{-x(1+q)t} \right) = [2]_q \sum_{l=0}^{d-1} (-1)^l q^{d-l+1} \zeta^l \chi(l) \frac{e^{-l(1+q)t}}{\zeta^d e^{-d(1+q)t} + q^d}. \quad (11)$$

Therefore, we give the following definition of the generating function for Dirichlet's type of twisted Eulerian polynomials.

**Definition 1.** Let  $G_{q,\zeta}(t|\chi) = \sum_{n=0}^{\infty} A_{n,\chi,\zeta}(-q) \frac{t^n}{n!}$  and  $\zeta \in T_p$ . Then we define the twisted Dirichlet's type of Eulerian polynomials by means of the following generating function:

$$\sum_{n=0}^{\infty} A_{n,\chi,\zeta}(-q) \frac{t^n}{n!} = [2]_q \sum_{l=0}^{d-1} (-1)^l q^{d-l+1} \zeta^l \chi(l) \frac{e^{-l(1+q)t}}{\zeta^d e^{-d(1+q)t} + q^d}.$$

By this definition, we have the following corollary.

**Corollary 1.** For any  $\zeta \in T_p$  and  $n \in \mathbb{N}^*$ , we have

$$A_{n,\chi,\zeta}(-q) = \left[ \frac{t^n}{n!} \right] [2]_q \sum_{l=0}^{d-1} (-1)^l q^{d-l+1} \zeta^l \chi(l) \frac{e^{-l(1+q)t}}{\zeta^d e^{-d(1+q)t} + q^d}$$

where  $[t^n] f(t)$  means the coefficient of  $t^n$  in  $f(t)$ .

Now we give an integral representation of the twisted Dirichlet's type of Eulerian polynomials.

**Theorem 1.** The following equalities hold:

$$I_{-q^{-1}} \left( \zeta^x \chi(x) x^n \right) = \int_{Z_p} \zeta^x \chi(x) x^n d\mu_{-q^{-1}}(x) = \frac{(-1)^n}{(1+q)^n} A_{n,\chi,\zeta}(-q) \quad (12)$$

**Proof.** It is evident from (11) and Definition 1.

By Definition 1, we have the following Theorem.

**Theorem 2.** For  $\zeta \in T_p$ , we have

$$G_{q,\zeta}(t|\chi) = \sum_{n=0}^{\infty} A_{n,\chi,\zeta}(-q) \frac{t^n}{n!} = [2]_q \sum_{m=0}^{\infty} \frac{(-1)^m \zeta^m \chi(m) e^{-m(1+q)t}}{q^{m-1}}. \quad (13)$$

**Proof.** From Definition 1, we discover:

$$\begin{aligned}
\sum_{n=0}^{\infty} A_{n,\chi,\zeta}(-q) \frac{t^n}{n!} &= [2]_q \sum_{l=0}^{d-1} (-1)^l q^{d-l+1} \zeta^l \chi(l) \frac{e^{-l(1+q)t}}{\zeta^d e^{-d(1+q)t} + q^d} \\
&= [2]_q \sum_{l=0}^{d-1} (-1)^l q^{-l+1} \zeta^l \chi(l) e^{-l(1+q)t} \sum_{m=0}^{\infty} (-1)^m \zeta^{md} q^{-md} e^{-md(1+q)t} \\
&= q [2]_q \sum_{m=0}^{\infty} \sum_{l=0}^{d-1} (-1)^{l+md} \chi(l+md) q^{-(l+md)} \zeta^{l+md} e^{-(l+md)(1+q)t} \\
&= q [2]_q \sum_{m=0}^{\infty} (-1)^m \zeta^m \chi(m) q^{-m} e^{-m(1+q)t}.
\end{aligned}$$

Thus, we arrive at the desired result.

Now we give the following Theorem, which is an explicit formula for twisted Dirichlet's type of Eulerian polynomials.

**Theorem 3.** For  $\zeta \in T_p$ , we have

$$\frac{(-1)^n}{q(1+q)^{n+1}} A_{n,\chi,\zeta}(-q) = \sum_{m=1}^{\infty} \frac{(-1)^m \zeta^m \chi(m) m^n}{q^m}. \quad (14)$$

**Proof.** By using Taylor expansion of  $e^{-m(1+q)t}$  in (13), we see that

$$\sum_{n=0}^{\infty} A_{n,\chi,\zeta}(-q) \frac{t^n}{n!} = [2]_q \sum_{n=0}^{\infty} \left( \sum_{m=0}^{\infty} \frac{(-1)^m \zeta^m \chi(m) (-m)^n (1+q)^n}{q^{m-1}} \right) \frac{t^n}{n!}$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we arrive at the desired result.

From (12) and (14), we have the following corollary.

**Corollary 2.** For  $\zeta \in T_p$  and  $n \in N$ , we have

$$\lim_{n \rightarrow \infty} \sum_{x=0}^{p^n-1} \frac{(-1)^x \zeta^x \chi(x) x^n}{q^x} = 2q^2 \sum_{m=1}^{\infty} \frac{(-1)^m \zeta^m \chi(m) m^n}{q^m}.$$

Now also, we give multiplication theorem for Dirichlet's type of twisted Eulerian polynomials.

**Theorem 4.** The following is true:

$$\frac{(-1)^n}{(1+q)^n} A_{n,\chi,\zeta}(-q) = \frac{d^n}{[d]_{-q^{-1}}} \sum_{a=0}^{d-1} (-1)^a \chi(a) \zeta^a q^{-a} \int_{Z_p} \left( \frac{a}{d} + x \right)^n \zeta^{dx} d\mu_{-q^{-d}}(x) \quad (15)$$

**Proof.** For each  $\zeta \in T_p$  and  $n \in N^*$  we see that

$$\begin{aligned}
& I_{-q^{-1}} \left( \zeta^x \chi(x) x^n \right) \\
&= \int_{Z_p} \zeta^x \chi(x) x^n d\mu_{-q^{-1}}(x) \\
&= \lim_{m \rightarrow \infty} \frac{1}{[dp^m]_{-q^{-1}}} \sum_{x=0}^{dp^m-1} (-1)^x \zeta^x \chi(x) x^n q^{-x} \\
&= \frac{d^n}{[d]_{-q^{-1}}} \sum_{a=0}^{d-1} (-1)^a \zeta^a \chi(a) q^{-a} \left( \lim_{m \rightarrow \infty} \frac{1}{[p^m]_{-q^{-d}}} \sum_{x=0}^{p^m-1} (-1)^x \zeta^{dx} \left( \frac{a}{d} + x \right)^n q^{-dx} \right) \\
&= \frac{d^n}{[d]_{-q^{-1}}} \sum_{a=0}^{d-1} (-1)^a \chi(a) \zeta^a q^{-a} \int_{Z_p} \left( \frac{a}{d} + x \right)^n \zeta^{dx} d\mu_{-q^{-d}}(x).
\end{aligned}$$

So we complete the proof of the theorem.

**Corollary 3.** For any  $\zeta \in T_p$  and  $n \in \mathbb{N}^*$ , we have

$$A_{n,\chi,\zeta}(-1) = (-2d)^n \sum_{a=0}^{d-1} (-1)^a \chi(a) \zeta^a E_{n,\zeta^d} \left( \frac{a}{d} \right).$$

**Proof.** It is interesting to point out that the case  $q = 1$  in (15) gives

$$\frac{(-1)^n}{2^n} A_{n,\chi,\zeta}(-1) = d^n \sum_{a=0}^{d-1} (-1)^a \chi(a) \zeta^a \int_{Z_p} \left( \frac{a}{d} + x \right)^n \zeta^{dx} d\mu_{-1}(x). \quad (16)$$

With the help of Rim and Kim's work [25], we get

$$E_{n,\zeta}(x) = \int_{Z_p} \zeta^y (x+y)^n d\mu_{-1}(y) \quad (17)$$

where  $\zeta \in T_p$ . Taking  $x=0$  in the above equation, we have  $E_{n,\zeta}(0) := E_{n,\zeta}$ , which is defined by the following generating function:

$$\sum_{n=0}^{\infty} E_{n,\zeta} \frac{t^n}{n!} = \frac{2 \sum_{l=0}^{d-1} (-1)^l \zeta^l e^{lt}}{\zeta^d e^{dt} + 1}, \quad |t| < \frac{\pi}{d}. \quad (18)$$

Combining (16), (17) and (18), we end the proof of Corollary 3.

## ON L-TYPE FUNCTION IN $C$

We describe  $L$ -type function by applying Mellin transformation to the generating function of Dirichlet's type of twisted Eulerian polynomials (stated in Definition 1), which is an interpolated function of Dirichlet's type of twisted Eulerian polynomials at negative integers for the corresponding Cauchy-Residue theorem. Further information about these topics can be obtained elsewhere [4, 6, 7, 15, 17, 19, 25, 31-33].

By (13), for  $s \in C$ , we consider the following complex contour integral representation (or known as Mellin transformation):

$$L_{E,\zeta}(s, \chi) = \frac{\int_0^{\infty} t^{s-1} G_{q,\zeta}(t | \chi) dt}{\int_0^{\infty} t^{s-1} e^{-t} dt}. \quad (19)$$

It follows from (19) that

$$L_{E,\zeta}(s, \chi) = q [2]_q \sum_{m=0}^{\infty} (-1)^m \chi(m) \zeta^m q^{-m} \left\{ \frac{\int_0^{\infty} t^{s-1} e^{-m(1+q)t} dt}{\int_0^{\infty} t^{s-1} e^{-t} dt} \right\}$$

$$= \frac{q}{(1+q)^{s-1}} \sum_{m=1}^{\infty} \frac{(-1)^m \chi(m) \zeta^m}{q^m m^s}.$$

On the other hand, we see that

$$L_{E,\zeta}(s, \chi) = \sum_{n=0}^{\infty} \frac{A_{n,\chi,\zeta}(-q)}{n!} \left( \frac{\int_0^{\infty} t^{s-n+1} dt}{\int_0^{\infty} t^{s-1} e^{-t} dt} \right). \quad (20)$$

Therefore, we give the definition of twisted Eulerian  $L$ -function as follows:

**Definition 2.** Let  $\zeta \in T_p$  and  $s \in C$ ; we define

$$L_{E,\zeta}(s, \chi) = \frac{q}{(1+q)^{s-1}} \sum_{m=1}^{\infty} \frac{(-1)^m \chi(m) \zeta^m}{q^m m^s}.$$

By (14) and Definition 2, we derive the following Theorem.

**Theorem 5.** The following equality holds:

$$L_{E,\zeta}(-n, \chi) = (-1)^n A_{n,\chi,\zeta}(-q).$$

**Proof.** From (14):

$$\frac{(-1)^n}{q(1+q)^{n+1}} A_{n,\chi,\zeta}(-q) = \sum_{m=1}^{\infty} \frac{(-1)^m \zeta^m \chi(m) m^n}{q^m}.$$

Putting  $s = -n$  in Definition 2 gives

$$L_{E,\zeta}(-n, \chi) = q(1+q)^{n+1} \sum_{m=1}^{\infty} \frac{(-1)^m \chi(m) \zeta^m m^n}{q^m}$$

from which we have

$$L_{E,\zeta}(-n, \chi) = (-1)^n A_{n,\chi,\zeta}(-q).$$

Thus, the proof of the theorem is completed.

## ACKNOWLEDGEMENTS

The authors would like to thank the referees for many valuable comments and suggestions, through which we have substantially improved the presentation of the paper.

## REFERENCES

1. S. Araci, "Novel identities involving Genocchi numbers and polynomials arising from applications of umbral calculus", *Appl. Math. Comput.*, **2014**, 233, 599-607.
2. S. Araci, "Novel identities for  $q$ -Genocchi numbers and polynomials", *J. Funct. Spaces Appl.*, **2012**, 2012, Article ID 214961.
3. S. Araci, M. Acikgoz and E. Sen, "On the extended Kim's  $p$ -adic  $q$ -deformed fermionic integrals in the  $p$ -adic integer ring", *J. Number Theory*, **2013**, 133, 3348-3361.
4. S. Araci, M. Acikgoz, K. H. Park and H. Jolany, "On the unification of two families of multiple twisted type polynomials by using  $p$ -adic  $q$ -integral at  $q = -1$ ", *Bull. Malays. Math. Sci. Soc.*,

**2014**, 37, 543-554.

5. S. Araci, E. Sen and M. Acikgoz, "Theorems on Genocchi polynomials of higher order arising from Genocchi basis", *Taiwan. J. Math.*, **2014**, 18, 473-482.
6. S. Araci, M. Acikgoz and H. Jolany, "On  $p$ -adic interpolating function associated with modified Dirichlet's type of twisted  $q$ -Euler numbers and polynomials with weight  $\alpha$ ", *J. Class. Anal.*, **2013**, 2, 35-48.
7. S. Araci, M. Acikgoz and E. Ağyüz, "On the Dirichlet's type of Eulerian polynomials", *Math. Sci.*, **2014**, 8, 131.
8. S. Araci, A. Bagdasaryan, C. Özel and H. M. Srivastava, "New symmetric identities involving  $q$ -zeta-type functions", *Appl. Math. Inform. Sci.*, **2014**, 8, 2803-2808.
9. S. Araci, M. Acikgoz, A. Bagdasaryan and E. Sen, "The Legendre polynomials associated with Bernoulli, Euler, Hermite and Bernstein polynomials", *Turk. J. Anal. Number Theory*, **2013**, 1, 1-3.
10. S. Araci and M. Acikgoz, "A note on the Frobenius-Euler numbers and polynomials associated with Bernstein polynomials", *Adv. Stud. Contemp. Math.*, **2012**, 22, 399-406.
11. T. Kim, "On a  $q$ -analogue of the  $p$ -adic log gamma functions and related integrals", *J. Number Theory*, **1999**, 76, 320-329.
12. T. Kim, " $q$ -Volkenborn integration", *Russian J. Math. Phys.*, **2002**, 19, 288-299.
13. T. Kim, "Non-archimedean  $q$ -integrals associated with multiple Changhee  $q$ -Bernoulli polynomials", *Russian J. Math. Phys.*, **2003**, 10, 91-98.
14. T. Kim, " $p$ -Adic  $q$ -integrals associated with the Changhee-Barnes'  $q$ -Bernoulli polynomials", *Integr. Transf. Special Funct.*, **2004**, 15, 415-420.
15. T. Kim, "Analytic continuation of multiple  $q$ -zeta functions and their values at negative integers", *Russian J. Math. Phys.*, **2004**, 11, 71-76.
16. T. Kim, "Some identities on the  $q$ -Euler polynomials of higher order and  $q$ -stirling numbers by the fermionic  $p$ -adic integral on  $Z_p$ ", *Russian J. Math. Phys.*, **2009**, 16, 484-491.
17. T. Kim, "On the  $q$ -extension of Euler and Genocchi numbers", *J. Math. Anal. Appl.*, **2007**, 326, 1458-1465.
18. T. Kim, "On the analogs of Euler numbers and polynomials associated with  $p$ -adic  $q$ -integral on  $Z_p$  at  $q = -1$ ", *J. Math. Anal. Appl.*, **2007**, 331, 779-792.
19. T. Kim, S. H. Lee, H. H. Han and C. S. Ryoo, "On the values of the weighted  $q$ -Zeta and  $L$ -functions", *Discrete Dynam. Nat. Soc.*, **2011**, 2011, Article ID 476381.
20. T. Kim, D. S. Kim, T. Mansour, S.-H. Rim and M. Schork, "Umbral calculus and Sheffer sequences of polynomials", *J. Math. Phys.*, **2013**, 54, 083504.
21. T. Kim, D. S. Kim, D. V. Dolgy and S. H. Rim, "Some identities on the Euler numbers arising from Euler basis polynomials", *Ars Combin.*, **2013**, 109, 433-446.
22. D. S. Kim, T. Kim, W. J. Kim and D. V. Dolgy, "A note on Eulerian polynomials", *Abstr. Appl. Anal.*, **2012**, 2012, Article ID 269640.
23. D. S. Kim and T. Kim, "A note on  $q$ -Eulerian polynomials", *Proc. Jangjeon Math. Soc.*, **2013**, 16, 445-450.
24. D. S. Kim and T. Kim, "A note on Boole polynomials", *Integr. Transf. Special Funct.*, **2014**, 25, 627-633.
25. S.H. Rim and T. Kim, "A note on  $q$ -Euler numbers associated with the basic  $q$ -zeta function",

*Appl. Math. Lett.*, **2007**, 20, 366-369.

26. Y. He and C. Wang, "Some formulae of products of the Apostol-Bernoulli and Apostol-Euler polynomials", *Discrete Dynam. Nat. Soc.*, **2012**, 2012, Article ID 927953.
27. H. Ozden, I. N. Cangul and Y. Simsek, "Multivariate interpolation functions of higher order  $q$ -Euler numbers and their applications", *Abstr. Appl. Anal.*, **2008**, 2008, Article ID 390857.
28. I. N. Cangul, H. Ozden and Y. Simsek, "A new approach to  $q$ -Genocchi numbers and their interpolation functions", *Nonlinear Anal.*, **2009**, 71, e793-e799.
29. I. N. Cangul, H. Ozden and Y. Simsek, "Generating functions of the  $(h, q)$  extension of twisted Euler polynomials and numbers", *Acta Math. Hungar.*, **2008**, 120, 281-299.
30. B.-N. Guo and F. Qi, "Some identities and an explicit formula for Bernoulli and Stirling numbers", *J. Comput. Appl. Math.*, **2014**, 255, 568-579.
31. H. M. Srivastava, "Some formulas for the Bernoulli and Euler polynomials at rational arguments", *Math. Proc. Cambridge Philos. Soc.*, **2000**, 129, 77-84.
32. H. M. Srivastava, "Some generalizations and basic (or  $q$ -) extensions of the Bernoulli, Euler and Genocchi polynomials", *Appl. Math. Inform. Sci.*, **2011**, 5, 390-444.
33. H. M. Srivastava and J. Choi, "Zeta and  $q$ -Zeta Functions and Associated Series and Integrals", Elsevier Science Publishers, Amsterdam, **2012**.