

## REFERENCES

1. Pisek P, Manosudprasit M. Comparison of dento-skeletal conditions of orthodontic patients with and without unilateral complete clefts of primary and secondary palates. **Thai J Oral Maxillofac Surg** 2008; 22: 36-44.
2. Boyarskiy S, Choi HJ, Park K. Evaluation of alveolar bone support of the permanent canine in cleft and noncleft patients. **Cleft Palate Craniofac J** 2006; 43(6): 678-82.
3. Adlam DM, Yau CK, Banks P. A retrospective study of the stability of midface osteotomies in cleft lip and palate patients. **Br J Oral Maxillofac Surg** 1989; 27: 265-76.
4. Subtelny JD. Orthodontic treatment of cleft lip and palate, Birth to adulthood. **Cleft Palate J** 1966; 36: 273-92.
5. Mars M, Houston WJB. A preliminary study of facial growth and morphology in unoperated male unilateral cleft lip and palate subjects over 13 years of age. **Cleft Palate J** 1990; 27: 7-10.
6. Graber TM. The congenital cleft palate deformity. **J Am Dent Assoc** 1954; 48: 375-95.
7. Roberts CT, Semb G, Shaw WC. Strategies for the advancement of surgical methods in cleft lip and palate. **Cleft Palate Craniofac J** 1991; 28: 141-9.
8. Shaw WC, Dahl E, Asher-McDade C, Brattstrom V, Mars M, Mc William J, et al. A six-center international study of treatment outcome in patients with clefts of the lip and palate. Part 5: General discussion and conclusions. **Cleft Palate Craniofac J** 1992; 29: 413-8.
9. Peltomaki T, Venditti BL, Grayson BH, Cutting CB, Brecht LE. Associations between severity of clefting and maxillary growth in patients with unilateral cleft lip and palate treated with infant orthopedics. **Cleft Palate Craniofac J** 2001; 38(6): 582-6.

10. Fonseca RJ, Turvey TA, Wolford LM. Orthognathic surgery in the cleft patient. In: Fonseca JR, Baker SB, Wolford LM, editors. **Oral and Maxillofacial surgery: cleft/craniofacial/cosmetic surgery**. Philadelphia: W.B.Saunders; 2000.
11. Bell WH, Jacobs JD, Quejada JG. Simultaneous repositioning of the maxilla, mandible and chin. Treatment planning and analysis of soft tissues. **Am J Orthod** 1986; 89:28-50.
12. Pieter J, Nollet PM, Katsaros C, Martin A, Kuijpers-Jagtman AM. Treatment outcome in unilateral cleft lip and palate evaluated with the Goslon Yardstick: A meta-analysis of 1236 patients. **Plast Reconstr Surg** 2005; 10: 1255-62.
13. Lines PA, Steinhauser EW. Soft tissue changes in relationship to movement of hard structures in orthognathic surgery: a preliminary report. **J Oral Surg** 1974; 32: 891-6.
14. McDonell JP, McNeill RW, West RA. Advancement genioplasty: a retrospective cephalometric analysis of osseous and soft tissue changes. **J Oral Surg** 1977; 37: 640-7.
15. Suckiel JM, Kohn MW. Soft tissue changes related to the surgical management of mandibular prognathism. **Am J Orthod** 1978; 73: 676-80.
16. Hunt NP, Rudge SJ. Facial profile and orthognathic surgery. **Br J Orthod** 1984; 11: 126-36.
17. Baker NJ, David S, Barnard DW, Birnie DJ, Robinson SN. Occlusal outcome in patients undergoing orthognathic surgery with internal fixation. **Br J Oral Maxillofac Surg** 1999; 37: 90-3.
18. Summers CJ. The occlusal index: a system for identifying and scoring occlusal disorders. **Am J Orthod** 1971; 59: 552-67.
19. Myrberg N, Thilander B. An evaluation of the duration and the results of orthodontic treatment. **Scand J Dent Res** 1973; 81: 85-91.
20. Gottlieb EL. Grading your orthodontic treatment results. **J clinic orthod** 1975; 9: 155-61.
21. Daniels C, Richmond S. The development of the Index of Complexity, Outcome and Need (ICON). **Br J Orthod** 2000; 27: 149-62.

22. **Orthodontics guidelines for treatment need.** [online]. 2007 January 3 [cited 2008 Dec 5] Available from:<http://www.dentalgain.org/orthodontics.html>.
23. DeGuzman L, Bahiraei D, Vig KWL, Vig PS, Weyant RJ, O'Brien K. The validation of the Peer Assessment Rating index for malocclusion severity and treatment difficulty. **Am J Orthod Dentofac Orthop** 1995; 107: 172-6.
24. Deguchi T, Honjo T, Fukunaga T, Miyawaki S, Roberts WE, Yamamotoa TT. Clinical assessment of orthodontic outcomes with the peer assessment rating, discrepancy index, objective grading system, and comprehensive clinical assessment. **Am J Orthod Dentofac Orthop** 2005; 127: 434-43.
25. Fogh-Anderson P. **Inheritance of harelip and cleft palate.** [n.p.]; Indiana: Copenhegan, Nyt Nordisk Forlag, Busck A, editors. 1942.
26. Vanderas AP. Incidence of cleft lip, cleft palate, and cleft lip and palate among races: a review. **Cleft Palate J** 1987; 24(3): 216-25.
27. Tan KL. Incidence and epidermology of cleft lip and palate in Singapore. **Ann Acad Med Singapore** 1988; 17: 311-4.
28. Chungsumanich A, Aojanepong C, Muangsombut S, Tomgpiew P. Epidemiology of cleft lip and palate in Thailand. **Ann Plast Surg** 1998; 41: 7-10.
29. Ruangsitt C, Phraserthsang P, Banpho Y, Lamduan W, Giathamnuay S, Nuwantha A. **Incidence of cleft lip and cleft palate in 3 hospitals in Khon Kaen.** Khon Kaen: Department of Orthodontics, Faculty of Dentistry, Khon Kaen University; 1993.
30. Kernahan D A, Stark R B. A new classification for cleft lip and palate. **Plast Reconstr Surg** 1958; 22: 435-41.
31. Santiago PE, Grayson BH, Cutting CB, Gianoutsos MP, Brecht LE, Kwon SM. Reduced need for alveolar bone grafting by presurgical orthopedics and primary gingivoperiosteoplasty. **Cleft Palate Craniofac J** 1998; 35: 77-80.
32. Tang E, So L. Prevalence and severity of malocclusion in children with cleft lip and/or palate in Hong Kong. **Cleft Palate Craniofac J** 1992; 29(3): 287-91.
33. Summers CJ. The occlusal index: A system for identifying and scoring occlusal disorders. **Am J Orthod** 1971; 59: 552-67.

34. Mars M, Plint DA, Houston JB, Bergland O, Semb G. The Goslon Yardstick: A New System of Assessing Dental Arch Relationships in Children with Unilateral Clefts of the Lip and Palate. **Cleft Palate Craniofac J** 1987; 24: 314-22.
35. Atack N, Hathorn I, Semb G, Dowell T, Sandy JR. A new index for assessing surgical outcome in unilateral cleft lip and palate subjects aged five: Reproducibility and validity. **Cleft Palate Craniofac J** 1997; 34: 242-6.
36. Huddart AG, Bodenham RS. The evaluation of arch form and occlusion in unilateral cleft palate subjects. **Cleft Palate J** 1972; 9: 194-209.
37. Tobiasen JM, Hiebert JM. Combined effects of severity of cleft impairment and facial attractiveness on social perception: An experimental study. **Cleft Palate Craniofac J** 1993; 30(1): 82-6.
38. Slade P, Bishop S, Jowett R. Relationships between cleft severity and attractiveness of newborns with unrepaired clefts. **Cleft Palate Craniofac J** 1995; 32(4): 318-22.
39. Ozturk Y, Cura N. Examination of Craniofacial Morphology in Children with Unilateral Cleft Lip and Palate. **Cleft Palate Craniofac J** 1996; 33(1): 32-6.
40. Edwards M, Watson A C H. **Advances in the Management of Cleft Palate**. New York: Churchill Livingstone; 1980.
41. Bishara SE. the influence of palatoplasty and cleft length on facial development. **Cleft Palate J** 1993; 10: 390-8.
42. Trotman CA, Collett AR. Analyses of craniofacial and dental morphology in monozygotic twin discordant for cleft lip and unilateral cleft lip and palate. **Angle Orthod** 1993; 63: 135-40.
43. Dahl E. Cranifacial morphology in congenital clefts of the lip and palate: An x-ray cephalometric study of young adult males. Copenhagen: **Acta Odontol Scandinavica** 1970; 28(Suppl 57): 11.
44. Aduss H. Craniofacial growth in complete unilateral cleft lip and palate. **Angle Orthod** 1971; 41: 202-13.

45. Hermann NV, Jensen BL, Dahl E, Bolund S, Kreiborg S. Craniofacial comparisons in 22-month-old lip-operated children with unilateral complete cleft lip and palate and unilateral incomplete cleft lip. **Cleft Palate Craniofac J** 2000; 37: 303-17.
46. Chowchuen B, Kijshawong S, Prathanee B, Ratanayatikul J, Pujjanasunthon N, Choosilp K, et al. Interdisciplinary Cleft Team Management of Center for Cleft Lip-Palate and Craniofacial Anomalies, Khon Kaen University. In: Chowchuen B, Prathanee B, Ratanayatikul C, editors. **Interdisciplinary care of cleft lip, cleft palate and craniofacial anomalies**. Khon Kaen: Siripan Offset; 2004. p.16-41.
47. Mars M, Asher-McDade C, Brattstrom V, Dahl E, McWilliam J, Molsted K, et al. A six-center international study of treatment outcome in patients with clefts of the lip and palate: part 3. Dental arch relationships. **Cleft Palate Craniofac J** 1992; 29: 405-8.
48. Ross RB. Treatment variables affecting facial growth in complete unilateral cleft lip and palate. Part 1-7. **Cleft Palate J** 1987; 24: 5-77.
49. Pruzansky S. Presurgical orthopedics and bone grafting for infants with cleft lip and palate: a dissent. **Cleft Palate J** 1964; 1: 411-8.
50. Atack N, Hathorn I, Mars M, Sandy J. Study models of 5 year old children as predictors of surgical outcome in unilateral cleft lip and palate. **Eur J Orthod** 1997; 19: 165-70.
51. Pae EK. Measurements must be interval, not ordinal. **Angle Orthod** 1999; 69: 397.
52. Llewellyn SK, Hamdan AM, Rock WP. An index of orthodontic treatment complexity. **Eur J Orthod** 2007; 29: 186-92.
53. Richmond S, Shaw WC, O'Brien KD, Buchanan IB, Jones R, Stephens CD, et al. The development of the PAR Index (Peer Assessment Rating): reliability and validity. **Eur J Orthod** 1992; 14: 125-39.
54. Richmond S, Shaw WC, Roberts CT, Andrews M. The PAR index (Peer Assessment Rating): methods to determine outcome of orthodontic treatment in terms of improvement and standards. **Eur J Orthod** 1992; 14: 180-7.

55. Richmond S, Andrews M. Orthodontic treatment standards in Norway. **Eur J Orthod** 1993; 15: 7-15.
56. Brook PH, Shaw WC. The development of an index of orthodontic treatment priority. **Eur J Orthod** 1989; 11: 309-20.
57. Holmes A. The prevalence of orthodontic treatment need. **Br J Orthod** 1992; 79: 177-82.
58. Draker HL. Handicapping labiolingual deviations: a proposed index for public health purposes. **Am J Orthod** 1960; 65: 295-305.
59. Draker HL. Handicapping labio-lingual deviations: a proposed index for public health purposes. **J Public Health Dent** 1958; 18: 1-7.
60. Theis JE, Huang G, King GJ, Omnell ML. Eligibility for publicly funded orthodontic treatment determined by the handicapping labiolingual deviation index. **Am J Orthod Dentofac Orthop** 2005; 128: 708-15.
61. Deacon S, Bessant P, Russell JI, Hathorn I. What are the occlusal outcomes for unilateral cleft lip and palate patients? A national project in the UK. **Br Dent J** 2007; 203: E18.
62. Holman JK, Hans MG, Nelson S, Powers MP. An assessment of extraction versus nonextraction orthodontic treatment using the peer assessment rating (PAR) index. **Angle Orthod** 1998; 68: 527-34.
63. Birkeland K, Furevik J, Boe OE, Wisth PJ. Evaluation of treatment and post-treatment changes by the PAR Index. **Eur J Orthod** 1997; 19: 279-88.

## **APPENDICES**

## **APPENDIX A**

### **Component of The Peer Assessment Rating (PAR) Index**


There are 11 components of the PAR Index to be assessed according to the criteria of Richmond et al.<sup>53</sup>

**Table A1** The PAR index components<sup>53</sup>

1.	Upper right segment
2.	Upper anterior segment
3.	Upper left segment
4.	Lower right segment
5.	Lower anterior segment
6.	Lower left segment
7.	Right buccal occlusion
8.	Overjet
9.	Overbite
10.	Centreline
11.	Left buccal occlusion

### ***Buccal segments***


The measuring zone is from the mesial anatomical contact point of the first permanent molar to the distal anatomical contact point of the canine (Figure 2).



**Figure 2** Showing the measuring zone of buccal segments in cleft lip and palate patient's models

### ***Anterior segments***

The measuring zone is from the mesial anatomical contact point of the canine on one side to the mesial anatomical contact point of the canine on the opposite side (Figure 3).



**Figure 3** Showing the measuring zone of anterior segments in cleft lip and palate patient's models

The occlusal features recorded are crowding, spacing, and impacted teeth. Displacements are recorded as the shortest distance between contact points of adjacent teeth. The greater the distance, the greater the PAR score.


An impacted tooth is recorded when the space for this tooth is less than or equal to 4 mm. Impacted canines are recorded in the anterior segment.

**Table A2** Tooth Displacement PAR scores<sup>53</sup>

Score	Discrepancy
0	0 mm to 1 mm
1	1.1 mm to 2 mm
2	2.1 mm to 4 mm
3	4.1 mm to 8 mm
4	greater than 8 mm
5	impacted teeth

### **Buccal occlusion**

The recording zone is from the canine to the last molar (first, second, or third molar)(Figure 4). All discrepancies are recorded in three planes when the teeth are in occlusion.



**Figure 4** Showing the measuring zone of buccal occlusion in cleft lip and palate patient's models


**Buccal occlusion assessments** (Temporary developmental stages and submerging deciduous teeth are excluded.)

**Table A3** Buccal occlusion PAR assessments<sup>53</sup>

Score	Discrepancy
Antero-posterior	
0	Good interdigitation Class I, II and III
1	Less than half unit discrepancy
2	Half a unit discrepancy (cusp to cusp)
Vertical	
0	No discrepancy in interdigitation
1	Lateral open bite on at least two teeth greater than 2 mm
Transverse	
0	No cross-bite
1	Cross-bite tendency
2	Single tooth in cross-bite
3	More than one tooth in cross-bite
4	More than one tooth in scissor bite

### *Overjet*

The recording zone is from the left to right lateral incisors (Figure 5). The most prominent aspect of any one incisor is recorded.



**Figure 5** Showing the measuring zone of overjet and overbite in cleft lip and palate patient's models

**Table A4** Overjet PAR measurements<sup>53</sup>

Score	Discrepancy
Over jet	
0	0 - 3 mm
1	3.1 - 5 mm
2	5.1 - 7 mm
3	7.1 - 9 mm
4	Greater than 9 mm
Anterior cross-bites	
0	No discrepancy
1	One or more teeth edge-to-edge
2	One single tooth in cross-bite
3	Two teeth in cross-bite
4	More than two teeth in cross-bite

### ***Overbite***

The recording zone includes the lateral incisors (Figure 5). The tooth with the greatest overlap is recorded as the vertical overlap or open bite.


**Table A5** Overbite PAR measurements<sup>53</sup>

Score	Discrepancy
Open bite	
0	No open bite
1	Open bite less than and equal to 1 mm
2	Open bite 1.1 – 2 mm
3	Open bite 2.1 – 3 mm
4	Open bite greater than or equal to 4 mm
Overbite	
1	Less than or equal to one third coverage of the lower incisor
2	Greater than one third, but less than two thirds coverage of the lower incisor
3	Greater than two thirds coverage of the lower incisor
4	Greater than or equal to full tooth coverage

### ***Centreline***

The center-line discrepancy is recorded in relation to the lower incisors (Figure

- 6). If a lower central incisor has been extracted the measurement is not recorded.

**Figure 6** Showing centreling measurment

**Table A6** Centre-line PAR assessments<sup>53</sup>

Score	Discrepancy
0	Coincident and up to one quarter lower incisor width
1	One quarter to one half lower incisor width
2	Greater than one half lower incisor width

**Table A7** Degree of improvement of PAR assessments<sup>53</sup>

Score reduction (Pretreatment - Posttreatment)	Degree of improvement
≥ 22	Greatly improved
< 22 and ≥ 30%	Improved
< 22 and ≤ 30%	Worse-no different

**Conventions for the PAR index<sup>53</sup>***General*

1. All scoring is accumulative.
2. There is no maximal cut-off level
3. The occlusion should be scored disregarding functional displacement (this cannot be determined from dental casts alone).
4. The contact points between first, second, and third molars are not recorded. The contact points between molars are so variable, however, severe deviations will produce a cross-bite, and will be noted in the buccal occlusions.
5. If the contact point displacement is as a result of poor restorative work (restorations or crowns), the displacement is not recorded.
6. Contact points between deciduous teeth are not recorded.
7. Extraction spaces are not recorded if the patient is to receive a prosthetic replacement. However, if space closure is intended, the distance between adjacent teeth should be noted.



### *Canines*

1. Where there are missing canines, displacements resulting from discrepancies between the mesial contact point to the first premolar and the distal of the lateral incisor should be recorded in the anterior segment.
2. Canine cross-bites should be recorded in the overjet section.
3. Contact points between the canines and premolars are scored as follows: the distal contact point of the canine to the midpoint on the mesial surface of the adjacent premolar. (These contact points are so variable. When untreated normal occlusions were assessed this relationship seemed to be the most acceptable.)

### *Impactions*

If a tooth is unerupted and displaced from the line of the arch either buccally or palatally due to insufficient space, this is regarded as an impaction. However, if the tooth is erupted and displaced, the displacement score is recorded.

### *Incisors*

1. If there is agenesis of the upper incisor or the tooth has been lost due to trauma or caries the procedure is as follows:
  - (a) if the space is maintained (for a prosthesis), the distance between adjacent teeth is not recorded;
  - (b) if the space is to be closed, the distance between adjacent teeth is recorded.
2. When recording an overjet, if the tooth falls on the line the lower grade is recorded.
3. If a lower incisor has been extracted or is missing, the centerline is not recorded.

### *Molars*

1. Contact points between first and second molars are not recorded. If the first molars have been extracted, the contact point of the second molar is recorded.
2. If the first molars have been extracted, the contact point of the second molar is recorded.

**APPENDIX B**

**Reliability of the PAR Index**

**Table B1** The reliability of PAR Index score between two trained examiners by Intra-class Correlation Coefficient (ICC)

<b>Reliability</b>	<b>ICC</b>	<b>95% CI of ICC</b>
Intra-examiner1	0.989	0.956 – 0.997
Intra-examiner2	0.994	0.977 – 0.999
Inter-examiner (1 <sup>st</sup> time)	0.989	0.957 – 0.997
Inter-examiner (2 <sup>nd</sup> time)	0.988	0.951 – 0.977

**APPENDIX C**

**Raw Data for Patient's General Characteristics**

**Table C1** Gender and age of all samples

No	Gender	Age (year)
1	Male	11
2	Female	11
3	Male	16
4	Male	13
5	Male	11
6	Female	18
7	Female	12
8	Female	12
9	Female	12
10	Male	12
11	Female	15
12	Male	18
13	Female	11
14	Male	21
15	Male	28
16	Female	18
17	Female	16
18	Female	17
19	Female	22
20	Male	13
21	Female	14
22	Female	17
23	Female	12
24	Female	13
25	Female	12
26	Female	15
27	Female	11
28	Male	16
29	Female	13
30	Female	18
31	Female	23
32	Female	30
33	Male	14
34	Male	11

**APPENDIX D**  
**Raw Data for PAR Index Score**

### PAR Index Score of Orthodontic Treatment Alone Group

**Table D1** Case No. 1

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 1, (5-4) 0, (4-3) 1	(6-5) 0, (5-4) 0, (4-3) 0		
Upper anterior segment	(3-2) 3, (2-1) 0, (1-1) 0 (1-2) 2, (2-3) 0	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Upper left segment	(6-5) 3, (5-4) 2, (4-3) 2	(6-5) 0, (5-4) 0, (4-3) 0		
Lower right segment	(6-5) 0, (5-4) 1, (4-3) 1	(6-5) 0, (5-4) 0, (4-3) 0		
Lower anterior segment	(3-2) 1, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 1	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5) 0, (5-4) 2, (4-3) 1	(6-5) 0, (5-4) 0, (4-3) 0		
Right buccal occlusion	AP 2 T 0 V 3	AP 0 V 0 T 0		
Overjet	OJ 0 x-bite 4	OJ 0 x-bite 0		
Overbite	openbite . 0 OB 0	openbite 0 OB 0		
Centerline	0	0		
Left buccal occlusion	AP 2 V 0 T 1	AP 0 V 0 T 0		
<b>Total</b>	<b>33</b>	<b>0</b>	<b>33</b>	<b>100</b>

**Table D2** Case No. 2

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) <b>5</b> , (5-4) <b>2</b> , (4-3) <b>0</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Upper anterior segment	(3-2) <b>5</b> , (2-1) <b>0</b> , (1-1) <b>0</b> (1-2) <b>1</b> , (2-3) <b>5</b>		(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>			
Upper left segment	(6-5) <b>5</b> , (5-4) <b>3</b> , (4-3) <b>2</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Lower right segment	(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Lower anterior segment	(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>		(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>			
Lower left segment	(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>		(6-5) <b>1</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Right buccal occlusion	AP <b>1</b> T <b>0</b> V <b>2</b>		AP <b>1</b> V <b>0</b> T <b>0</b>			
Overjet	OJ <b>0</b> x-bite <b>4</b>		OJ <b>0</b> x-bite <b>0</b>			
Overbite	openbite <b>0</b> OB <b>2</b>		openbite <b>0</b> OB <b>0</b>			
Centerline		<b>0</b>		<b>0</b>		
Left buccal occlusion	AP <b>2</b> V <b>0</b> T <b>3</b>		AP <b>1</b> V <b>0</b> T <b>0</b>			
<b>Total</b>	<b>42</b>		<b>3</b>		<b>39</b>	<b>92.86</b>

**Table D3** Case No. 3

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 0, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Upper anterior segment	(3-2) 1, (2-1) 1, (1-1) 2 (1-2) 0, (2-3) 0	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Upper left segment	(6-5) 3, (5-4) 0, (4-3) 0	(6-5) 1, (5-4) 0, (4-3) 0		
Lower right segment	(6-5) 2, (5-4) 2, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Lower anterior segment	(3-2) 1, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 2	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5) 0, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Right buccal occlusion	AP 0 T 0 V 0	AP 0 V 0 T 0		
Overjet	OJ 0 x-bite 4	OJ 0 x-bite 0		
Overbite	openbite 0 OB 0	openbite 0 OB 0		
Centerline	2	2		
Left buccal occlusion	AP 0 V 0 T 2	AP 0 V 0 T 0		
<b>Total</b>	<b>22</b>	<b>3</b>	<b>19</b>	<b>86.36</b>

**Table D4** Case No. 4

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) <b>2</b> , (5-4) <b>1</b> , (4-3) <b>2</b>	(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>		
Upper anterior segment	(3-2) <b>1</b> , (2-1) <b>0</b> , (1-1) <b>3</b> (1-2) <b>3</b> , (2-3) <b>0</b>	(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>		
Upper left segment	(6-5) <b>2</b> , (5-4) <b>1</b> , (4-3) <b>2</b>	(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>		
Lower right segment	(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>	(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>		
Lower anterior segment	(3-2) <b>1</b> , (2-1) <b>0</b> , (1-1) <b>0</b> (1-2) <b>1</b> , (2-3) <b>1</b>	(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>		
Lower left segment	(6-5) <b>1</b> , (5-4) <b>0</b> , (4-3) <b>2</b>	(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>		
Right buccal occlusion	AP <b>2</b> T <b>0</b> V <b>2</b>	AP <b>1</b> V <b>0</b> T <b>0</b>		
Overjet	OJ <b>0</b> x-bite <b>3</b>	OJ <b>0</b> x-bite <b>0</b>		
Overbite	openbite <b>0</b> OB <b>0</b>	openbite <b>0</b> OB <b>0</b>		
Centerline		<b>1</b>	<b>0</b>	
Left buccal occlusion	AP <b>2</b> V <b>0</b> T <b>3</b>	AP <b>0</b> V <b>0</b> T <b>0</b>		
<b>Total</b>	<b>36</b>		<b>1</b>	<b>35</b> <b>97.22</b>

**Table D5** Case No. 5

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 0, (5-4) 1, (4-3) 2	(6-5) 0, (5-4) 0, (4-3) 0		
Upper anterior segment	(3-2) 2, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 5	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Upper left segment	(6-5) 0, (5-4) 1, (4-3) 0	(6-5) 1, (5-4) 0, (4-3) 0		
Lower right segment	(6-5) 0, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Lower anterior segment	(3-2) 0, (2-1) 0, (1-1) 1 (1-2) 0, (2-3) 1	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5) 0, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Right buccal occlusion	AP 1 T 0 V 0	AP 0 V 0 T 0		
Overjet	OJ 0 x-bite 4	OJ 0 x-bite 0		
Overbite	openbite 0 OB 2	openbite 0 OB 0		
Centerline	1	0		
Left buccal occlusion	AP 0 V 0 T 3	AP 0 V 0 T 0		
<b>Total</b>	<b>24</b>	<b>1</b>	<b>23</b>	<b>95.83</b>

**Table D6** Case No. 6

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 0, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Upper anterior segment	(3-2) 1, (2-1) 4, (1-1) 2 (1-2) 4, (2-3) 1	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Upper left segment	(6-5) 1, (5-4) 0, (4-3) 0	(6-5) 1, (5-4) 0, (4-3) 0		
Lower right segment	(6-5) 1, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Lower anterior segment	(3-2) 1, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 1	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5) 2, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Right buccal occlusion	AP 0 T 0 V 2	AP 2 V 0 T 0		
Overjet	OJ 3 x-bite 4	OJ 1 x-bite 0		
Overbite	openbite 4 OB 2	openbite 0 OB 0		
Centerline	1	0		
Left buccal occlusion	AP 0 V 0 T 2	AP 0 V 0 T 0		
<b>Total</b>	<b>36</b>	<b>4</b>	<b>32</b>	<b>88.89</b>

**Table D7** Case No. 7

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) <b>5</b> , (5-4) <b>2</b> , (4-3) <b>2</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Upper anterior segment	(3-2) <b>1</b> , (2-1) <b>2</b> , (1-1) <b>3</b>		(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b>			
	(1-2) <b>0</b> , (2-3) <b>3</b>		(1-2) <b>0</b> , (2-3) <b>0</b>			
Upper left segment	(6-5) <b>1</b> , (5-4) <b>0</b> , (4-3) <b>1</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Lower right segment	(6-5) <b>0</b> , (5-4) <b>1</b> , (4-3) <b>2</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Lower anterior segment	(3-2) <b>0</b> , (2-1) <b>1</b> , (1-1) <b>0</b>		(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b>			
	(1-2) <b>0</b> , (2-3) <b>1</b>		(1-2) <b>0</b> , (2-3) <b>0</b>			
Lower left segment	(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Right buccal occlusion	AP <b>2</b>		AP <b>0</b>			
	T <b>0</b>		V <b>0</b>			
	V <b>4</b>		T <b>0</b>			
Overjet	OJ <b>0</b>		OJ <b>1</b>			
	x-bite <b>4</b>		x-bite <b>0</b>			
Overbite	openbite <b>2</b>		openbite <b>0</b>			
	OB <b>0</b>		OB <b>0</b>			
Centerline	<b>2</b>		<b>0</b>			
Left buccal occlusion	AP <b>1</b>		AP <b>0</b>			
	V <b>0</b>		V <b>0</b>			
	T <b>2</b>		T <b>0</b>			
<b>Total</b>	<b>42</b>		<b>1</b>		<b>41</b>	<b>97.62</b>

**Table D8** Case No. 8

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 0, (5-4) 3, (4-3) 3	(6-5) 0, (5-4) 0, (4-3) 0		
Upper anterior segment	(3-2) 1, (2-1) 0, (1-1) 3	(3-2) 0, (2-1) 0, (1-1) 0		
	(1-2) 0, (2-3) 0	(1-2) 0, (2-3) 0		
Upper left segment	(6-5) 2, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Lower right segment	(6-5) 0, (5-4) 2, (4-3) 3	(6-5) 0, (5-4) 0, (4-3) 0		
Lower anterior segment	(3-2) 0, (2-1) 0, (1-1) 0	(3-2) 0, (2-1) 0, (1-1) 0		
	(1-2) 0, (2-3) 0	(1-2) 0, (2-3) 0		
Lower left segment	(6-5) 0, (5-4) 0, (4-3) 2	(6-5) 0, (5-4) 0, (4-3) 0		
Right buccal occlusion	AP 0	AP 0		
	T 0	V 0		
	V 2	T 0		
Overjet	OJ 0	OJ 0		
	x-bite 4	x-bite 0		
Overbite	openbite 0	openbite 0		
	OB 3	OB 0		
Centerline	2	0		
Left buccal occlusion	AP 0	AP 0		
	V 0	V 0		
	T 0	T 0		
<b>Total</b>	<b>30</b>	<b>1</b>	<b>29</b>	<b>96.67</b>

**Table D9** Case No. 9

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 0, (5-4) 1, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Upper anterior segment	(3-2) 1, (2-1) 0, (1-1) 3 (1-2) 5, (2-3) 1	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 1		
Upper left segment	(6-5) 0, (5-4) 0, (4-3) 2	(6-5) 0, (5-4) 0, (4-3) 1		
Lower right segment	(6-5) 0, (5-4) 0, (4-3) 2	(6-5) 0, (5-4) 0, (4-3) 0		
Lower anterior segment	(3-2) 0, (2-1) 1, (1-1) 0 (1-2) 1, (2-3) 1	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5) 0, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Right buccal occlusion	AP 0 T 0 V 2	AP 1 V 0 T 0		
Overjet	OJ 0 x-bite 4	OJ 0 x-bite 0		
Overbite	openbite 0 OB 2	openbite 0 OB 0		
Centerline	2	1		
Left buccal occlusion	AP 1 V 0 T 0	AP 1 V 0 T 0		
<b>Total</b>	<b>29</b>	<b>5</b>	<b>24</b>	<b>82.76</b>

**Table D10 Case No. 10**

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) <b>0</b> , (5-4) <b>1</b> , (4-3) <b>0</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Upper anterior segment	(3-2) <b>1</b> , (2-1) <b>2</b> , (1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>		(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>			
Upper left segment	(6-5) <b>1</b> , (5-4) <b>2</b> , (4-3) <b>1</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Lower right segment	(6-5) <b>5</b> , (5-4) <b>3</b> , (4-3) <b>0</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Lower anterior segment	(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>		(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>			
Lower left segment	(6-5) <b>3</b> , (5-4) <b>0</b> , (4-3) <b>2</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Right buccal occlusion	AP <b>0</b> T <b>0</b> V <b>0</b>		AP <b>0</b> V <b>0</b> T <b>0</b>			
Overjet	OJ <b>0</b> x-bite <b>4</b>		OJ <b>0</b> x-bite <b>0</b>			
Overbite	openbite <b>0</b> OB <b>1</b>		openbite <b>0</b> OB <b>0</b>			
Centerline	<b>0</b>		<b>0</b>			
Left buccal occlusion	AP <b>0</b> V <b>0</b> T <b>2</b>		AP <b>1</b> V <b>0</b> T <b>0</b>			
<b>Total</b>	<b>28</b>		<b>1</b>		<b>27</b>	<b>96.43</b>

**Table D11 Case No. 11**

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) 1, (5-4) 0, (4-3) 0		(6-5) 0, (5-4) 0, (4-3) 0			
Upper anterior segment	(3-2) 1, (2-1) 0, (1-1) 0 (1-2) 1, (2-3) 1		(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0			
Upper left segment	(6-5) 0, (5-4) 1, (4-3) 0		(6-5) 0, (5-4) 0, (4-3) 0			
Lower right segment	(6-5) 0, (5-4) 0, (4-3) 1		(6-5) 0, (5-4) 0, (4-3) 0			
Lower anterior segment	(3-2) 1, (2-1) 0, (1-1) 2 (1-2) 0, (2-3) 1		(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0			
Lower left segment	(6-5) 1, (5-4) 1, (4-3) 2		(6-5) 0, (5-4) 0, (4-3) 0			
Right buccal occlusion	AP 0 T 0 V 0		AP 0 V 0 T 0			
Overjet	OJ 0 x-bite 4		OJ 0 x-bite 0			
Overbite	openbite 1 OB 0		openbite 0 OB 0			
Centerline	1		0			
Left buccal occlusion	AP 0 V 0 T 0		AP 0 V 0 T 0			
<b>Total</b>	<b>20</b>		<b>1</b>	<b>19</b>	<b>95.00</b>	

**Table D12 Case No. 12**

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) 0, (5-4) 1, (4-3) 2		(6-5) 0, (5-4) 0, (4-3) 0			
Upper anterior segment	(3-2) 0, (2-1) 2, (1-1) 0 (1-2) 0, (2-3) 0		(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0			
Upper left segment	(6-5) 1, (5-4) 1, (4-3) 2		(6-5) 0, (5-4) 0, (4-3) 0			
Lower right segment	(6-5) 1, (5-4) 2, (4-3) 0		(6-5) 1, (5-4) 0, (4-3) 0			
Lower anterior segment	(3-2) 1, (2-1) 1, (1-1) 0 (1-2) 0, (2-3) 1		(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0			
Lower left segment	(6-5) 0, (5-4) 2, (4-3) 2		(6-5) 1, (5-4) 0, (4-3) 0			
Right buccal occlusion	AP 0 T 0 V 0		AP 0 V 0 T 0			
Overjet	OJ 0 x-bite 1		OJ 0 x-bite 0			
Overbite	openbite 0 OB 2		openbite 0 OB 0			
Centerline	1		0			
Left buccal occlusion	AP 2 V 0 T 4		AP 0 V 0 T 0			
<b>Total</b>	<b>29</b>		<b>2</b>		<b>27</b>	<b>93.10</b>

**Table D13 Case No. 13**

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 3, (5-4) 3, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Upper anterior segment	(3-2) 0, (2-1) 0, (1-1) 5 (1-2) 5, (2-3) 5	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Upper left segment	(6-5) 0, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Lower right segment	(6-5) 5, (5-4) 2, (4-3) 2	(6-5) 0, (5-4) 0, (4-3) 0		
Lower anterior segment	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5) 0, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Right buccal occlusion	AP 2 T 0 V 4	AP 2 V 0 T 0		
Overjet	OJ 3 x-bite 0	OJ 0 x-bite 0		
Overbite	openbite 1 OB 2	openbite 0 OB 0		
Centerline	2	1		
Left buccal occlusion	AP 2 V 0 T 1	AP 1 V 0 T 0		
<b>Total</b>	<b>45</b>	<b>4</b>	<b>41</b>	<b>91.11</b>

**Table D14** Case No. 14

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 1, (5-4) 2, (4-3) 1	(6-5) 0, (5-4) 1, (4-3) 0		
Upper anterior segment	(3-2) 0, (2-1) 0, (1-1) 2 (1-2) 4, (2-3) 0	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 2, (2-3) 0		
Upper left segment	(6-5) 1, (5-4) 2, (4-3) 1	(6-5) 0, (5-4) 0, (4-3) 0		
Lower right segment	(6-5) 2, (5-4) 2, (4-3) 0	(6-5) 2, (5-4) 2, (4-3) 0		
Lower anterior segment	(3-2) 0, (2-1) 0, (1-1) 1 (1-2) 0, (2-3) 0	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5) 1, (5-4) 0, (4-3) 1	(6-5) 0, (5-4) 0, (4-3) 0		
Right buccal occlusion	AP 2 T 0 V 0	AP 1 V 0 T 0		
Overjet	OJ 0 x-bite 0	OJ 0 x-bite 0		
Overbite	openbite 0 OB 3	openbite 0 OB 2		
Centerline	2	0		
Left buccal occlusion	AP 0 V 0 T 0	AP 0 V 0 T 0		
<b>Total</b>	<b>26</b>	<b>10</b>	<b>16</b>	<b>61.54</b>

**Table D15** Case No. 15

PAR components	Pre-treatment		Post-treatment		Improvement	
	scores		scores		scores	%
Upper right segment	(6-5) 1, (5-4) 0, (4-3) 1		(6-5) 0, (5-4) 0, (4-3) 0			
Upper anterior segment	(3-2) 0, (2-1) 0, (1-1) 2		(3-2) 0, (2-1) 0, (1-1) 0			
	(1-2) 2, (2-3) 2		(1-2) 0, (2-3) 0			
Upper left segment	(6-5) 0, (5-4) 0, (4-3) 1		(6-5) 0, (5-4) 0, (4-3) 0			
Lower right segment	(6-5) 0, (5-4) 0, (4-3) 0		(6-5) 0, (5-4) 0, (4-3) 0			
Lower anterior segment	(3-2) 2, (2-1) 2, (1-1) 0		(3-2) 0, (2-1) 0, (1-1) 0			
	(1-2) 0, (2-3) 2		(1-2) 0, (2-3) 0			
Lower left segment	(6-5) 1, (5-4) 2, (4-3) 0		(6-5) 0, (5-4) 0, (4-3) 0			
Right buccal occlusion	AP 0		AP 2			
	T 0		V 0			
	V 0		T 4			
Overjet	OJ 0		OJ 0			
	x-bite 3		x-bite 0			
Overbite	openbite 1		openbite 0			
	OB 0		OB 0			
Centerline	1		0			
Left buccal occlusion	AP 0		AP 2			
	V 0		V 0			
	T 0		T 0			
<b>Total</b>	<b>23</b>		<b>8</b>		<b>15</b>	<b>65.22</b>

**Table D16 Case No. 16**

PAR components	Pre-treatment		Post-treatment		Improvement	
	scores		scores		scores	%
Upper right segment	(6-5) 2, (5-4) 1, (4-3) 2		(6-5) 0, (5-4) 0, (4-3) 0			
Upper anterior segment	(3-2) 2, (2-1) 1, (1-1) 0		(3-2) 0, (2-1) 0, (1-1) 0			
	(1-2) 0, (2-3) 3		(1-2) 0, (2-3) 0			
Upper left segment	(6-5) 0, (5-4) 0, (4-3) 2		(6-5) 0, (5-4) 0, (4-3) 0			
Lower right segment	(6-5) 0, (5-4) 0, (4-3) 1		(6-5) 0, (5-4) 0, (4-3) 0			
Lower anterior segment	(3-2) 0, (2-1) 0, (1-1) 0		(3-2) 0, (2-1) 0, (1-1) 0			
	(1-2) 0, (2-3) 0		(1-2) 0, (2-3) 0			
Lower left segment	(6-5) 2, (5-4) 1, (4-3) 0		(6-5) 0, (5-4) 1, (4-3) 0			
Right buccal occlusion	AP 2		AP 2			
	T 0		V 0			
	V 0		T 0			
Overjet	OJ 0		OJ 0			
	x-bite 4		x-bite 0			
Overbite	openbite 0		openbite 0			
	OB 1		OB 0			
Centerline	2		0			
Left buccal occlusion	AP 0		AP 0			
	V 0		V 0			
	T 0		T 1			
<b>Total</b>	<b>26</b>		<b>4</b>		<b>22</b>	<b>84.62</b>

**Table D17 Case No. 17**

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) <b>5</b> , (5-4) <b>2</b> , (4-3) <b>2</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Upper anterior segment	(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b>		(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b>			
	(1-2) <b>0</b> , (2-3) <b>0</b>		(1-2) <b>0</b> , (2-3) <b>0</b>			
Upper left segment	(6-5) <b>2</b> , (5-4) <b>1</b> , (4-3) <b>2</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Lower right segment	(6-5) <b>2</b> , (5-4) <b>1</b> , (4-3) <b>2</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Lower anterior segment	(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b>		(3-2) <b>0</b> , (2-1) <b>0</b> , (1-1) <b>0</b>			
	(1-2) <b>0</b> , (2-3) <b>0</b>		(1-2) <b>0</b> , (2-3) <b>0</b>			
Lower left segment	(6-5) <b>1</b> , (5-4) <b>0</b> , (4-3) <b>1</b>		(6-5) <b>0</b> , (5-4) <b>0</b> , (4-3) <b>0</b>			
Right buccal occlusion	AP <b>1</b>		AP <b>1</b>			
	T <b>0</b>		V <b>0</b>			
	V <b>3</b>		T <b>0</b>			
Overjet	OJ <b>0</b>		OJ <b>0</b>			
	x-bite <b>4</b>		x-bite <b>0</b>			
Overbite	openbite <b>0</b>		openbite <b>0</b>			
	OB <b>0</b>		OB <b>0</b>			
Centerline		<b>1</b>		<b>1</b>		
Left buccal occlusion	AP <b>0</b>		AP <b>0</b>			
	V <b>0</b>		V <b>0</b>			
	T <b>3</b>		T <b>0</b>			
<b>Total</b>	<b>33</b>		<b>2</b>		<b>31</b>	<b>93.94</b>

**Table D18 Case No. 18**

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) 0, (5-4) 0, (4-3) 2		(6-5) 0, (5-4) 0, (4-3) 0			
Upper anterior segment	(3-2) 0, (2-1) 0, (1-1) 0		(3-2) 0, (2-1) 0, (1-1) 0			
	(1-2) 0, (2-3) 0		(1-2) 0, (2-3) 0			
Upper left segment	(6-5) 0, (5-4) 0, (4-3) 1		(6-5) 0, (5-4) 0, (4-3) 0			
Lower right segment	(6-5) 0, (5-4) 0, (4-3) 0		(6-5) 0, (5-4) 0, (4-3) 0			
Lower anterior segment	(3-2) 0, (2-1) 0, (1-1) 0		(3-2) 0, (2-1) 0, (1-1) 0			
	(1-2) 0, (2-3) 0		(1-2) 0, (2-3) 0			
Lower left segment	(6-5) 3, (5-4) 0, (4-3) 1		(6-5) 0, (5-4) 0, (4-3) 0			
Right buccal occlusion	AP 2		AP 0			
	T 0		V 0			
	V 0		T 1			
Overjet	OJ 0		OJ 0			
	x-bite 4		x-bite 0			
Overbite	openbite 0		openbite 0			
	OB 0		OB 0			
Centerline	0		0			
Left buccal occlusion	AP 0		AP 2			
	V 0		V 0			
	T 0		T 0			
<b>Total</b>	<b>13</b>		<b>3</b>		<b>10</b>	<b>76.92</b>

**Table D19** Case No. 19

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) 0, (5-4) 1, (4-3) 1		(6-5) 0, (5-4) 0, (4-3) 0			
Upper anterior segment	(3-2) 0, (2-1) 1, (1-1) 1 (1-2) 3, (2-3) 5		(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0			
Upper left segment	(6-5) 0, (5-4) 0, (4-3) 2		(6-5) 0, (5-4) 0, (4-3) 0			
Lower right segment	(6-5) 0, (5-4) 2, (4-3) 2		(6-5) 0, (5-4) 0, (4-3) 0			
Lower anterior segment	(3-2) 0, (2-1) 0, (1-1) 1 (1-2) 1, (2-3) 0		(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0			
Lower left segment	(6-5) 0, (5-4) 3, (4-3) 1		(6-5) 0, (5-4) 0, (4-3) 0			
Right buccal occlusion	AP 2 T 0 V 0		AP 2 V 0 T 0			
Overjet	OJ 0 x-bite 3		OJ 0 x-bite 0			
Overbite	openbite 0 OB 2		openbite 0 OB 0			
Centerline	0		0			
Left buccal occlusion	AP 2 V 0 T 3		AP 0 V 0 T 0			
<b>Total</b>	<b>36</b>		<b>2</b>		<b>34</b>	<b>94.44</b>

**Table D20** Case No. 20

<b>PAR components</b>	<b>Pre-treatment scores</b>	<b>Post-treatment scores</b>	<b>Improvement</b>	
			<b>scores</b>	<b>%</b>
Upper right segment	(6-5) 0, (5-4) 0, (4-3) 2	(6-5) 0, (5-4) 0, (4-3) 0		
Upper anterior segment	(3-2) 3, (2-1) 3, (1-1) 1 (1-2) 0, (2-3) 3	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Upper left segment	(6-5) 1, (5-4) 0, (4-3) 1	(6-5) 0, (5-4) 0, (4-3) 0		
Lower right segment	(6-5) 2, (5-4) 2, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Lower anterior segment	(3-2) 0, (2-1) 0, (1-1) 1 (1-2) 1, (2-3) 1	(3-2) 0, (2-1) 0, (1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5) 1, (5-4) 0, (4-3) 0	(6-5) 0, (5-4) 0, (4-3) 0		
Right buccal occlusion	AP 1 T 0 V 2	AP 1 V 0 T 0		
Overjet	OJ 0 x-bite 4	OJ 0 x-bite 0		
Overbite	openbite 0 OB 2	openbite 0 OB 0		
Centerline	1	0		
Left buccal occlusion	AP 2 V 0 T 3	AP 0 V 0 T 3		
<b>Total</b>	<b>37</b>	<b>4</b>	<b>33</b>	<b>89.19</b>

**Table D21** Case No. 21

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5)0, (5-4) 1, (4-3) 3	(6-5) 0,(5-4)0,(4-3) 0		
Upper anterior segment	(3-2) 1,(2-1) 4, (1-1) 1 (1-2) 0, (2-3) 2	(3-2) 0,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 0		
Upper left segment	(6-5) 1, (5-4) 1, (4-3) 1	(6-5)0,(5-4) 0,(4-3) 0		
Lower right segment	(6-5) 0, (5-4)2, (4-3) 2	(6-5) 0,(5-4)0,(4-3) 0		
Lower anterior segment	(3-2) 2, (2-1)1, (1-1) 0 (1-2) 2, (2-3) 0	(3-2) 0,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5)0, (5-4) 0, (4-3) 1	(6-5)0,(5-4) 0,(4-3) 0		
Right buccal occlusion	AP 2 T 0 V 0	AP 2 V 0 T 0		
Overjet	OJ 4 x-bite 0	OJ 0 x-bite 0		
Overbite	openbite 0 OB 0	openbite 0 OB 0		
Centerline	1	0		
Left buccal occlusion	AP 0 V 0 T 0	AP 1 V 0 T 0		
<b>Total</b>	<b>32</b>	<b>3</b>	<b>29</b>	<b>90.63</b>

**Table D22** Case No. 22

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) <b>1</b> ,(5-4) <b>0</b> ,(4-3) <b>2</b>	(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>		
Upper anterior segment	(3-2) <b>2</b> ,(2-1) <b>3</b> ,(1-1) <b>3</b>  (1-2) <b>0</b> , (2-3) <b>0</b>	(3-2) <b>0</b> ,(2-1) <b>0</b> ,(1-1) <b>0</b>  (1-2) <b>0</b> , (2-3) <b>0</b>		
Upper left segment	(6-5) <b>2</b> ,(5-4) <b>0</b> ,(4-3) <b>2</b>	(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>		
Lower right segment	(6-5) <b>1</b> ,(5-4) <b>1</b> ,(4-3) <b>0</b>	(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>		
Lower anterior segment	(3-2) <b>2</b> ,(2-1) <b>1</b> ,(1-1) <b>2</b>  (1-2) <b>1</b> , (2-3) <b>1</b>	(3-2) <b>0</b> ,(2-1) <b>0</b> ,(1-1) <b>0</b>  (1-2) <b>0</b> , (2-3) <b>0</b>		
Lower left segment	(6-5) <b>1</b> ,(5-4) <b>1</b> ,(4-3) <b>1</b>	(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>		
Right buccal occlusion	AP <b>2</b>  T <b>0</b>  V <b>0</b>	AP <b>2</b>  V <b>0</b>  T <b>0</b>		
Overjet	OJ <b>1</b>  x-bite <b>2</b>	OJ <b>0</b>  x-bite <b>0</b>		
Overbite	openbite <b>0</b>  OB <b>1</b>	openbite <b>0</b>  OB <b>0</b>		
Centerline	<b>0</b>	<b>0</b>		
Left buccal occlusion	AP <b>1</b>  V <b>0</b>  T <b>0</b>	AP <b>0</b>  V <b>0</b>  T <b>0</b>		
<b>Total</b>	<b>34</b>	<b>2</b>	<b>32</b>	<b>94.12</b>

**Table D23** Case No. 23

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) 2,(5-4) 0,(4-3) 3		(6-5) 0,(5-4) 0,(4-3) 0			
Upper anterior segment	(3-2) 0,(2-1) 0,(1-1) 2 (1-2) 2, (2-3) 0		(3-2) 0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 0			
Upper left segment	(6-5) 5,(5-4) 2,(4-3) 2		(6-5) 0,(5-4) 0,(4-3) 0			
Lower right segment	(6-5) 0,(5-4) 0,(4-3) 1		(6-5) 0,(5-4) 0,(4-3) 0			
Lower anterior segment	(3-2) 0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 1		(3-2) 0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 0			
Lower left segment	(6-5) 2,(5-4) 2,(4-3) 2		(6-5) 0,(5-4) 0,(4-3) 0			
Right buccal occlusion	AP 1 T 0 V 0		AP 0 V 0 T 0			
Overjet	OJ 0 x-bite 3		OJ 0 x-bite 0			
Overbite	openbite 3 OB 0		openbite 0 OB 0			
Centerline	0		0			
Left buccal occlusion	AP 0 V 0 T 2		AP 1 V 0 T 0			
<b>Total</b>	<b>35</b>		<b>1</b>		<b>34</b>	<b>97.14</b>

**Table D24** Case No. 24

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 0,(5-4)0,(4-3) 2	(6-5) 0,(5-4)0,(4-3) 0		
Upper anterior segment	(3-2) 0,(2-1)2,(1-1) 3	(3-2) 0,(2-1)0,(1-1) 0		
	(1-2) 5, (2-3) 4	(1-2) 0, (2-3) 0		
Upper left segment	(6-5) 0,(5-4)0,(4-3) 3	(6-5) 0,(5-4)0,(4-3) 0		
Lower right segment	(6-5)0,(5-4) 0,(4-3) 0	(6-5) 0,(5-4)0,(4-3) 0		
Lower anterior segment	(3-2)1,(2-1) 0,(1-1) 1	(3-2) 0,(2-1)0,(1-1) 0		
	(1-2) 1, (2-3) 1	(1-2) 0, (2-3) 0		
Lower left segment	(6-5)0,(5-4) 0,(4-3) 0	(6-5) 0,(5-4)0,(4-3) 0		
Right buccal occlusion	AP 2	AP 0		
	T 0	V 0		
	V 0	T 0		
Overjet	OJ 0	OJ 0		
	x-bite 4	x-bite 1		
Overbite	openbite 1	openbite 0		
	OB 0	OB 0		
Centerline	0	0		
Left buccal occlusion	AP 1	AP 0		
	V 0	V 0		
	T 2	T 0		
<b>Total</b>	<b>33</b>	<b>1</b>	<b>32</b>	<b>96.97</b>

**Table D25** Case No. 25

PAR components	Pre-treatment		Post-treatment		Improvement	
	scores		scores		scores	%
Upper right segment	(6-5) <b>5</b> ,(5-4) <b>2</b> ,(4-3) <b>0</b>		(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>			
Upper anterior segment	(3-2) <b>0</b> ,(2-1) <b>1</b> ,(1-1) <b>0</b> (1-2) <b>2</b> , (2-3) <b>0</b>		(3-2) <b>0</b> ,(2-1) <b>0</b> ,(1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>			
Upper left segment	(6-5) <b>0</b> ,(5-4) <b>2</b> ,(4-3) <b>4</b>		(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>			
Lower right segment	(6-5) <b>1</b> ,(5-4) <b>0</b> ,(4-3) <b>3</b>		(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>			
Lower anterior segment	(3-2) <b>2</b> ,(2-1) <b>1</b> ,(1-1) <b>0</b> (1-2) <b>2</b> , (2-3) <b>2</b>		(3-2) <b>0</b> ,(2-1) <b>0</b> ,(1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>			
Lower left segment	(6-5) <b>2</b> ,(5-4) <b>0</b> ,(4-3) <b>3</b>		(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>			
Right buccal occlusion	AP <b>1</b> T <b>0</b> V <b>0</b>		AP <b>0</b> V <b>0</b> T <b>0</b>			
Overjet	OJ <b>0</b> x-bite <b>4</b>		OJ <b>0</b> x-bite <b>0</b>			
Overbite	openbite <b>0</b> OB <b>3</b>		openbite <b>0</b> OB <b>0</b>			
Centerline		<b>1</b>		<b>0</b>		
Left buccal occlusion	AP <b>1</b> V <b>0</b> T <b>3</b>		AP <b>0</b> V <b>0</b> T <b>0</b>			
<b>Total</b>	<b>45</b>		<b>0</b>		<b>45</b>	<b>100</b>

**Table D26** Case No. 26

PAR components	Pre-treatment		Post-treatment		Improvement	
	scores		scores		scores	%
Upper right segment	(6-5) 2,(5-4) 0,(4-3) 3		(6-5) 0,(5-4) 0,(4-3) 0			
Upper anterior segment	(3-2) 4,(2-1) 0,(1-1) 3 (1-2) 3, (2-3) 3		(3-2) 0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 0			
Upper left segment	(6-5) 2,(5-4) 0,(4-3) 3		(6-5) 1,(5-4) 0,(4-3) 0			
Lower right segment	(6-5) 0,(5-4) 0,(4-3) 1		(6-5) 0,(5-4) 0,(4-3) 0			
Lower anterior segment	(3-2) 0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 2		(3-2) 0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 0			
Lower left segment	(6-5) 2,(5-4) 0,(4-3) 1		(6-5) 0,(5-4) 0,(4-3) 0			
Right buccal occlusion	AP 1 T 0 V 0		AP 0 V 0 T 0			
Overjet	OJ 4 x-bite 0		OJ 0 x-bite 0			
Overbite	openbite -4 OB 0		openbite 1 OB 0			
Centerline	1		0			
Left buccal occlusion	AP 1 V 0 T 0		AP 0 V 0 T 0			
<b>Total</b>	<b>40</b>		<b>2</b>		<b>38</b>	<b>95</b>

**Table D27** Case No. 27

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 5,(5-4)2,(4-3) 3	(6-5)0,(5-4)0,(4-3) 0		
Upper anterior segment	(3-2) 2,(2-1)1,(1-1) 0 (1-2) 0, (2-3) 0	(3-2) 0,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 1		
Upper left segment	(6-5) 5,(5-4)2,(4-3) 0	(6-5)0,(5-4) 0,(4-3) 0		
Lower right segment	(6-5)5,(5-4) 0,(4-3) 0	(6-5) 0,(5-4)0,(4-3) 0		
Lower anterior segment	(3-2) 0,(2-1)1,(1-1) 0 (1-2) 3, (2-3) 2	(3-2) 0,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5)5,(5-4) 0,(4-3) 0	(6-5) 0,(5-4)0,(4-3) 0		
Right buccal occlusion	AP 0 T 0 V 0	AP 1 V 0 T 0		
Overjet	OJ 0 x-bite 3	OJ 0 x-bite 0		
Overbite	openbite 0 OB 2	openbite 0 OB 0		
Centerline	1	0		
Left buccal occlusion	AP 0 V 0 T 0	AP 1 V 0 T 0		
<b>Total</b>	<b>42</b>	<b>3</b>	<b>39</b>	<b>92.86</b>

## PAR Index Score of Combined Orthognathic Surgery Group

**Table D28** Case No. 28

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) <b>1</b> ,(5-4)0,(4-3) <b>0</b>	(6-5) <b>0</b> ,(5-4)0,(4-3) <b>0</b>		
Upper anterior segment	(3-2) <b>5</b> ,(2-1) <b>0</b> ,(1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>	(3-2) <b>0</b> ,(2-1)0,(1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>		
Upper left segment	(6-5) <b>0</b> ,(5-4)0,(4-3) <b>0</b>	(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>		
Lower right segment	(6-5) <b>1</b> ,(5-4) <b>2</b> ,(4-3) <b>1</b>	(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>		
Lower anterior segment	(3-2) <b>0</b> ,(2-1) <b>1</b> ,(1-1) <b>0</b> (1-2) <b>1</b> , (2-3) <b>1</b>	(3-2) <b>0</b> ,(2-1)0,(1-1) <b>0</b> (1-2) <b>0</b> , (2-3) <b>0</b>		
Lower left segment	(6-5) <b>0</b> ,(5-4) <b>2</b> ,(4-3) <b>1</b>	(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>		
Right buccal occlusion	AP <b>2</b> T <b>0</b> V <b>3</b>	AP <b>0</b> V <b>0</b> T <b>0</b>		
Overjet	OJ <b>0</b> x-bite <b>4</b>	OJ <b>0</b> x-bite <b>0</b>		
Overbite	openbite <b>0</b> OB <b>0</b>	openbite <b>0</b> OB <b>0</b>		
Centerline			<b>0</b>	
Left buccal occlusion	AP <b>2</b> V <b>0</b> T <b>3</b>	AP <b>1</b> V <b>0</b> T <b>0</b>		
<b>Total</b>	<b>30</b>	<b>1</b>	<b>29</b>	<b>96.67</b>

**Table D29** Case No. 29

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) 2,(5-4)2,(4-3) 0		(6-5) 0,(5-4)0,(4-3) 0			
Upper anterior segment	(3-2)2,(2-1) 0,(1-1) 0 (1-2) 5, (2-3) 0		(3-2) 1,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 0			
Upper left segment	(6-5) 5,(5-4)1,(4-3) 1		(6-5) 0,(5-4)0,(4-3) 0			
Lower right segment	(6-5) 0,(5-4)1,(4-3) 0		(6-5) 0,(5-4)0,(4-3) 0			
Lower anterior segment	(3-2) 1,(2-1)1,(1-1) 0 (1-2) 1, (2-3) 0		(3-2) 0,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 0			
Lower left segment	(6-5) 0,(5-4)1,(4-3) 0		(6-5) 0,(5-4)0,(4-3) 0			
Right buccal occlusion	AP 2 T 0 V 3		AP 0 V 0 T 0			
Overjet	OJ 0 x-bite 4		OJ 1 x-bite 0			
Overbite	openbite 0 OB 3		openbite 0 OB 0			
Centerline		2		1		
Left buccal occlusion	AP 2 V 0 T 3		AP 0 V 0 T 0			
<b>Total</b>	<b>42</b>		<b>3</b>		<b>39</b>	<b>92.86</b>

**Table D30** Case No. 30

PAR components	Pre-treatment		Post-treatment		Improvement	
	scores		scores		scores	%
Upper right segment	(6-5) <b>4</b> ,(5-4) <b>0</b> ,(4-3) <b>3</b>		(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>			
Upper anterior segment	(3-2)3,(2-1) <b>0</b> ,(1-1) <b>2</b>		(3-2) <b>0</b> ,(2-1) <b>0</b> ,(1-1) <b>0</b>			
	(1-2) <b>4</b> , (2-3) <b>0</b>		(1-2) <b>0</b> , (2-3) <b>0</b>			
Upper left segment	(6-5)3,(5-4) <b>0</b> ,(4-3) <b>1</b>		(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>			
Lower right segment	(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>		(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>			
Lower anterior segment	(3-2) <b>2</b> ,(2-1) <b>1</b> ,(1-1) <b>0</b>		(3-2) <b>0</b> ,(2-1) <b>0</b> ,(1-1) <b>0</b>			
	(1-2) <b>0</b> , (2-3) <b>1</b>		(1-2) <b>0</b> , (2-3) <b>0</b>			
Lower left segment	(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>		(6-5) <b>0</b> ,(5-4) <b>0</b> ,(4-3) <b>0</b>			
Right buccal occlusion	AP <b>2</b>		AP <b>0</b>			
	T <b>0</b>		V <b>0</b>			
	V <b>3</b>		T <b>1</b>			
Overjet	OJ <b>0</b>		OJ <b>0</b>			
	x-bite <b>4</b>		x-bite <b>1</b>			
Overbite	openbite <b>0</b>		openbite <b>0</b>			
	OB <b>3</b>		OB <b>0</b>			
Centerline		<b>1</b>		<b>0</b>		
Left buccal occlusion	AP <b>2</b>		AP <b>1</b>			
	V <b>0</b>		V <b>0</b>			
	T <b>3</b>		T <b>2</b>			
<b>Total</b>		<b>42</b>		<b>5</b>	<b>37</b>	<b>88.10</b>

**Table D31** Case No. 31

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5) 0,(5-4)0,(4-3) 3		(6-5) 0,(5-4)0,(4-3) 0			
Upper anterior segment	(3-2)0,(2-1) 1,(1-1) 0 (1-2) 1, (2-3) 1		(3-2) 0,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 0			
Upper left segment	(6-5) 2,(5-4)3,(4-3) 0		(6-5) 0,(5-4)0,(4-3) 0			
Lower right segment	(6-5) 1,(5-4)0,(4-3) 0		(6-5) 0,(5-4)0,(4-3) 0			
Lower anterior segment	(3-2) 2,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 2		(3-2) 0,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 0			
Lower left segment	(6-5) 0,(5-4)2,(4-3) 0		(6-5) 1,(5-4)0,(4-3) 0			
Right buccal occlusion	AP 1 T 0 V 0		AP 0 V 0 T 0			
Overjet	OJ 0 x-bite 1		OJ 0 x-bite 0			
Overbite	openbite 0 OB 0		openbite 0 OB 0			
Centerline	1		0			
Left buccal occlusion	AP 0 V 0 T 3		AP 0 V 0 T 0			
<b>Total</b>	<b>24</b>		<b>1</b>		<b>23</b>	<b>95.83</b>

**Table D32** Case No. 32

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 0,(5-4)5,(4-3) 3	(6-5) 0,(5-4)0,(4-3) 0		
Upper anterior segment	(3-2) 3,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 0	(3-2) 0,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 0		
Upper left segment	(6-5) 5,(5-4)2,(4-3) 0	(6-5) 0,(5-4)0,(4-3) 0		
Lower right segment	(6-5) 2,(5-4)3,(4-3) 0	(6-5) 0,(5-4)0,(4-3) 0		
Lower anterior segment	(3-2)0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 2	(3-2) 0,(2-1)0,(1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5)3,(5-4) 1,(4-3) 0	(6-5) 1,(5-4)0,(4-3) 0		
Right buccal occlusion	AP 2 T 0 V 3	AP 0 V 0 T 0		
Overjet	OJ 0 x-bite 4	OJ 0 x-bite 0		
Overbite	openbite 0 OB 0	openbite 0 OB 0		
Centerline	1	1		
Left buccal occlusion	AP 2 V 0 T 3	AP 0 V 0 T 0		
<b>Total</b>	<b>44</b>	<b>3</b>	<b>41</b>	<b>93.18</b>

**Table D33** Case No. 33

PAR components	Pre-treatment scores		Post-treatment scores		Improvement	
					scores	%
Upper right segment	(6-5)2,(5-4)2,(4-3) 1		(6-5) 0,(5-4)0,(4-3) 0			
Upper anterior segment	(3-2) 5,(2-1)0,(1-1) 2 (1-2) 3, (2-3) 0		(3-2)0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 0			
Upper left segment	(6-5)5,(5-4) 2,(4-3) 2		(6-5) 0,(5-4)0,(4-3) 0			
Lower right segment	(6-5)2,(5-4) 1,(4-3) 1		(6-5) 0,(5-4)0,(4-3) 0			
Lower anterior segment	(3-2)0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 1		(3-2)0,(2-1)0, (1-1) 0 (1-2) 0, (2-3) 0			
Lower left segment	(6-5)0,(5-4) 2,(4-3) 1		(6-5)0,(5-4) 0,(4-3) 0			
Right buccal occlusion	AP 2 T 0 V 3		AP 0 V 0 T 1			
Overjet	OJ 0 x-bite 4		OJ 0 x-bite 0			
Overbite	openbite 0 OB 2		openbite 0 OB 0			
Centerline	2		0			
Left buccal occlusion	AP 1 V 0 T 3		AP 0 V 0 T 0			
<b>Total</b>	<b>49</b>		<b>1</b>		<b>48</b>	<b>97.96</b>

**Table D34** Case No. 34

PAR components	Pre-treatment scores	Post-treatment scores	Improvement	
			scores	%
Upper right segment	(6-5) 5,(5-4) 2,(4-3) 2	(6-5) 0,(5-4) 0,(4-3) 0		
Upper anterior segment	(3-2) 3,(2-1) 0,(1-1) 2 (1-2) 3, (2-3) 3	(3-2) 0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 0		
Upper left segment	(6-5) 5,(5-4) 1,(4-3) 0	(6-5) 0,(5-4) 0,(4-3) 0		
Lower right segment	(6-5) 0,(5-4) 2,(4-3) 3	(6-5) 0,(5-4) 0,(4-3) 0		
Lower anterior segment	(3-2) 0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 1	(3-2) 0,(2-1) 0,(1-1) 0 (1-2) 0, (2-3) 0		
Lower left segment	(6-5) 0,(5-4) 2,(4-3) 1	(6-5) 0,(5-4) 0,(4-3) 0		
Right buccal occlusion	AP 2 T 0 V 2	AP 1 V 0 T 0		
Overjet	OJ 0 x-bite 4	OJ 0 x-bite 0		
Overbite	openbite 0 OB 0	openbite 1 OB 0		
Centerline	1	0		
Left buccal occlusion	AP 2 V 0 T 2	AP 1 V 0 T 0		
<b>Total</b>	<b>48</b>	<b>3</b>	<b>45</b>	<b>93.75</b>

**APPENDIX E**  
**Raw Data for PAR Index Score Improvement Categories**

**Table E1** PAR index score improvement categories

No.	improvement score (%)	improvement catagories
1	33 (100.00)	1
2	39 (92.86)	2
3	19 (86.36)	2
4	35 (97.22)	2
5	23 (95.83)	2
6	32 (88.89)	2
7	41 (97.62)	2
8	29 (96.67)	2
9	24 (82.76)	2
10	27 (96.43)	2
11	19 (95.00)	2
12	27 (93.10)	2
13	41 (91.11)	2
14	16 (61.54)	2
15	15 (65.22)	2
16	22 (84.62)	2
17	31 (93.94)	2
18	10 (76.92)	2
19	34 (94.44)	2
20	33 (89.19)	2
21	29 (90.63)	2
22	32 (94.12)	2
23	34 (97.14)	2
24	32 (96.97)	2
25	45 (100.00)	1
26	38 (95.00)	2
27	39 (92.86)	2
28	29 (96.67)	2
29	39 (92.86)	2
30	37 (88.10)	2
31	23 (95.83)	2
32	41 (93.18)	2
33	48 (97.96)	2
34	45 (93.75)	2

**APPENDIX F**  
**Ethical Committee Approval Documents**



**มหาวิทยาลัยขอนแก่น  
หนังสือบันทึกไว้เพื่อแสดงว่า**

**โครงการวิจัยเรื่อง:** การประเมินผลการรักษาทางทันตกรรมจัดฟันขั้นสุดท้ายในผู้ป่วยปากแหว่งเพดานโหว์ ที่ได้รับการรักษาโดยการจัดฟันเพียงอย่างเดียวและที่ได้รับการรักษาโดยการจัดฟันร่วมกับการผ่าตัดขากรรไกรโดยดัชนีชี้วัดพาร์

**EVALUATION OF THE FINAL PHASE TREATMENT OUTCOMES OF DENTO-SKELETAL PROBLEMS IN COMPLETE CLEFT LIP AND PALATE PATIENTS WITH AND WITHOUT SURGERY USING THE PEER ASSESSMENT RATING (PAR) INDEX**

**ผู้วิจัย:**

1. นางสาวศุทธินันท์ ค่าวนรัตนกุล  
คณะทันตแพทยศาสตร์ มหาวิทยาลัยขอนแก่น
2. รองศาสตราจารย์ทันตแพทย์หญิงทักษิณี วงศ์ริมวงศ์  
คณะทันตแพทยศาสตร์ มหาวิทยาลัยขอนแก่น
3. รองศาสตราจารย์ทันตแพทย์มนเทียร โนนศุลประสิทธิ์  
คณะทันตแพทยศาสตร์ มหาวิทยาลัยขอนแก่น

**สำหรับเอกสาร:**

1. เอกสาร ขอปรับปรุงแก้ไขโครงการวิจัยที่ผ่านการรับรองจากคณะกรรมการจัดการวิจัยในมนุษย์ มหาวิทยาลัยขอนแก่น ฉบับที่ ศธ 0514.13.3/31 ลงวันที่ 21 มกราคม พ.ศ. 2554
2. แบบรายงานการแก้ไขปรับปรุงโครงการวิจัย ต่อคณะกรรมการจัดการวิจัยในมนุษย์ มหาวิทยาลัยขอนแก่น ลงวันที่ 21 มกราคม พ.ศ. 2554
3. ประวัติ และความชำนาญของนักวิจัย
  - ศ.ดร. มนเทียร โนนศุลประสิทธิ์
  - ศ.ดร. พญ. คร. ทักษิณี วงศ์ริมวงศ์

ลักษณะที่ 4.6.04: 05/2554

เลขที่: HE 532082

วันที่มาตถูก: 14 มีนาคม พ.ศ.2554

คณะกรรมการจัดการวิจัยในมนุษย์มหาวิทยาลัยขอนแก่น

สำนักงาน: อาคารสนับสนุนทางวิจัย บรมราชชนนี อุบลราชธานี (ชั้น 17)

โทรศัพท์: (043) 366616, (043) 366617 โทรสาร: (043) 366617

Institutional Review Board Number: IRB00001189

Federal Wide Assurance: FWA00003418

ได้ผ่านการรับรองจากคณะกรรมการจริยธรรมการวิจัยในมนุษย์นิหารวิทยาลักษณ์แก่น โดยยึดหลักเกณฑ์ตาม  
คำประกาศเซลลินกี (Declaration of Helsinki) และแนวทางการปฏิบัติการวิจัยทางคลินิกที่ดี (ICH GCP)

ให้ไว้ ณ วันที่ 7 กุมภาพันธ์ พ.ศ. 2554

(นายแพพต์เกรียงศักดิ์ เวทีกุณาจารย์)

ประธานคณะกรรมการจริยธรรมการวิจัยในมนุษย์นิหารวิทยาลักษณ์แก่น

เดือนที่ 4.6.04: 05/2554

เลขที่: HE 532082

วันหมดอายุ: 14 มีนาคม พ.ศ.2554

คณะกรรมการจริยธรรมการวิจัยในมนุษย์นิหารวิทยาลักษณ์แก่น

Institutional Review Board Number; IRB00001189

สำนักงาน: สถาบันสหคุณภาพคุณภาพนานาชาติ อนุสรณ์ (ชั้น 17)

Federal Wide Assurance; FWA00003418

โทร. (043) 366616, (043) 366617 โทรสาร (043) 366617

## **RESEARCH PUBLICATIONS**

1. **Danthumrongkul S**, Manosudprasit M, Wangsrimongkol T. Evaluation of the final phase treatment outcomes of dento-skeletal problems in complete cleft lip and palate patients with and without surgery using the Peer Assessment Rating (PAR) Index. Proceedings of the 12<sup>th</sup> graduate research conference; 2011 Jan 28<sup>th</sup>; Khon Kaen, Thailand. Khon Kaen University.
2. **Danthumrongkul S**, Manosudprasit M, Wangsrimongkol T. Evaluation of the final phase treatment outcomes of dento-skeletal problems in complete cleft lip and palate patients with and without surgery using the Peer Assessment Rating (PAR) Index. Final programme booklet of the 87<sup>th</sup> Congress of the European Orthodontic Society; 2011 Jun 19<sup>th</sup>-23<sup>rd</sup>; Istanbul, Turkey. European Orthodontic Society.

## VITAE



Name: Miss Suthinun Danthumrongkul.

Date of Birth: February 29<sup>th</sup>, 1980

Place of Birth: Bangkok Province, Thailand.

Address: 1/75, Moo 7, Ramindra Road, Tarang, Bangkaen, Bangkok. Thailand 10220.

Education:

1997-2003 Bachelor Degree of Doctor of Dental Surgery.  
(Second Honor), Faculty of Dentistry,  
Chulalongkorn University, Bangkok, Thailand.

2008-2011 Student of the Master of Science in Orthodontics,  
Department of Orthodontics, Faculty of Dentistry,  
Khon Kaen University, Khon Kaen, Thailand.

