

Full Paper

## Some convolution properties of a subclass of $p$ -valent functions

Khalida Inayat Noor and Nazar Khan\*

Department of Mathematics COMSATS, Institute of Information Technology, Park Road Islamabad, Pakistan

\* Corresponding author, e-mail: [nazarmaths@gmail.com](mailto:nazarmaths@gmail.com)

Received: 15 January 2014 / Accepted: 24 June 2015 / Published: 3 July 2015

---

**Abstract:** In this paper we introduce and study a new subclass  $R_p$  of  $p$ -valent functions that are analytic in the open unit disk  $E = \{z: |z| < 1\}$ . Some interesting results by using convolution technique for this subclass  $R_p$  are obtained. Also, we point out some known consequences of our main results.

**Keywords:** analytic functions,  $p$ -valent functions,  $p$ -valently starlike functions,  $p$ -valently convex functions, convolution

---

### INTRODUCTION

Let  $A(p)$  denote the class of functions

$$f(z) = z^p + \sum_{n=1}^{\infty} a_{n+p} z^{n+p} \quad (p \in N = \{1, 2, \dots\}), \quad (1)$$

which are analytic and  $p$ -valent in the open unit disk  $E = \{z: |z| < 1\}$ . A function  $f(z) \in A(p)$  is said to be  $p$ -valently starlike of order  $\alpha$  ( $0 \leq \alpha < p$ ) in  $E$  if and only if

$$\operatorname{Re} \left( \frac{zf'(z)}{f(z)} \right) > \alpha \quad (z \in E).$$

We denote the class of all  $p$ -valent starlike functions of order  $\alpha$  by  $S_p^*(\alpha)$ . Further, a function  $f(z) \in A(p)$  is said to be  $p$ -valently convex of order  $\alpha$  ( $0 \leq \alpha < p$ ) in  $E$  if and only if

$$\operatorname{Re} \left( 1 + \frac{zf''(z)}{f'(z)} \right) > \alpha \quad (z \in E).$$

We denote by  $C_p(\alpha)$  the subclass of  $A(p)$ , consisting of all  $p$ -valently convex functions of order  $\alpha$  in  $E$ . It follows from the definition that if  $f(z)$  is  $p$ -valently convex function, then  $zf'(z)$  is  $p$ -valently starlike in  $E$ . The classes  $S_p^*(\alpha)$  and  $C_p(\alpha)$  were first introduced by Owa [1].

It is easy to see that  $S_p^*(0) = S_p^*$  and  $C_p(0) = C_p$  are, respectively, the classes of  $p$ -valently starlike and  $p$ -valently convex functions in  $E$ . We also note that  $S_p^* = S^*$  and  $C_1 = C$  are, respectively, the well-known classes of starlike and convex functions in  $E$ .

We say that a function

$$h(z) = 1 + \sum_{n=1}^{\infty} c_n z^n \quad (z \in E)$$

belongs to the class  $P(\alpha)$  if  $h(z)$  satisfies the following condition

$$\operatorname{Re} h(z) > \alpha, \quad 0 \leq \alpha < 1 \quad (z \in E).$$

Let  $f(z), g(z) \in A(p)$ , where  $f(z)$  is given by (1) and  $g(z)$  is defined by

$$g(z) = z^p + \sum_{n=1}^{\infty} b_{n+p} z^{n+p} \quad (z \in E).$$

Then the Hadamard product (or convolution)  $f * g$  of the functions  $f(z)$  and  $g(z)$  is defined by

$$(f * g)(z) = z^p + \sum_{n=1}^{\infty} a_{n+p} b_{n+p} z^{n+p} = (g * f)(z).$$

In recent years many interesting subclasses of analytic multivalent functions associated with the linear operator and their many special cases were investigated by, for example Liu [2] and Sokol et al. [3], and also by Arif et al. [4, 5] and others [e.g. 6-8] using convolution technique.

We now define the following:

**Definition 1.** A function  $f(z)$  given by (1) is said to belong to the class  $R_p$  if

$$\operatorname{Re} \left( \frac{f^{(p)}(z) + z f^{(p+1)}(z)}{p!} \right) > 0 \quad (z \in E, p \in N = \{1, 2, \dots\}), \quad (2)$$

where  $f^{(p)}(z)$  is the  $p$ th derivative of  $f(z)$ .

As a special case, the class  $R_1 = R$  was studied by Singh and Singh [9] in 1989. Using essentially their technique and that of Lashin [10], we prove the main results for the class  $R_p$ , which is the main motivation of this paper.

## PRELIMINARY RESULTS

**Lemma 1** [11]. Let  $\{d_n\}_0^{\infty}$  be a convex null sequence. Then the function

$$q(z) = \frac{d_0}{2} + \sum_{n=1}^{\infty} d_n z^n$$

is analytic in  $E$  and  $\operatorname{Re} q(z) > 0 \quad (z \in E)$ .

**Lemma 2** [12]. If  $N(z)$  and  $D(z)$  are analytic in  $E$ ,  $N(0) = D(0) = 0$ ,  $D(z)$  is starlike in  $E$  and

$$\operatorname{Re} \left( \frac{N'(z)}{D'(z)} \right) > 0, \quad \text{then} \quad \operatorname{Re} \left( \frac{N(z)}{D(z)} \right) > 0 \quad (z \in E).$$

**Lemma 3** [9]. If  $h(z)$  is analytic in  $E$ ,  $h(0) = 1$  and  $\operatorname{Re} h(z) > \frac{1}{2} \quad (z \in E)$ , then for any function  $F$  analytic in  $E$ , the function  $h * F$  takes values in the convex hull of the image of  $E$  under  $F$ .

**Lemma 4** [13]. Let  $\beta < 1$ . If the function  $h(z)$  is analytic in  $E$ , with  $h(0) = 1$ , and

$$\operatorname{Re} (h(z) + z h'(z)) > \beta \quad (z \in E).$$

Then

$$\operatorname{Re} h(z) > (2\beta - 1) + 2(1 - \beta) \ln 2 \quad (z \in E).$$

The result is sharp.

**Lemma 5** [14]. For  $\alpha \leq 1$  and  $\beta \leq 1$ ,

$$P(\alpha) * P(\beta) \subset P(\delta), \quad \delta = 1 - 2(1 - \alpha)(1 - \beta).$$

The result is sharp.

## MAIN RESULTS

**Theorem 1.** Let  $f(z) \in R_p$ ; then

$$\operatorname{Re} \left( \frac{f^{(p)}(z)}{p!} \right) > -1 + 2\log 2 \quad (z \in E).$$

The constant  $-1 + 2\log 2$  cannot be replaced by any larger one.

**Proof.** Let  $f(z) \in R_p$ ; then we have

$$\operatorname{Re} \left( \frac{f^{(p)}(z) + z f^{(p+1)}(z)}{p!} \right) > 0 \quad (z \in E),$$

which can be written as

$$\operatorname{Re} \left( 1 + \sum_{n=1}^{\infty} \frac{(p+n)!(n+1)}{p!n!} a_{p+n} z^n \right) > 0 \quad (z \in E), \quad (3)$$

or equivalently,

$$\operatorname{Re} \left( 1 + \frac{1}{2} \sum_{n=1}^{\infty} \frac{(p+n)!(n+1)}{p!n!} a_{p+n} z^n \right) > \frac{1}{2} \quad (z \in E). \quad (4)$$

Consider the function

$$h(z) = 1 + 2 \sum_{n=1}^{\infty} \frac{1}{n+1} z^n. \quad (5)$$

Clearly,  $h(z)$  is analytic in  $E$ ,  $h(0) = 1$  and

$$\begin{aligned} \operatorname{Re} h(z) &= \operatorname{Re} \left( 1 - \frac{2}{z} \{z + \log(1 - z)\} \right) \\ &> -1 + 2\log 2 \quad [15]. \end{aligned} \quad (6)$$

From (4) and (5) we obtain

$$\frac{f^{(p)}(z)}{p!} = \left( 1 + \frac{1}{2} \sum_{n=1}^{\infty} \frac{(p+n)!(n+1)}{p!n!} a_{p+n} z^n \right) * \left( 1 + 2 \sum_{n=1}^{\infty} \frac{1}{n+1} z^n \right),$$

from which it follows, in view of (4), (6) and Lemma 3, that

$$\operatorname{Re} \left( \frac{f^{(p)}(z)}{p!} \right) > -1 + 2\log 2 \quad (z \in E).$$

The constant  $-1 + 2\log 2$  cannot be replaced by any larger one, which follows from the fact that the function  $f_1$  defined by  $\frac{z f_1^{(p)}(z)}{p!} = -z - 2\log(1 - z)$  is in the class  $R_p$ .

**Corollary 1** [9]. If  $f(z) \in R$ , then

$$\operatorname{Re} f'(z) > -1 + 2\log 2 = 0.39... \quad (z \in E).$$

The constant  $-1 + 2\log 2$  cannot be replaced by any larger one.

**Theorem 2.** Let  $f(z) \in R_p$ , then

$$\operatorname{Re} \left( \frac{f^{(p-1)}(z)}{z} \right) > \frac{p!}{2} \quad (z \in E).$$

**Proof.** Since the sequence  $\{d_n\}_0^\infty$  defined by  $d_0 = 1, d_n = \frac{2}{(n+1)^2}, n \geq 1$  is a convex null sequence, using Lemma 1 we have

$$\operatorname{Re} \left( 1 + 2 \sum_{n=1}^{\infty} \frac{1}{(n+1)^2} z^n \right) > \frac{1}{2} \quad (z \in E). \quad (7)$$

We can write

$$\frac{f^{(p-1)}(z)}{p!z} = \left( 1 + \frac{1}{2} \sum_{n=1}^{\infty} \frac{(p+n)!(n+1)}{p!n!} a_{p+n} z^n \right) * \left( 1 + 2 \sum_{n=1}^{\infty} \frac{1}{(n+1)^2} z^n \right).$$

From (4), (7) and Lemma 3, we have the required result.

**Theorem 3.** Let  $f(z) \in R_p$ , then for every  $n \geq 1$ , the  $n^{\text{th}}$  partial sum of  $f(z)$  satisfies  $\operatorname{Re} S_n^{(p)}(z, f) > 0, z \in E$  and hence  $S_n(z, f)$  is  $p$ -valent in  $E$ .

**Proof.** From (3) and (5) we can write

$$\frac{S_n^{(p)}(z, f)}{p!} = \left( 1 + \sum_{n=1}^{\infty} \frac{(p+n)!(n+1)}{p!n!} a_{p+n} z^n \right) * \left( 1 + \sum_{n=1}^{\infty} \frac{1}{(n+1)} z^n \right). \quad (8)$$

Putting  $z = re^{i\theta}$ ,  $0 \leq r \leq 1$ ,  $0 \leq |\theta| \leq \pi$ , and using the minimum principle for harmonic functions with the result in the literature[16], we have

$$\begin{aligned} \operatorname{Re} \left( 1 + \sum_{n=1}^k \frac{z^n}{n+1} \right) &= \operatorname{Re} \left( 1 + \sum_{n=1}^k \frac{r^n e^{in\theta}}{n+1} \right) \\ &= \operatorname{Re} \left( 1 + \sum_{n=1}^k \frac{r^n}{n+1} (\cos n\theta + i \sin n\theta) \right) \\ &= 1 + \sum_{n=1}^k \frac{r^n}{n+1} \cos n\theta \quad (0 \leq \theta \leq \pi) \\ &= 1 + \sum_{n=1}^k \frac{\cos n\theta}{n+1} \geq \frac{1}{2}. \end{aligned} \quad (9)$$

Using (3), (8), (9) and Lemma 3, we deduce that  $\operatorname{Re} (S_n^{(p)}(z, f)) > 0, z \in E$ . From the result given [17], we see that  $S_n(z, f)$  is  $p$ -valent in  $E$  for every  $n \geq 1$ . If  $p = 1$ , then Theorem 2 and Theorem 3 were proved [9].

**Theorem 4.** If  $f(z) \in A(p)$  and

$$\operatorname{Re} \left( \frac{f^{(p)}(z) + z f^{(p+1)}(z)}{p!} \right) > -\frac{1}{4} \quad (z \in E), \quad (10)$$

then  $f(z) \in S_p^*(p-1)$ .

**Proof.** Let  $f(z) \in A(p)$  given by(1). It follows from the hypothesis of the theorem that

$$\operatorname{Re} \left( 1 + \frac{2}{5} \sum_{n=1}^{\infty} \frac{(p+n)!(n+1)}{p!n!} a_{p+n} z^n \right) > \frac{1}{2} \quad (z \in E). \quad (11)$$

Also, the sequence  $\{d_n\}_{n \geq 1}$ , where  $d_0 = 1$  and  $d_n = \frac{5}{2} \frac{1}{(n+1)^2}, n \geq 1$ , is a convex null sequence such as

$$\operatorname{Re} \left( 1 + \frac{5}{2} \sum_{n=1}^{\infty} \frac{1}{(n+1)^2} z^n \right) > \frac{1}{2} \quad (z \in E). \quad (12)$$

From (11), (12) and Lemma 1, we obtain

$$\begin{aligned} \operatorname{Re} \left( \frac{f^{(p-1)}(z)}{p!z} \right) &= \operatorname{Re} \left( 1 + \frac{2}{5} \sum_{n=1}^{\infty} \frac{(p+n)!(n+1)}{p!n!} a_{p+n} z^n \right) * \left( 1 + \frac{5}{2} \sum_{n=1}^{\infty} \frac{1}{(n+1)^2} z^n \right) \\ &> \frac{1}{2} \quad (z \in E). \end{aligned} \quad (13)$$

Now we define a function  $w(z)$  by

$$w(z) = \frac{z f^{(p)}(z) - f^{(p-1)}(z)}{z f^{(p)}(z) + f^{(p-1)}(z)},$$

which can be written as

$$\frac{zf^{(p)}(z)}{f^{(p-1)}(z)} = \frac{1+w(z)}{1-w(z)} \quad (z \in E). \quad (14)$$

Clearly,  $w(0) = 0$ . Since  $f(z)$  is  $p$ -valent in  $E$ , we have  $w(z) \neq 1$  in  $E$ . From (14) we obtain

$$\frac{f^{(p)}(z)+zf^{(p+1)}(z)}{p!} = \frac{f^{(p-1)}(z)}{p!z} \left( \left( \frac{1+w(z)}{1-w(z)} \right)^2 + \frac{2zw'(z)}{(1-w(z))^2} \right). \quad (15)$$

We claim that  $|w(z)| < 1$  in  $E$ . If this is not true, then there exists a point  $z_0 \in E$  such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1.$$

Then from the result given [18], we have

$$zw'(z_0) = kw(z_0), \text{ where } k \geq 1, w(z_0) = e^{i\theta}, 0 < \theta < 2\pi.$$

Putting  $z = z_0$  in (15), we obtain

$$\begin{aligned} \operatorname{Re} \left( \frac{f^{(p)}(z_0)+z_0f^{(p+1)}(z_0)}{p!} \right) &= \operatorname{Re} \left( \frac{f^{(p-1)}(z_0)}{p!z_0} \left( \left( \frac{1+e^{i\theta}}{1-e^{i\theta}} \right)^2 + \frac{2ke^{i\theta}}{(1-e^{i\theta})^2} \right) \right) \\ &\leq -\frac{\cos\theta+1+k}{\cos\theta-1} \operatorname{Re} \left( \frac{f^{(p-1)}(z_0)}{z_0} \right) \\ &\leq -\frac{k}{2} \operatorname{Re} \left( \frac{f^{(p-1)}(z_0)}{z_0} \right) \\ &\leq -\frac{1}{4}. \end{aligned} \quad (16)$$

Since  $k \geq 1$  and from (13),  $\operatorname{Re} \left( \frac{f^{(p-1)}(z)}{p!z} \right) > \frac{1}{2}$ ,  $z \in E$ . Inequality (16) contradicts inequality (10); thus,  $|w(z)| < 1$  in  $E$ . Equation (14) then implies that  $f(z) \in S_p^*(p-1)$  [19].

Using the Alexander type relation, we obtain the following corollary.

**Corollary 2.** If  $g(z) \in A(p)$  and

$$\operatorname{Re} \left( \frac{g^{(p)}(z)+3zg^{(p+1)}(z)+z^2g^{(p+2)}(z)}{p!} \right) > -\frac{1}{4} \quad (z \in E),$$

then  $g(z) \in C_p(p-1)$ .

**Corollary 3** [9]. If  $f(z) \in A$  and let

$$\operatorname{Re}(f'(z) + zf''(z)) > -\frac{1}{4} \quad (z \in E),$$

then  $f(z) \in S^*$ .

Our next result shows that the class  $R_p$  is closed with respect to Hadamard product.

**Theorem 5.** If  $f(z)$  and  $g(z)$  belong to the class  $R_p$  and

$$h^{(p-1)}(z) = f^{(p-1)}(z) * g^{(p-1)}(z),$$

then  $h(z)$  also belongs to the class  $R_p$ .

**Proof.** Since

$$h^{(p-1)}(z) = f^{(p-1)}(z) * g^{(p-1)}(z),$$

it follows that

$$zh^{(p)}(z) = zf^{(p)}(z) * g^{(p-1)}(z).$$

A simple computation gives

$$\operatorname{Re} \left( \frac{h^{(p)}(z)+zh^{(p+1)}(z)}{p!} \right) = \operatorname{Re} \left( \left( \frac{f^{(p)}(z)+zf^{(p+1)}(z)}{p!} \right) * \left( \frac{g^{(p-1)}(z)}{p!z} \right) \right). \quad (17)$$

From (17), using (1), (13) and Lemma 3, we have the desired result.

**Corollary 4** [9]. If  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$  and  $g(z) = z + \sum_{n=2}^{\infty} b_n z^n$  belong to  $R$ , then their Hadamard product

$$h(z) = (f * g)(z)$$

also belongs to  $R$ .

From the proof of Theorem 5, it is clear that the following more general result holds.

**Theorem 6.** If  $f(z) \in R_p$ ,  $g(z) \in A(p)$  and  $Re\left(\frac{g^{(p-1)}(z)}{p!z}\right) > \frac{1}{2}$ ,  $z \in E$ , then

$$f^{(p-1)}(z) * g^{(p-1)}(z)$$

also belongs to the class  $R_p$ .

**Theorem 7.** Let  $f(z) \in R_p$ . Then  $Re(f^{(p)}(z)) > 0$ ,  $z \in E$  and hence  $f(z)$  is  $p$ -valent in  $E$ .

**Proof.** Using the result given [17] and applying Lemma 2 with  $N(z) = \frac{zf^{(p)}(z)}{p!}$  and  $D(z) = z$  proves Theorem 7. If  $p = 1$ , then Theorem 7 was proved earlier [20].

Ruscheweyh and Small [21] have proved that the class  $C$  is closed with respect to Hadamard product. In what follows we prove that the Hadamard product of functions of the class  $R_p$  belongs to the class  $C_p(\alpha)$ .

**Theorem 8.** If  $f(z)$  and  $g(z)$  belong to  $R_p$  and

$$h^{(p-1)}(z) = f^{(p-1)}(z) * g^{(p-1)}(z),$$

then  $h(z) \in C_p(p-1)$ .

**Proof.** In view of Corollary 3, it is sufficient to show that

$$Re\left(\frac{h^{(p)}(z) + 3zh^{(p+1)}(z) + z^2h^{(p+2)}(z)}{p!}\right) > -\frac{1}{4} \quad (z \in E).$$

Equivalently, this can be written as

$$Re\left(1 + \sum_{n=1}^{\infty} (n+1) \left(\frac{(n+p)!}{n!p!}\right)^2 a_{n+p} b_{n+p} z^n\right) > -\frac{1}{4} \quad (z \in E). \quad (18)$$

Since  $f(z), g(z) \in R_p$ , we have

$$Re\left(1 + \frac{1}{2} \sum_{n=1}^{\infty} \frac{(n+p)!(n+1)}{p!n!} a_{n+p} z^n\right) > \frac{1}{2} \quad (z \in E),$$

and

$$Re\left(1 + \frac{1}{2} \sum_{n=1}^{\infty} \frac{(n+p)!(n+1)}{p!n!} b_{n+p} z^n\right) > \frac{1}{2} \quad (z \in E).$$

Using Lemma 3, we obtain

$$Re\left(1 + \frac{1}{4} \sum_{n=1}^{\infty} (n+1)^2 \left(\frac{(n+p)!}{n!p!}\right)^2 a_{n+p} b_{n+p} z^n\right) > \frac{1}{2} \quad (z \in E). \quad (19)$$

Consider the function

$$h(z) = \left(1 + 4 \sum_{n=1}^{\infty} \frac{1}{n+1} z^n\right). \quad (20)$$

Clearly,  $h(z)$  is analytic in  $E$ ,  $h(0) = 1$  and

$$\begin{aligned} Re\left(1 + 4 \sum_{n=1}^{\infty} \frac{1}{n+1} z^n\right) &= Re\left(-3 - \frac{4}{z} \log(1-z)\right) \\ &> -3 + 4 \log 2 \quad [15] \\ &> -\frac{1}{4} \quad (z \in E). \end{aligned} \quad (21)$$

From (19) and (20), we can write

$$\begin{aligned} & \left(1 + \sum_{n=1}^{\infty} (n+1) \left(\frac{(n+p)!}{n!p!}\right)^2 a_{n+p} b_{n+p} z^n\right) \\ &= \left(1 + \frac{1}{4} \sum_{n=1}^{\infty} (n+1)^2 \left(\frac{(n+p)!}{n!p!}\right)^2 a_{n+p} b_{n+p} z^n\right) * \left(1 + 4 \sum_{n=1}^{\infty} \frac{1}{n+1} z^n\right). \end{aligned} \quad (22)$$

Using (19), (21), (22) and Lemma 3, we see that (18) holds for all  $z \in E$ . This completes the proof.

**Corollary 5** [9]. If  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$  and  $g(z) = z + \sum_{n=2}^{\infty} b_n z^n$  belong to  $R$ , then

$$h(z) = (f * g)(z) \in C.$$

We now define the integral operator  $F(z)$  as follows. Let  $f(z) \in A(p)$ ; then

$$\begin{aligned} \frac{F^{(p-1)}(z)}{p!} &= \frac{c+1}{z^c} \int_0^z t^{c-1} \left(\frac{f^{(p-1)}(t)}{p!}\right) dt, \quad (c > -p, p \in N = \{1, 2, \dots\}). \\ &= z + \sum_{n=1}^{\infty} \frac{(c+1)(n+p)!}{(n+c+1)(n+1)!p!} a_{n+p} z^{n+1}. \end{aligned} \quad (23)$$

For  $p=1$ , the operator defined in (23) is a generalised form of operator by Bernardi [22]. A comprehensive study of operators with applications can be found in the literature [23-30].

**Theorem 9.** Let  $f(z) \in R_p$  and

$$\frac{F^{(p-1)}(z)}{p!} = \frac{c+1}{z^c} \int_0^z t^{c-1} \left(\frac{f^{(p-1)}(t)}{p!}\right) dt \quad (z \in E). \quad (24)$$

Then  $F(z) \in R_p$ .

**Proof.** Let

$$\frac{F^{(p)}(z) + zF^{(p+1)}(z)}{p!} = h(z).$$

Then  $h(z)$  is analytic in  $E$  and  $h(0) = 1$ .

From (24), we have

$$\left(\frac{z^c F^{(p-1)}(z)}{p!}\right)' = (c+1)z^{c-1} \frac{f^{(p-1)}(z)}{p!}.$$

A simple computation gives us:

$$\operatorname{Re} \left( h(z) + \frac{zh'(z)}{c+1} \right) = \operatorname{Re} \left( \frac{f^{(p)}(z) + zF^{(p+1)}(z)}{p!} \right).$$

From the hypothesis of Theorem 9 with the result given [31], we have

$$\operatorname{Re} \left( \frac{F^{(p)}(z) + zF^{(p+1)}(z)}{p!} \right) > 0 \quad (z \in E).$$

This completes the proof. It is easy to see that if  $0 \leq \lambda \leq 1$ , and  $f(z)$  and  $g(z)$  are in  $R_p$ , then  $G(z) = \lambda g(z) + (1-\lambda)f(z)$  is also in  $R_p$ . This shows that the class  $R_p$  is a convex set.

**Theorem 10.** Let  $f(z), g(z) \in A(p)$  and  $\alpha, \beta < 1$ . If

$$\frac{f^{(p)}(z)}{p!} \in P(\alpha), \quad \frac{g^{(p)}(z)}{p!} \in P(\beta),$$

and

$$\phi^{(p-1)}(z) = f^{(p-1)}(z) * g^{(p-1)}(z),$$

then  $\phi(z) \in \mathcal{S}_p^*(p-1)$ , provided that

$$(1-\alpha)(1-\beta) < \frac{3}{8(\ln 2 - 1)^2 + 4}. \quad (25)$$

**Proof.** Using the given hypothesis on  $f(z)$  and  $g(z)$  and Lemma 5, we have

$$\begin{aligned} \operatorname{Re} \left( \frac{f^{(p)}(z)}{p!} * \frac{g^{(p)}(z)}{p!} \right) &= \operatorname{Re} \left( \frac{\phi^{(p)}(z) + z\phi^{(p+1)}(z)}{p!} \right) \\ &> 1 - 2(1 - \alpha)(1 - \beta). \end{aligned} \quad (26)$$

By using Lemma 4, from (26) we have

$$\operatorname{Re} \left( \frac{\phi^{(p)}(z)}{p!} \right) > 1 + 4(1 - \alpha)(1 - \beta)(\ln 2 - 1) \quad (z \in E),$$

or equivalently, this can be written as

$$\operatorname{Re} \left( \frac{\phi^{(p-1)}(z)}{p!z} + z \left( \frac{\phi^{(p-1)}(z)}{p!z} \right)' \right) > 1 + 4(1 - \alpha)(1 - \beta)(\ln 2 - 1) \quad (z \in E). \quad (27)$$

Using Lemma 4, (27) becomes

$$\operatorname{Re} \left( \frac{\phi^{(p-1)}(z)}{p!z} \right) > 1 - 8(1 - \alpha)(1 - \beta)(\ln 2 - 1)^2.$$

Suppose

$$h(z) = \frac{z\phi^{(p)}(z)}{\phi^{(p-1)}(z)} \quad \text{and} \quad q(z) = \frac{\phi^{(p-1)}(z)}{z}.$$

Then  $h(z)$  is analytic in  $E$ ,  $h(0) = 1$  and

$$\operatorname{Re} q(z) > 1 - 8(1 - \alpha)(1 - \beta)(\ln 2 - 1)^2. \quad (28)$$

A simple computation gives us:

$$\begin{aligned} \phi^{(p)}(z) + z\phi^{(p+1)}(z) &= q(z)(h^2(z) + zh'(z)) \\ &= \Psi(h(z), zh'(z), z). \end{aligned} \quad (29)$$

By taking  $u = h(z)$  and  $v = zh'(z)$ ,  $\Psi(u, v; z) = q(z)(u^2 + v)$ . From (25) and (29) we have

$$\operatorname{Re} \left( \Psi(h(z), zh'(z), z) \right) > 1 - 2(1 - \alpha)(1 - \beta) \quad (z \in E).$$

Now for real  $x, y \leq -\frac{1}{2}(1 + x^2)$ , we have

$$\begin{aligned} \operatorname{Re}(\Psi(ix, y, z)) &= (-x^2 + y)\operatorname{Re} q(z) \\ &\leq -\frac{1}{2}(1 + 3x^2)\operatorname{Re} q(z) \\ &\leq -\frac{1}{2}\operatorname{Re} q(z) \quad (z \in E). \end{aligned} \quad (30)$$

From (28) and (30) we obtain

$$\operatorname{Re}(\Psi(ix, y, z)) \leq 1 - 2(1 - \alpha)(1 - \beta), \quad \text{for all } z \in E.$$

By using the results given [19, 31], we have  $\phi(z) \in S_p^*(p-1)$  ( $z \in E$ ).

**Corollary 6.** Let  $f(z), g(z) \in A(p)$  and  $\alpha, \beta < 1$ . If

$$\frac{f^{(p)}(z)}{p!} \in P(\alpha), \quad \frac{g^{(p)}(z)}{p!} \in P(\beta),$$

and

$$\psi^{(p-1)}(z) = \int_0^z \frac{(f^{(p-1)} + g^{(p-1)})(t)}{t} dt,$$

then  $\psi(z) \in C_p(p-1)$ , provided that

$$(1 - \alpha)(1 - \beta) < \frac{3}{8(\ln 2 - 1)^2 + 4}.$$

The proof is simple by taking  $z\psi^{(p)}(z) = \phi^{(p-1)}(z)$ .

**Theorem 11.** Let  $f(z), g(z), h(z) \in A(p)$ ,  $\alpha, \beta, \gamma < 1$ . If

$$\frac{f^{(p)}(z)}{p!} \in P(\alpha), \frac{g^{(p)}(z)}{p!} \in P(\beta), \frac{h^{(p)}(z)}{p!} \in P(\gamma),$$

and

$$\varnothing^{(p-1)}(z) = (f^{(p-1)} * g^{(p-1)} * h^{(p-1)})(z),$$

then  $\varnothing(z) \in S_p^*(p-1)$ , provided that

$$(1-\alpha)(1-\beta)(1-\gamma) < \frac{3}{\{8(\ln 2-1)^2+4\}(-4)(\ln 2-1)}.$$

**Proof.** By the hypotheses on  $f, g$  and  $h$  and Lemma 5, we obtain

$$\begin{aligned} \operatorname{Re} \left( \frac{K^{(p)}(z)}{p!} * \frac{h^{(p)}(z)}{p!} \right) &= \operatorname{Re} \left( \frac{\varnothing^{(p)}(z) + z\varnothing^{(p+1)}(z)}{p!} \right) \\ &> 1 - 2(1-\alpha_1)(1-\beta), \end{aligned} \quad (31)$$

where  $K^{(p-1)}(z) = (f^{(p-1)} * g^{(p-1)})(z)$  and  $\operatorname{Re} \frac{K^{(p)}(z)}{p!} > \alpha_1, \alpha_1 = 1 + 4(1-\alpha)(1-\beta)(\ln 2 - 1)$ .

From (31), together with Lemma 4, we have

$$\operatorname{Re} \left( \frac{\varnothing^{(p)}(z)}{p!} \right) > 1 - 16(1-\alpha)(1-\beta)(1-\gamma)(\ln 2 - 1)^2 \quad (z \in E),$$

Using the same technique similar to that of Theorem 10, we obtain the required result.

**Corollary 7.** Let  $f(z), g(z), h(z) \in A(p)$ ,  $\alpha, \beta, \gamma < 1$ . If

$$\frac{f^{(p)}(z)}{p!} \in P(\alpha), \frac{g^{(p)}(z)}{p!} \in P(\beta), \frac{h^{(p)}(z)}{p!} \in P(\gamma),$$

and

$$\varphi^{(p-1)}(z) = (f^{(p-1)} * g^{(p-1)} * h^{(p-1)})(z),$$

Then  $\varphi(z) \in C_p(p-1)$ , provided that

$$(1-\alpha)(1-\beta)(1-\gamma) < \frac{3}{16(\ln 2-1)^2+8}.$$

For proving  $\varphi(z) \in C_p(p-1)$ , it is sufficient to show that

$$\zeta^{(p-1)}(z) = z\varphi^{(p)}(z) \in S_p^*(p-1).$$

By the hypotheses on  $f(z), g(z)$  and  $h(z)$  and Lemma 5, we obtain

$$\begin{aligned} \operatorname{Re} \left( \frac{(f^{(p)} * g^{(p)} * h^{(p)})(z)}{p!} \right) &= \operatorname{Re} \left( \frac{\zeta^{(p)}(z) + z\zeta^{(p+1)}(z)}{p!} \right) \\ &> 1 - 4(1-\alpha)(1-\beta)(1-\gamma), \end{aligned}$$

and the proof is completed similarly to that of Theorem 10. If  $p = 1$ , then Theorem 10 and Corollary 7 were given [10].

**Theorem 12.** Let  $f_1(z), f_2(z), \dots, f_n(z) \in A(p)$ ,  $\alpha_1, \alpha_1, \dots, \alpha_n < 1$ . If

$$\frac{f_1^{(p)}(z)}{p!} \in P(\alpha_1), \frac{f_2^{(p)}(z)}{p!} \in P(\alpha_2), \dots, \frac{f_n^{(p)}(z)}{p!} \in P(\alpha_n),$$

and

$$\tau^{(p-1)}(z) = (f_1^{(p-1)} * f_2^{(p-1)} * \dots * f_n^{(p-1)})(z), \quad (32)$$

then  $\tau(z) \in S_p^*(p-1)$ , provided that

$$(1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_n) < \frac{3}{\{8(\ln 2 - 1)^2 + 4\}((-4)(\ln 2 - 1))^{n-2}}, \quad n \geq 2. \quad (33)$$

**Proof.** For proving the above Theorem, we use the principle of mathematical induction. For  $n = 2$ , we have proved Theorem 10; thus, (32) holds for  $n = 2$ . Suppose that (32) holds true for  $n = k$ ; that is,

$$\tau^{(p-1)}(z) = (f_1^{(p-1)} * f_2^{(p-1)} * \dots * f_k^{(p-1)})(z),$$

then  $\tau(z) \in S_p^*(p-1)$ , provided that inequality (33) is satisfied.

We have to prove that (32) holds true for  $n = k + 1$ . For this, consider

$$\tau^{(p-1)}(z) = (f_1^{(p-1)} * f_2^{(p-1)} * \dots * f_{k+1}^{(p-1)})(z).$$

Now using the given hypothesis on  $f_j(z)$ ,  $j = 1, 2, \dots, k$  and Lemma 5, we have

$$\begin{aligned} \operatorname{Re} \left( \frac{M^{(p)}(z)}{p!} * \frac{f_{k+1}^{(p)}(z)}{p!} \right) &= \operatorname{Re} \left( \frac{\tau^{(p)}(z) + z\tau^{(p+1)}(z)}{p!} \right) \\ &> 1 - 2(1 - \alpha^*)(1 - \alpha_{k+1}), \end{aligned} \quad (34)$$

where  $M^{(p-1)}(z) = (f_1^{(p-1)} * f_2^{(p-1)} * \dots * f_k^{(p-1)})$  and

$$\operatorname{Re} \frac{M^{(p)}(z)}{p!} > \alpha^*, \quad \alpha^* = 1 + 4(1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_k)(\ln 2 - 1)^{k-1}(-4)^{k-2}.$$

By using Lemma 4, from (34) we have

$$\operatorname{Re} \frac{\tau^{(p)}(z)}{p!} > 1 + 4(1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_k)(\ln 2 - 1)^k(-4)^{k-1}, \quad z \in E, \quad k \geq 2. \quad (35)$$

Applying Lemma 4, (35) can be written as

$$\operatorname{Re} \frac{\tau^{(p-1)}(z)}{zp!} > 1 - 8(1 - \alpha^*)(1 - \alpha_{k+1}).$$

Now with the same procedure used in Theorem 10, we have  $\tau(z) \in S_p^*(p-1)$ , provided that

$$(1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_k)(1 - \alpha_{k+1}) < \frac{3}{\{8(\ln 2 - 1)^2 + 4\}((-4)(\ln 2 - 1))^{k-1}}.$$

Therefore, the result is true for  $n = k + 1$  and hence by using mathematical induction, (32) holds true for all  $n \geq 2$ . This completes the proof.

#### ACKNOWLEDGEMENTS

The authors would like to thank Dr. S. M. Junaid Zaidi, rector of Comsats Institute of Information Technology, for providing excellent research facilities. This research was carried out under the Higher Education Commission project grant no. NRPU No. 20-1966/R&D/11-2553. The authors are also grateful to the referees for their helpful comments and suggestions which improved the presentation of the paper.

#### REFERENCES

1. S. Owa, "On certain classes of  $p$ -valent functions with negative coefficients", *Simon Stevin.*, **1985**, 59, 385-402.
2. J. L. Liu, "Certain convolution properties of multivalent analytic functions associated with a linear operator", *General Math.*, **2009**, 17, 41-52.
3. J. Sokol, J. Dziok and M. K. Aouf, "On certain classes of meromorphically  $p$ -valent starlike and  $p$ -valent quasi-convex functions", *Tamkang J. Math.*, **2013**, 44, 261-269.

4. M. Arif, "Sufficiency criteria for a class of p-valent analytic functions of complex order", *Abst. Appl. Anal.*, **2013**, doi: 10.1155/2013/517296.
5. M. Raza, M. Arif and M. Darus, "Fekete-Szego inequality for a subclass of p-valent analytic functions", *J. Appl. Math.*, **2013**, 2013, Art.ID 127615.
6. S. Hussain, N. Anjum and I. Z. Cheema, "Certain classes of multivalent functions related with a linear operator", *Acta. Uni. Apul.*, **2011**, 28, 367-377.
7. Z. Shareef, S. Hussain and M. Darus, "Convolution operators in the geometric functions theory", *J. Ineq. Appl.*, **2012**, doi: 10.1186/1029-242X-2012-213.
8. J. Sokol, "Classes of multivalent functions associated with a convolution operator", *Comput. Math. Appl.*, **2010**, 60, 1343-1350.
9. R. Singh and S. Singh, "Convolution properties of a class of starlike functions", *Proc. Amer. Math. Soc.*, **1989**, 106, 145-152.
10. A. Y. Lashin, "Some convolution properties of analytic functions", *Appl. Math. Lett.*, **2005**, 18, 135-138.
11. L. Fejer, "Über die positivitat von summen, die nach trigonometrischen order Legendreschen funktionen fortschreiten", *Acta Litt. Ac. Sci. Szeged*, **1925**, 2, 75-86.
12. R. J. Libera, "Some classes of regular univalent functions", *Proc. Amer. Math. Soc.*, **1965**, 16, 755-758.
13. L. Yi and S. Ding, "A new criterion for starlike functions", *Int. J. Math. Math. Sci.*, **1996**, 19, 613-614.
14. S. Ponnusamy and V. Singh, "Convolution properties of some classes of analytic functions", *J. Math. Sci.*, **1998**, 89, 1008-1020.
15. M. S. Robertson, "An extremal problem for functions with positive real part", *Michigan Math. J.*, **1964**, 11, 327-335.
16. W. Rogosinski and G. Szego, "Über die abschimlte von potenzreihen die in ernein kreise be schrankt bleiben", *Math. Z.*, **1928**, 28, 73-94.
17. S. Ozaki, "On the theory of multivalent functions", *Sci. Rep. Tokyo Bunrika Daigaku A.*, **1935**, 40, 167-188.
18. I. S. Jack, "Functions starlike and convex of order  $\alpha$ ", *J. London Math. Soc.*, **1971**, s2-3, 469-474.
19. M. Nunokawa, S. Owa, T. Sekine, R. Yamakawa, H. Saitoh and J. Nishiwaki, "On certain multivalent functions", *Int. J. Math. Math. Sci.*, **2007**, 2007, Art. ID 72393.
20. P. N. Chichra, "New subclasses of the class of close-to-convex functions", *Proc. Amer. Math. Soc.*, **1977**, 62, 37-43.
21. S. Ruscheweyh and T. Sheil-Small, "Hadamard products of schlicht functions and the Polya-Schoenberg conjecture", *Comm. Math. Helv.*, **1973**, 48, 119-135.
22. S. D. Bernardi, "Convex and starlike univalent functions", *Trans. Amer. Math. Soc.*, **1969**, 135, 429-446.
23. B. C. Carlson and D. B. Shaffer, "Starlike and prestarlike hypergeometric functions", *SIAM J. Math. Anal.*, **1984**, 15, 737-745.

24. N. E. Cho, O. S. Kwon and H. M. Srivastava, "Inclusion relationships and argument properties for certain subclasses of multivalent functions associated with a family of linear operators", *J. Math. Anal. Appl.*, **2004**, 292, 470-483.
25. J. Dziok and H. M. Srivastava, "Classes of analytic functions associated with the generalized hypergeometric function", *Appl. Math. Comput.*, **1999**, 103, 1-13.
26. J. L. Liu and H. M. Srivastava, "A linear operator and associated families of meromorphically multivalent functions", *J. Math. Anal. Appl.*, **2001**, 259, 566-581.
27. K. I. Noor, "Some classes of p-valent analytic functions defined by certain integral operator", *Appl. Math. Comput.*, **2004**, 157, 835-840.
28. K. I. Noor, "On new classes of integral operators", *J. Natur. Geom.*, **1999**, 16, 71-80.
29. K. I. Noor and M. A. Noor, "On integral operators", *J. Math. Anal. Appl.*, **1999**, 238, 341-352.
30. J. Sokol, K. I. Noor and H. M. Srivastava, "A family of convolution operators for multivalent analytic functions", *Eur. J. Pure Appl. Math.*, **2012**, 5, 469-479.
31. D. J. Hallenbeck and S. Ruscheweyh, "Subordination by convex functions", *Proc. Amer. Math. Soc.*, **1975**, 52, 191-195.