

Effect of Spinosad on the heat tolerance and cold tolerance of *Sitophilus oryzae* (Coleoptera: Curculionidae)

Wijayaratne, L.K.W.*#

Department of Plant Sciences, Faculty of Agriculture, Rajarata University of Sri Lanka, Puliyankulama, Anuradhapura, Sri Lanka

*Corresponding author, Email: wollylk@yahoo.com

#Presenting author, Email: wollylk@yahoo.com

DOI: xx.xxxx/xxx.2014.xxx.xxx.xxx

Abstract

Rice Weevil *Sitophilus oryzae* (Coleoptera: Curculionidae) is a serious insect pest of stored products in Sri Lanka. Currently Pirimiphos Methyl is recommended to control this species but grain handlers seek for alternatives. Exposure to high or low temperature is popular in stored-product insect pest management but are expensive. Therefore it is important to look for treatments that synergize extreme temperature sensitivity. Spinosad is effective against certain stored-product insects but has not yet been tested for its synergy with heat or cold. This experiment was conducted to evaluate effect of Spinosad on the heat tolerance and cold tolerance of *Sitophilus oryzae* adults. Experiment was laid out in a complete randomized design with four replicates. The concentration of Spinosad and the duration of exposure were changed making them a factorial set of treatments. Four week old *Sitophilus oryzae* adults were first exposed to a series of concentration of Spinosad. Later they were held at 40°C or at 11°C for different durations. Mortality was counted 24 hours after the termination of heat or cold exposure. Increase in the exposure period to heat or cold increased mortality of *Sitophilus oryzae* adults. Pre exposure to Spinosad at 12.5 ppm or above synergized the adult mortality at high or low temperature showing a dose response. This study shows that heat tolerance and cold tolerance of *Sitophilus oryzae* adults is reduced by pre exposure to Spinosad. Therefore, Spinosad can be used as a grain protectant in combination with high or low temperatures against *Sitophilus oryzae* in particular and need to test for its efficacy on other stored-product insect species as well.

Keywords: *Sitophilus oryzae*, Spinosad, cold tolerance, heat tolerance, synergy

1. Introduction

Rice Weevil *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae) is a serious insect pest of stored products in Sri Lanka (Gunarathna and Karunaratne, 2009). Currently Pirimiphos Methyl is recommended to control this species (Rubasinghe et al., 2006) but grain handlers seek for alternatives. Exposure to high (Beckett et al., 2007) or low temperature (Fields, 1992) is popular in stored-product insect pest management but are expensive. Therefore it is important to look for treatments that synergize extreme temperature sensitivity of these insects. Spinosad is a fermented bacterial formulation derived from *Saccharopolyspora spinosa* Mertz and Yao (Mertz and Yao, 1990). Spinosad is effective against certain stored-product insects (Huang and Subramanyam, 2007) but has not yet been tested for its synergy with heat or cold. The objectives of this study were to evaluate effect of Spinosad on the heat tolerance and cold tolerance of *S. oryzae* adults.

2. Materials and methods

A commercial preparation of Spinosad available at the local market in Sri Lanka was used as the test compound. A concentration series was prepared by diluting the test compound in distilled water. Red rice (variety Basmati) was used in the experiment to spray Spinosad. One of the selected Spinosad concentrations (0.5 mL) was sprayed onto rice placed in each Petri dish, covered with the lid, and was shaken well to ensure uniform distribution of sprayed concentration. Spraying was conducted from the lowest to the highest concentration of Spinosad. The distilled water was sprayed as the control. There were six exposure periods for heat tolerance experiment and seven exposure periods for the cold tolerance experiment. For each treatment, with a particular concentration and exposure period, there were four replicates. Twenty *S. oryzae* adults were introduced to each Petri dish containing the sprayed rice medium, and were held at room temperature ($32\pm 1^\circ\text{C}$) for 36-48 h. Later, they were kept in an oven maintained at 40°C or in a refrigerator at $11\pm 0.5^\circ\text{C}$ for different durations, and mortality counted. Temperature was recorded by a portable digital thermometer (TECPEL 506B CE, Taiwan).

3. Results and Discussion

Increased duration at high or low temperature increased the mortality of *S. oryzae* adults. Pre exposure to Spinosad synergized the adult mortality at high or low temperature.




Figure 1 Heat tolerance of *S. oryzae* adults.

Cold tolerance




Figure 2 Effect of cold tolerance on *Sitophilus oryzae*.

This study shows that heat tolerance and cold tolerance of *S. oryzae* adults are reduced by prior exposure to Spinosad. Therefore, Spinosad can be used as a grain protectant in particular against *S. oryzae* in combination with high or low temperatures and need to be tested for its efficacy on other stored-product insect species as well.

Acknowledgements

This research was carried out by using the funds allocated as the Research Allowance to the author. The findings were presented at the IWCSPP held in Chiang Mai, Thailand using the funds provided by the University Grants Commission of Sri Lanka, under the Encouragement Awards Scheme (International Mobility Support). The author also wishes to thank Dr. A. Balasuriya, Head, Department of Plant Sciences, Rajarata University of Sri Lanka for the administrative support; Mr. M.H.J.P. Gunarathne, Head, Department of Agric. Engineering and Soil Science for providing the oven for the study; Hayleys Agriculture Holdings Limited, Sri Lanka for providing the commercial product of Spinosad; and Mr. Sarath Dayawansa for technical support.

References

- Beckett, S.J., Fields, P.G., Subramanyam, B., 2007. Disinfestation of stored products and associated structures using heat. In: Tang, J., Mitcham, E., Wang, S., Lurie, S. (Eds.), Heat Treatments for Postharvest Pest Control: Theory and Practice. CABI, Wallingford, pp. 182-237.
- Fields, P.G., 1992. The control of stored-product insects and mites with extreme temperatures. *Journal of Stored Products Research* 28, 89-118.
- Gunarathna, T.V.N.M., Karunaratne, M.M.S.C., 2009. Laboratory evaluation of some Sri Lankan plants as post-harvest grain protectants for the control of rice weevil *Sitophilus oryzae*. *Vidyodaya* 14, 69-83.
- Huang, F., Subramanyam, B., 2007. Effectiveness of spinosad against seven major stored-grain insects on corn. *Insect Science* 14, 225-230.
- Mertz, E.P., Yao, R.C., 1990. *Saccharopholyspora spinosa* sp. nov. isolated from soil collected in a sugar rum still. *International Journal of Systemic Bacteriology* 40, 34-39.
- Rubasinghe, G.R.S., Paranagama, P.A., Abeywickrama, K.P., 2006. Physiochemical changes of stored cowpea, *Vigna unguiculata* treated with selected essential oils to control cowpea bruchid, *Callosobruchus maculatus* (F.). *Journal of Food Agriculture and Environment* 4, 41-44.