



REFERENCES

- Aebersold R, Mann M. Mass spectrometry-based proteomics. **Nature** 2003; 422(6928): 198-207.
- Alur M, Nguyen MM, Eggener SE, Jiang F, Dadras SS, Stern J, et al. Suppressive roles of calreticulin in prostate cancer growth and metastasis. **Am J Pathol** 2009; 175(2): 882-90.
- Andersen JS, Wilkinson CJ, Mayor T, Mortensen P, Nigg EA, Mann M. Proteomic characterization of the human centrosome by protein correlation profiling. **Nature** 2003; 426(6966): 570-4.
- Anderson L, Seilhamer J. A comparison of selected mRNA and protein abundances in human liver. **Electrophoresis** 1997; 18(3): 533-7.
- Attwood HD, Chou ST. The longevity of *Clonorchis sinensis*. **Pathology** 1978; 10(2): 153-6.
- Bai Z, Ye Y, Liang B, Xu F, Zhang H, Zhang Y, et al. Proteomics-based identification of a group of apoptosis-related proteins and biomarkers in gastric cancer. **Int J Oncol** 2011; 38(2): 375-83.
- Banks RE, Dunn MJ, Forbes MA, Stanley A, Pappin D, Naven T, et al. The potential use of laser capture microdissection to selectively obtain distinct populations of cells for proteomic analysis--preliminary findings. **Electrophoresis** 1999; 20(4-5): 689-700.
- Bao H, Jiang M, Zhu M, Sheng F, Ruan J, Ruan C. Overexpression of annexin II affects the proliferation, apoptosis, invasion and production of proangiogenic factors in multiple myeloma. **Int J Hematol** 2009; 90(2): 177-85.
- Bisca A, D'Ambrosio C, Scaloni A, Puglisi F, Aprile G, Piga A, et al. Proteomic evaluation of core biopsy specimens from breast lesions. **Cancer Lett** 2004; 204(1): 79-86.
- Bishop JG, Corces VG. The nucleotide sequence of a *Drosophila melanogaster* enolase gene. **Nucleic Acids Res** 1990; 18(1): 191.

- Blechacz B, Gores GJ. Cholangiocarcinoma: advances in pathogenesis, diagnosis, and treatment. **Hepatology** 2008; 48(1): 308-21.
- _____, Gores GJ. Cholangiocarcinoma. **Clin Liver Dis** 2008; 12(1): 131-50.
- Bonner RF, Emmert-Buck M, Cole K, Pohida T, Chuaqui R, Goldstein S, et al. Laser capture microdissection: molecular analysis of tissue. **Science** 1997; 278(5342): 1481-83.
- Bouvard V, Baan R, Straif K, Grosse Y, Secretan B, El Ghissassi F, et al. A review of human carcinogens--Part B: biological agents. **Lancet Oncol** 2009; 10(4): 321-2.
- Bower M, Rustin GJ, Newlands ES, Holden L, Short D, Foskett M, et al. Chemotherapy for gestational trophoblastic tumours hastens menopause by 3 years. **Eur J Cancer** 1998; 34(8): 1204-7.
- Boylan KL, Andersen JD, Anderson LB, Higgins L, Skubitz AP. Quantitative proteomic analysis by iTRAQ(R) for the identification of candidate biomarkers in ovarian cancer serum. **Proteome Sci** 2010; 8(3): 1-9.
- Braden AR, Kafka MT, Cunningham L, Jones H, Vishwanatha JK. Polymeric nanoparticles for sustained down-regulation of annexin A2 inhibit prostate tumor growth. **J Nanosci Nanotechnol** 2009; 9(5): 2856-65.
- Brownstein C, Falcone DJ, Jacovina A, Hajjar KA. A mediator of cell surface-specific plasmin generation. **Ann N Y Acad Sci** 2001; 11(14): 1136-44.
- Burgess JA, Lescuyer P, Hainard A, Burkhard PR, Turck N, Michel P, et al. Identification of brain cell death associated proteins in human post-mortem cerebrospinal fluid. **J Proteome Res** 2006; 5(7): 1674-81.
- Cardone RA, Bellizzi A, Busco G, Weinman EJ, Dell'Aquila ME, Casavola V, et al. The NHERF1 PDZ2 domain regulates PKA-RhoA-p38-mediated NHE1 activation and invasion in breast tumor cells. **Mol Biol Cell** 2007; 18(5): 1768-80.
- Cargile BJ, Talley DL, Stephenson JL, Jr. Immobilized pH gradients as a first dimension in shotgun proteomics and analysis of the accuracy of pI predictability of peptides. **Electrophoresis** 2004; 25(6): 936-45.
- Cha MK, Suh KH, Kim IH. Overexpression of peroxiredoxin I and thioredoxin1 in human breast carcinoma. **J Exp Clin Cancer Res** 2009; 28(93): 1-12.

- Chang GC, Liu KJ, Hsieh CL, Hu TS, Charoenfuprasert S, Liu HK, et al. Identification of alpha-enolase as an autoantigen in lung cancer: its overexpression is associated with clinical outcomes. **Clin Cancer Res** 2006; 12(19): 5746-54.
- Chang JW, Jeon HB, Lee JH, Yoo JS, Chun JS, Kim JH, et al. Augmented expression of peroxiredoxin I in lung cancer. **Biochem Biophys Res Commun** 2001; 289(2): 507-12.
- Chang MC, Xiao S, Nose V. Clinicopathologic and immunohistochemical correlation in sporadic pancreatic endocrine tumors: possible roles of utrophin and cyclin D1 in malignant progression. **Hum Pathol** 2007; 38(5): 732-40.
- Chang YS, Wu W, Walsh G, Hong WK, Mao L. Enolase-alpha is frequently down-regulated in non-small cell lung cancer and predicts aggressive biological behavior. **Clin Cancer Res** 2003; 1(9): 3641-4.
- Chenau J, Michelland S, Sidibe J, Seve M. Peptides OFFGEL electrophoresis: a suitable pre-analytical step for complex eukaryotic samples fractionation compatible with quantitative iTRAQ labeling. **Proteome Sci** 2008; 6(9): 1-8.
- Chhipa RR, Lee KS, Onate S, Wu Y, Ip C. Prx1 enhances androgen receptor function in prostate cancer cells by increasing receptor affinity to dihydrotestosterone. **Mol Cancer Res** 2009; 7(9): 1543-52.
- Chong PK, Gan CS, Pham TK, Wright PC. Isobaric tags for relative and absolute quantitation (iTRAQ) reproducibility: implication of multiple injections. **J Proteome Res** 2006; 5(5): 1232-40.
- David CJ, Chen M, Assanah M, Canoll P, Manley JL. HnRNP proteins controlled by c-myc deregulate pyruvate kinase mRNA splicing in cancer. **Nature** 463(7279): 364-8.
- Deora AB, Kreitzer G, Jacovina AT, Hajjar KA. An annexin 2 phosphorylation switch mediates p11-dependent translocation of annexin 2 to the cell surface. **J Biol Chem** 2004; 279(42): 43411-8.
- Desmetz C, Bibeau F, Boissiere F, Bellet V, Rouanet P, Maudelonde T, et al. Proteomics-based identification of HSP60 as a tumor-associated antigen in early stage breast cancer and ductal carcinoma *in situ*. **J Proteome Res** 2008; 7(9): 3830-7.

- Dunn BK, Jegalian K, Greenwald P. Biomarkers for early detection and as surrogate endpoints in cancer prevention trials: issues and opportunities. **Recent Results Cancer Res** 2011; 188: 21-47.
- Elias JE, Gygi SP. Target-decoy search strategy for increased confidence in large-scale protein identifications by mass spectrometry. **Nat Methods** 2007; 4(3): 207-14.
- Elliott MH, Smith DS, Parker CE, Borchers C. Current trends in quantitative proteomics. **J Mass Spectrom** 2009; 44(12): 1637-60.
- Emoto K, Sawada H, Yamada Y, Fujimoto H, Takahama Y, Ueno M, et al. Annexin II overexpression is correlated with poor prognosis in human gastric carcinoma. **Anticancer Res** 2001a; 21(2): 1339-45.
- _____, Yamada Y, Sawada H, Fujimoto H, Ueno M, Takayama T, et al. Annexin II overexpression correlates with stromal tenascin-C overexpression: a prognostic marker in colorectal carcinoma. **Cancer** 2001b; 92(6): 1419-26.
- Erikson E, Erikson RL. Identification of a cellular protein substrate phosphorylated by the avian sarcoma virus-transforming gene product. **Cell** 1980; 21(3): 829-36.
- Ernault E, Bourreau A, Gamelin E, Guette C. A proteomic approach for plasma biomarker discovery with iTRAQ labelling and OFFGEL fractionation. **J Biomed Biotechnol** 2010; (9): 27-917.
- Fedor HL, De Marzo AM. Practical methods for tissue microarray construction. **Methods Mol Med** 2005; 10(3): 89-101.
- Fontana S, Pucci-Minafra I, Becchi M, Freyria AM, Minafra S. Effect of collagen substrates on proteomic modulation of breast cancer cells. **Proteomics** 2004; 4(3): 849-60.
- Friedman DB, Hill S, Keller JW, Merchant NB, Levy SE, Coffey RJ, et al. Proteome analysis of human colon cancer by two-dimensional difference gel electrophoresis and mass spectrometry. **Proteomics** 2004; 4(3): 793-811.
- Gan CS, Chong PK, Pham TK, Wright PC. Technical, experimental, and biological variations in isobaric tags for relative and absolute quantitation (iTRAQ). **J Proteome Res** 2007; 6(2): 821-7.

- Gasteiger E, Gattiker A, Hoogland C, Ivanyi I, Appel RD, Bairoch A. ExpPASy: the proteomics server for in-depth protein knowledge and analysis. **Nucleic Acids Res** 2003; 31(13): 3784-8.
- Georgescu MM, Morales FC, Molina JR, Hayashi Y. Roles of NHERF1/EBP50 in cancer. **Curr Mol Med** 2008; 8(6): 459-68.
- _____. NHERF1: molecular brake on the PI3K pathway in breast cancer. **Breast Cancer Res** 2008; 10(2): 106-7.
- Glen A, Gan CS, Hamdy FC, Eaton CL, Cross SS, Catto JW, et al. iTRAQ-facilitated proteomic analysis of human prostate cancer cells identifies proteins associated with progression. **J Proteome Res** 2008; 7(3): 897-907.
- Gonzalez AM, Otey C, Edlund M, Jones JC. Interactions of a hemidesmosome component and actinin family members. **J Cell Sci** 2001; 114(23): 4197-206.
- Grubman SA, Perrone RD, Lee DW, Murray SL, Rogers LC, Wolkoff LI, et al. Regulation of intracellular pH by immortalized human intrahepatic biliary epithelial cell lines. **Am J Physiol** 1994; 266(6): 1060-70.
- Harinasuta T, Riganti M, Bunnag D. Opisthorchis viverrini infection: pathogenesis and clinical features. **Arzneimittelforschung** 1984; 34(9): 1167-9.
- Hay MM, Chen TR, McClintock P and Reid Y. American Type Culture Collection Catalogue of Cell Lines and Hybridomas. **6th ed, Rockville, Md: American Type Culture Collection** 1988: 156-83.
- Hood BL, Darfler MM, Guiel TG, Furusato B, Lucas DA, Ringeisen BR, et al. Proteomic analysis of formalin-fixed prostate cancer tissue. **Mol Cell Proteomics** 2005; 4(11): 1741-53.
- Hu HD, Ye F, Zhang DZ, Hu P, Ren H, Li SL. iTRAQ quantitative analysis of multidrug resistance mechanisms in human gastric cancer cells. **J Biomed Biotechnol** 2010; 2010(571343): 1-10.
- Huang HY, Li CF, Fang FM, Tsai JW, Li SH, Lee YT, et al. Prognostic implication of ezrin overexpression in myxofibrosarcomas. **Ann Surg Oncol** 2010; 17(12): 3212-9.
- Huang JS, Chao CC, Su TL, Yeh SH, Chen DS, Chen CT, et al. Diverse cellular transformation capability of overexpressed genes in human hepatocellular carcinoma. **Biochem Biophys Res Commun** 2004; 315(4): 950-8.

- Huang Q, Huang Q, Chen W, Wang L, Lin W, Lin J, et al. Identification of transgelin as a potential novel biomarker for gastric adenocarcinoma based on proteomics technology. **J Cancer Res Clin Oncol** 2008; 134(11): 1219-27.
- Imbeaud S, Graudens E, Boulanger V, Barlet X, Zaborski P, Eveno E, et al. Towards standardization of RNA quality assessment using user-independent classifiers of microcapillary electrophoresis traces. **Nucleic Acids Res** 2005; 33(6): 1-12.
- Ishihama Y, Oda Y, Tabata T, Sato T, Nagasu T, Rappsilber J, et al. Exponentially modified protein abundance index (emPAI) for estimation of absolute protein amount in proteomics by the number of sequenced peptides per protein. **Mol Cell Proteomics** 2005; 4(9): 1265-72.
- Jeong HC, Kim GI, Cho SH, Lee KH, Ko JJ, Yang JH, et al. Proteomic analysis of human small cell lung cancer tissues: up-regulation of coactosin-like protein-1. **J Proteome Res** 2011; 10(1): 269-76.
- Ji NY, Park MY, Kang YH, Lee CI, Kim DG, Yeom YI, et al. Evaluation of annexin II as a potential serum marker for hepatocellular carcinoma using a developed sandwich ELISA method. **Int J Mol Med** 2009; 24(6): 765-71.
- Jia H, Liu Y, Xia R, Tong C, Yue T, Jiang J, et al. Casein kinase 2 promotes Hedgehog signaling by regulating both smoothed and Cubitus interruptus. **J Biol Chem** 2010; 285(48): 37218-26.
- Jolodar A, Fischer P, Bergmann S, Buttner DW, Hammerschmidt S, Brattig NW. Molecular cloning of an alpha-enolase from the human filarial parasite *Onchocerca volvulus* that binds human plasminogen. **Biochim Biophys Acta** 2003; 1627(2): 111-20.
- Jongsuksuntigul P, Imsomboon T. Opisthorchiasis control in Thailand. **Acta Trop** 2003; 88(3): 229-32.
- Keshamouni VG, Michailidis G, Grasso CS, Anthwal S, Strahler JR, Walker A, et al. Differential protein expression profiling by iTRAQ-2DLC-MS/MS of lung cancer cells undergoing epithelial-mesenchymal transition reveals a migratory/invasive phenotype. **J Proteome Res** 2006; 5(5): 1143-54.

- Kim JH, Bogner PN, Ramnath N, Park Y, Yu J, Park YM. Elevated peroxiredoxin 1, but not NF-E2-related factor 2, is an independent prognostic factor for disease recurrence and reduced survival in stage I non-small cell lung cancer. **Clin Cancer Res** 2007; 13(13): 3875-82.
- Kim YJ, Lee WS, Ip C, Chae HZ, Park EM, Park YM. Prx1 suppresses radiation-induced c-Jun NH2-terminal kinase signaling in lung cancer cells through interaction with the glutathione S-transferase Pi/c-Jun NH2-terminal kinase complex. **Cancer Res** 2006; 66(14): 7136-42.
- Kischel P, Guillonneau F, Dumont B, Bellahcene A, Stresing V, Clezardin P, et al. Cell membrane proteomic analysis identifies proteins differentially expressed in osteotropic human breast cancer cells. **Neoplasia** 2008; 10(9): 1014-20.
- Kreimann EL, Morales FC, de Orbeta-Cruz J, Takahashi Y, Adams H, Liu TJ, et al. Cortical stabilization of beta-catenin contributes to NHERF1/EBP50 tumor suppressor function. **Oncogene** 2007; 26(36): 5290-9.
- Krijgsveld J, Gauci S, Dormeyer W, Heck AJ. In-gel isoelectric focusing of peptides as a tool for improved protein identification. **J Proteome Res** 2006; 5(7): 1721-30.
- Lawrie LC, Curran S, McLeod HL, Fothergill JE, Murray GI. Application of laser capture microdissection and proteomics in colon cancer. **Mol Pathol** 2001; 54(4): 253-8.
- Lazar CS, Cresson CM, Lauffenburger DA, Gill GN. The Na⁺/H⁺ exchanger regulatory factor stabilizes epidermal growth factor receptors at the cell surface. **Mol Biol Cell** 2004; 15(12): 5470-80.
- Lee EK, Han GY, Park HW, Song YJ, Kim CW. Transgelin promotes migration and invasion of cancer stem cells. **J Proteome Res** 2010; 9(10): 5108-17.
- Lengqvist J, Uhlen K, Lehtio J. iTRAQ compatibility of peptide immobilized pH gradient isoelectric focusing. **Proteomics** 2007; 7(11): 1746-52.
- Liu JW, Shen JJ, Tanzillo-Swartz A, Bhatia B, Maldonado CM, Person MD, et al. Annexin II expression is reduced or lost in prostate cancer cells and its re-expression inhibits prostate cancer cell migration. **Oncogene** 2003; 22(10): 1475-85.

- Lomnytska MI, Becker S, Hellman K, Hellstrom AC, Souchelnytskyi S, Mints M, et al. Diagnostic protein marker patterns in squamous cervical cancer. **Proteomics Clin Appl** 2010; 4(1): 17-31.
- Lu Q, Dobbs LJ, Gregory CW, Lanford GW, Revelo MP, Shappell S, et al. Increased expression of delta-catenin/neural plakophilin-related armadillo protein is associated with the down-regulation and redistribution of E-cadherin and p120ctn in human prostate cancer. **Hum Pathol** 2005; 36(10): 1037-48.
- _____, Paredes M, Medina M, Zhou J, Cavallo R, Peifer M, et al. Delta-catenin, an adhesive junction-associated protein which promotes cell scattering. **J Cell Biol** 1999; 144(3): 519-32.
- Mai J, Waisman DM, Sloane BF. Cell surface complex of cathepsin B/annexin II tetramer in malignant progression. **Biochim Biophys Acta** 2000; 1477(1): 215-30.
- Marchler-Bauer A, Anderson JB, Derbyshire MK, DeWeese-Scott C, Gonzales NR, Gwadz M, et al. CDD: a conserved domain database for interactive domain family analysis. **Nucleic Acids Res** 2007; 35(Database issue): D237-40.
- Maudsley S, Zamah AM, Rahman N, Blitzer JT, Luttrell LM, Lefkowitz RJ, et al. Platelet-derived growth factor receptor association with Na(+)/H(+) exchanger regulatory factor potentiates receptor activity. **Mol Cell Biol** 2000; 20(22): 8352-63.
- Merseburger AS, Hennenlotter J, Simon P, Ohneseit PA, Kuehs U, Kruck S, et al. Cathepsin D expression in renal cell cancer-clinical implications. **Eur Urol** 2005; 48(3): 519-26.
- Michel PE, Reymond F, Arnaud IL, Josserand J, Girault HH, Rossier JS. Protein fractionation in a multicompartiment device using Off-Gel isoelectric focusing. **Electrophoresis** 2003; 24(1): 3-11.
- Mohammad HS, Kurokohchi K, Yoneyama H, Tokuda M, Morishita A, Jian G, et al. Annexin A2 expression and phosphorylation are up-regulated in hepatocellular carcinoma. **Int J Oncol** 2008; 33(6): 1157-63.
- Molloy MP, Witzmann FA. Proteomics: technologies and applications. **Brief Funct Genomic Proteomic** 2002; 1(1): 23-39.

- Mortz E, Vorm O, Mann M, Roepstorff P. Identification of proteins in polyacrylamide gels by mass spectrometric peptide mapping combined with database search. **Biol Mass Spectrom** 1994; 23(5): 249-61.
- Naito S, Yokomizo A, Koga H. Mechanisms of drug resistance in chemotherapy for urogenital carcinoma. **Int J Urol** 1999; 6(9): 427-39.
- Nakanuma Y, Sripa B, Vatanasapt V, Leong A, Ponchon T, Ishak K. Intrahepatic cholangiocarcinoma. In: **Hamilton SR, Aaltonen LA, eds WHO Classification of Tumors, Pathology and Genetics Tumours of the Digestive System Lyon: IARC Press** 2000; 173-80.
- Nathan H, Pawlik TM, Wolfgang CL, Choti MA, Cameron JL, Schulick RD. Trends in survival after surgery for cholangiocarcinoma: a 30-year population-based SEER database analysis. **J Gastrointest Surg** 2007; 11(11): 1488-96.
- Nedjadi T, Kitteringham N, Campbell F, Jenkins RE, Park BK, Navarro P, et al. S100A6 binds to annexin 2 in pancreatic cancer cells and promotes pancreatic cancer cell motility. **Br J Cancer** 2009; 101(7): 1145-54.
- Noh DY, Ahn SJ, Lee RA, Kim SW, Park IA, Chae HZ. Overexpression of peroxiredoxin in human breast cancer. **Anticancer Res** 2001; 21(3B): 2085-90.
- Oliva D, Barba G, Barbieri G, Giallongo A, Feo S. Cloning, expression and sequence homologies of cDNA for human gamma enolase. **Gene** 1989; 79(2): 355-60.
- Olnes MJ, Erlich R. A review and update on cholangiocarcinoma. **Oncology** 2004; 66(3): 167-79.
- Olson MF, Sahai E. The actin cytoskeleton in cancer cell motility. **Clin Exp Metastasis** 2009; 26(4): 273-87.
- Ong SE, Blagoev B, Kratchmarova I, Kristensen DB, Steen H, Pandey A, et al. Stable isotope labeling by amino acids in cell culture, SILAC, as a simple and accurate approach to expression proteomics. **Mol Cell Proteomics** 2002; 1(5): 376-86.
- _____, Mann M. Mass spectrometry-based proteomics turns quantitative. **Nat Chem Biol** 2005; 1(5): 252-62.

- Pak JH, Moon JH, Hwang SJ, Cho SH, Seo SB, Kim TS. Proteomic analysis of differentially expressed proteins in human cholangiocarcinoma cells treated with *Clonorchis sinensis* excretory-secretory products. **J Cell Biochem** 2009; 108(6): 1376-88.
- Palmer-Toy DE, Sarracino DA, Sgroi D, LeVangie R, Leopold PE. Direct acquisition of matrix-assisted laser desorption/ionization time-of-flight mass spectra from laser capture microdissected tissues. **Clin Chem** 2000; 46(9): 1513-6.
- Pan Y, Weinman EJ, Dai JL. Na⁺/H⁺ exchanger regulatory factor 1 inhibits platelet-derived growth factor signaling in breast cancer cells. **Breast Cancer Res** 2008; 10(1): 1-12.
- Pancholi V. Multifunctional alpha-enolase: its role in diseases. **Cell Mol Life Sci** 2001; 58(7): 902-20.
- Pastorelli R, Saletta F, Carpi D, Campagna R, dell'Osta C, Schiarea S, et al. Proteome characterization of a human urothelial cell line resistant to the bladder carcinogen 4-aminobiphenyl. **Proteome Sci** 2007; 5(6): 1-9.
- Paweletz CP, Ornstein DK, Roth MJ, Bichsel VE, Gillespie JW, Calvert VS, et al. Loss of annexin 1 correlates with early onset of tumorigenesis in esophageal and prostate carcinoma. **Cancer Res** 2000; 60(22): 6293-7.
- Peng J, Elias JE, Thoreen CC, Licklider LJ, Gygi SP. Evaluation of multidimensional chromatography coupled with tandem mass spectrometry (LC/LC-MS/MS) for large-scale protein analysis: the yeast proteome. **J Proteome Res** 2003; 2(1): 43-50.
- Pierce A, Unwin RD, Evans CA, Griffiths S, Carney L, Zhang L, et al. Eight-channel iTRAQ enables comparison of the activity of six leukemogenic tyrosine kinases. **Mol Cell Proteomics** 2008; 7(5): 853-63.
- Pinlaor S, Hiraku Y, Ma N, Yongvanit P, Semba R, Oikawa S, et al. Mechanism of NO-mediated oxidative and nitrative DNA damage in hamsters infected with *Opisthorchis viverrini*: a model of inflammation-mediated carcinogenesis. **Nitric Oxide** 2004a; 11(2): 175-83.
- _____, Sripa B, Sithithaworn P, Yongvanit P. Hepatobiliary changes, antibody response, and alteration of liver enzymes in hamsters re-infected with *Opisthorchis viverrini*. **Exp Parasitol** 2004b; 108(1-2): 32-9.

- Pitteri S, Hanash S. A systems approach to the proteomic identification of novel cancer biomarkers. **Dis Markers** 2010; 28(4): 233-9.
- Pucci-Minafra I, Cancemi P, Fontana S, Minafra L, Feo S, Becchi M, et al. Expanding the protein catalogue in the proteome reference map of human breast cancer cells. **Proteomics** 2006; 6(8): 2609-25.
- _____, Fontana S, Cancemi P, Basirico L, Caricato S, Minafra S. A contribution to breast cancer cell proteomics: detection of new sequences. **Proteomics** 2002; 2(7): 919-27.
- Qi Y, Chiu JF, Wang L, Kwong DL, He QY. Comparative proteomic analysis of esophageal squamous cell carcinoma. **Proteomics** 2005; 5(11): 2960-71.
- Reczek D, Berryman M, Bretscher A. Identification of EBP50: A PDZ-containing phosphoprotein that associates with members of the ezrin-radixin-moesin family. **J Cell Biol** 1997; 139(1): 169-79.
- Redlitz A, Fowler BJ, Plow EF, Miles LA. The role of an enolase-related molecule in plasminogen binding to cells. **Eur J Biochem** 1995; 227(1): 407-15.
- Rhee SG, Kang SW, Chang TS, Jeong W, Kim K. Peroxiredoxin, a novel family of peroxidases. **IUBMB Life** 2001; 52(1): 35-41.
- Rodova M, Kelly KF, VanSaun M, Daniel JM, Werle MJ. Regulation of the rapsyn promoter by kaiso and delta-catenin. **Mol Cell Biol** 2004; 24(16): 7188-96.
- Rodrigo Tapia JP, Pena Alonso E, Garcia-Pedrero JM, Florentino Fresno M, Suarez Nieto C, Owen Morgan R, et al. Annexin A2 expression in head and neck squamous cell carcinoma. **Acta Otorrinolaringol Esp** 2007; 58(6): 257-62.
- Ross PL, Huang YN, Marchese JN, Williamson B, Parker K, Hattan S, et al. Multiplexed protein quantitation in *Saccharomyces cerevisiae* using amine-reactive isobaric tagging reagents. **Mol Cell Proteomics** 2004; 3(12): 1154-69.
- Ruan Y, Wan M. An optimized procedure for solubilization, reduction, and transfer of human breast cancer membrane-enriched fraction by 2-DE. **Electrophoresis** 2007; 28(18): 3333-40.
- Sardana G, Marshall J, Diamandis EP. Discovery of candidate tumor markers for prostate cancer via proteomic analysis of cell culture-conditioned medium. **Clin Chem** 2007; 53(3): 429-37.

- Shibata T, Chuma M, Kokubu A, Sakamoto M, Hirohashi S. EBP50, a beta-catenin-associating protein, enhances Wnt signaling and is over-expressed in hepatocellular carcinoma. **Hepatology** 2003; 38(1): 178-86.
- Shilov IV, Seymour SL, Patel AA, Loboda A, Tang WH, Keating SP, et al. The Paragon Algorithm, a next generation search engine that uses sequence temperature values and feature probabilities to identify peptides from tandem mass spectra. **Mol Cell Proteomics** 2007; 6(9): 1638-55.
- Shimonishi T, Miyazaki K, Nakanuma Y. Cytokeratin profile relates to histological subtypes and intrahepatic location of intrahepatic cholangiocarcinoma and primary sites of metastatic adenocarcinoma of liver. **Histopathology** 2000; 37(1): 55-63.
- Simamura E, Hirai K, Shimada H, Koyama J, Niwa Y, Shimizu S. Furanonaphthoquinones cause apoptosis of cancer cells by inducing the production of reactive oxygen species by the mitochondrial voltage-dependent anion channel. **Cancer Biol Ther** 2006; 5(11): 1523-9.
- Simone NL, Paweletz CP, Charboneau L, Petricoin EF, 3rd, Liotta LA. Laser capture microdissection: beyond functional genomics to proteomics. **Mol Diagn** 2000a; 5(4): 301-7.
- _____, Remaley AT, Charboneau L, Petricoin EF, 3rd, Glickman JW, Emmert-Buck MR, et al. Sensitive immunoassay of tissue cell proteins procured by laser capture microdissection. **Am J Pathol** 2000b; 156(2): 445-52.
- Song J, Bai J, Yang W, Gabrielson EW, Chan DW, Zhang Z. Expression and clinicopathological significance of oestrogen-responsive ezrin-radixin-moesin-binding phosphoprotein 50 in breast cancer. **Histopathology** 2007; 51(1): 40-53.
- Sousa LP, Brasil BS, Silva Bde M, Nogueira SV, Andrade AA, Ferreira PC, et al. Characterization of alpha-enolase as an interferon-alpha 2 alpha 1 regulated gene. **Front Biosci** 2005a; 337(4): 1065-71.
- _____, Silva BM, Brasil BS, Nogueira SV, Ferreira PC, Kroon EG, et al. Plasminogen/plasmin regulates alpha-enolase expression through the MEK/ERK pathway. **Biochem Biophys Res Commun** 2005b; 337(4): 1065-71.

- Sriamporn S, Parkin DM, Pisani P, Vatanasapt V, Suwanrungruang K, Kamsa-ard P, et al. A prospective study of diet, lifestyle, and genetic factors and the risk of cancer in Khon Kaen Province, northeast Thailand: description of the cohort. **Asian Pac J Cancer Prev** 2005; 6(3): 295-303.
- _____, Pisani P, Pipitgool V, Suwanrungruang K, Kamsa-ard S, Parkin DM. Prevalence of *Opisthorchis viverrini* infection and incidence of cholangiocarcinoma in Khon Kaen, Northeast Thailand. **Trop Med Int Health** 2004; 9(5): 588-94.
- Sripa B, Kaewkes S, Sithithaworn P, Mairiang E, Laha T, Smout M, et al. Liver fluke induces cholangiocarcinoma. **PLoS Med** 2007; 4(7): e201.
- _____, Leungwattanawanit S, Nitta T, Wongkham C, Bhudhisawasdi V, Puapairoj A, et al. Establishment and characterization of an opisthorchiasis-associated cholangiocarcinoma cell line (KKU-100). **World J Gastroenterol** 2005; 11(22): 3392-7.
- _____, Pairojkul C. Cholangiocarcinoma: lessons from Thailand. **Curr Opin Gastroenterol** 2008; 24(3): 349-56.
- _____, Sithithaworn P, Sirisinha S. *Opisthorchis viverrini* and opisthorchiasis: the 21st century review. **Acta Trop** 2003; 88(3): 169-70.
- Srisomsap C, Sawangareetrakul P, Subhasitanont P, Chokchaichamnankit D, Chiablaem K, Bhudhisawasdi V, et al. Proteomic studies of cholangiocarcinoma and hepatocellular carcinoma cell secretomes. **J Biomed Biotechnol** 2010; 73(12) 2291-305.
- Srisomsap C, Sawangareetrakul P, Subhasitanont P, Panichakul T, Keeratichamroen S, Lirdprapamongkol K, et al. Proteomic analysis of cholangiocarcinoma cell line. **Proteomics** 2004; 4(4): 1135-44.
- Stemmer-Rachamimov AO, Wiederhold T, Nielsen GP, James M, Pinney-Michalowski D, Roy JE, et al. NHE-RF, a merlin-interacting protein, is primarily expressed in luminal epithelia, proliferative endometrium, and estrogen receptor-positive breast carcinomas. **Am J Pathol** 2001; 158(1): 57-62.

- Sun S, Xu MZ, Poon RT, Day PJ, Luk JM. Circulating Lamin B1 (LMNB1) biomarker detects early stages of liver cancer in patients. **J Proteome Res** 2010; 9(1): 70-8.
- Takahashi Y, Morales FC, Kreimann EL, Georgescu MM. PTEN tumor suppressor associates with NHERF proteins to attenuate PDGF receptor signaling. **EMBO J** 2006; 25(4): 910-20.
- Tan TL, Chen WN. A proteomics analysis of cellular proteins associated with HBV genotype-specific HBX: potential in identification of early diagnostic markers for HCC. **J Clin Virol** 2005; 33(4): 293-8.
- Tsai ST, Chien IH, Shen WH, Kuo YZ, Jin YT, Wong TY, et al. ENO1, a potential prognostic head and neck cancer marker, promotes transformation partly via chemokine CCL20 induction. **Eur J Cancer** 2010; 46(9): 1712-23.
- Tzivion G, Gupta VS, Kaplun L, Balan V. 14-3-3 proteins as potential oncogenes. **Semin Cancer Biol** 2006; 16(3): 203-13.
- Uttaravichien T, Bhudhisawasdi V, Pairojkul C, Pugkhem A. Intrahepatic cholangiocarcinoma in Thailand. **J Hepatobiliary Pancreat Surg** 1999; 6(2): 128-35.
- van Diest PJ, van Dam P, Henzen-Logmans SC, Berns E, van der Burg ME, Green J, et al. A scoring system for immunohistochemical staining: consensus report of the task force for basic research of the EORTC-GCCG. European Organization for Research and Treatment of Cancer-Gynaecological Cancer Cooperative Group. **J Clin Pathol** 1997; 50(10): 801-4.
- Waisman DM. Annexin II tetramer: structure and function. **Mol Cell Biochem** 1995; 149-150: 301-22.
- Wang AG, Yoon SY, Oh JH, Jeon YJ, Kim M, Kim JM, et al. Identification of intrahepatic cholangiocarcinoma related genes by comparison with normal liver tissues using expressed sequence tags. **Biochem Biophys Res Commun** 2006; 345(3): 1022-32.
- Wang B, Yang Y, Friedman PA. Na/H exchange regulatory factor 1, a novel AKT-associating protein, regulates extracellular signal-regulated kinase signaling through a B-Raf-mediated pathway. **Mol Biol Cell** 2008; 19(4): 1637-45.

- Wang Y, Geppert H, Bajorath J. Shannon entropy-based fingerprint similarity search strategy. **J Chem Inf Model** 2009; 49(7): 1687-91.
- Watanapa P. Cholangiocarcinoma in patients with opisthorchiasis. **Br J Surg** 1996; 83(8): 1062-64.
- Wen F, Shen A, Shanas R, Bhattacharyya A, Lian F, Hostetter G, et al. Higher expression of the heterogeneous nuclear ribonucleoprotein k in melanoma. **Ann Surg Oncol** 2010; 17(10): 2619-27.
- Wertz A, Borst A, Haag J. Nonlinear integration of binocular optic flow by DNOVS2, a descending neuron of the fly. **J Neurosci** 2008; 28(12): 3131-40.
- Westley B, Rochefort H. A secreted glycoprotein induced by estrogen in human breast cancer cell lines. **Cell** 1980; 20(2): 353-62.
- Westley BR, May FE. Cathepsin D and breast cancer. **Eur J Cancer** 1996; 32(1): 15-24.
- Wolters DA, Washburn MP, Yates JR, 3rd. An automated multidimensional protein identification technology for shotgun proteomics. **Anal Chem** 2001; 73(23): 5683-90.
- Yanagawa T, Ishikawa T, Ishii T, Tabuchi K, Iwasa S, Bannai S, et al. Peroxiredoxin I expression in human thyroid tumors. **Cancer Lett** 1999; 145(1): 127-32.
- Yanay O, Vaughan DJ, Diab M, Brownstein D, Brogan TV. Retained wooden foreign body in a child's thigh complicated by severe necrotizing fasciitis: a case report and discussion of imaging modalities for early diagnosis. **Pediatr Emerg Care** 2001; 17(5): 354-5.
- Yonglitthipagon P, Pairojkul C, Chamgramol Y, Mulvenna J, Sripa B. Up-regulation of annexin A2 in cholangiocarcinoma caused by *Opisthorchis viverrini* and its implication as a prognostic marker. **Int J Parasitol** 2010; 40(10): 1203-12.
- Yu LR, Conrads TP, Uo T, Kinoshita Y, Morrison RS, Lucas DA, et al. Global analysis of the cortical neuron proteome. **Mol Cell Proteomics** 2004; 3(9): 896-907.
- Zeng Y, Abdallah A, Lu JP, Wang T, Chen YH, Terrian DM, et al. Delta-catenin promotes prostate cancer cell growth and progression by altering cell cycle and survival gene profiles. **Mol Cancer** 2009; 8(19): 1-8.

- Zhang F, Zhang L, Zhang B, Wei X, Yang Y, Qi RZ, et al. Anxa2 plays a critical role in enhanced invasiveness of the multidrug resistant human breast cancer cells. **J Proteome Res** 2009; 8(11): 5041-7.
- Zhang JY, Wang Y, Zhang D, Yang ZQ, Dong XJ, Jiang GY, et al. Delta-catenin promotes malignant phenotype of non-small cell lung cancer by non-competitive binding to E-cadherin with p120ctn in cytoplasm. **J Pathol** 2010; 222(1): 76-88.
- Zhang Y, Li M, Liu Y, Han N, Zhang K, Xiao T, et al. ENO1 protein levels in the tumor tissues and circulating plasma samples of non-small cell lung cancer patients. **Zhongguo Fei Ai Za Zhi** 2010; 13(12): 1089-93.
- Zhong L, Peng X, Hidalgo GE, Doherty DE, Stromberg AJ, Hirschowitz EA. Antibodies to HSP70 and HSP90 in serum in non-small cell lung cancer patients. **Cancer Detect Prev** 2003; 27(4): 285-90.
- Zhuang PY, Zhang JB, Zhu XD, Zhang W, Wu WZ, Tan YS, et al. Two pathologic types of hepatocellular carcinoma with lymph node metastasis with distinct prognosis on the basis of CK19 expression in tumor. **Cancer** 2008; 112(12): 2740-8.

APPENDICES

Reagents for laboratory experiments

I. Reagents for cell culture

1. Complete RPMI-1640 medium

RPMI-1640 (Gibco BRL)		1 pack
NaHCO ₃		1.176 g
Adjust pH to 7.4 by HCl		
Distilled water	to	1 liter
Filter through 0.22- μ m filter		
Filtered RPMI-1640		100 ml
Fetal bovine serum		10 ml
Penicillin (200,000 U/ml)		50 μ l
Streptomycin (200,000 μ g/ml)		50 μ l
Store at 4°C.		

2. Antibiotics

a. Penicillin 200,000 U/ml

Penicillin		1,000,000 U
Sterile distilled water	to	5 ml
Store at -20°C.		

b. Streptomycin (200,000 μ g/ml)

Streptomycin		1 g
Sterile distilled water	to	5 ml
Store at -20°C.		

3. 0.25% Trypsin-EDTA

2.5% Trypsin solution (Gibco BRL)		2 ml
Versine EDTA (Gibco BRL)		50 ml
Filter through 0.22- μ m filter.		
Store at 4°C.		

4. Phosphate buffered saline (PBS) pH 7.4 for cell culture

NaCl	8.0 g
KCl	0.2 g
KH ₂ PO ₄	0.2 g
Na ₂ HPO ₄	1.15 g

Note: Adjust pH to 7.4 by NaOH, add distilled water to final volume of 1 liter. Sterilized by autoclaving at 121°C, 15 pounds per square inch for 15 min and store at room temperature.

II. Reagents for Proteomics

1. Sample preparation solution with urea and thiourea

Urea	10.5 g
Thiourea	3.8 g
CHAPS ^a	1 g
IPG Buffer ^b	500 µl
DTT	154 mg
Double-distilled water	to 25 ml

Note: ^aOther neutral or zwitterionic detergents may be used at concentrations up to 2% (w/v). ^bUse IPG Buffer in the pH range corresponding to the pH range of the IEF separation to be performed.

2. Thiourea rehydration stock solution

Urea	10.5 g
Thiourea	3.8 g
CHAPS	0.5 g
IPG Buffer	125 µl
1% Bromophenol blue stock solution	50 µl
Double-distilled water	to 25 ml

Note: DTT is added just prior to use: add 7 mg DTT per 2.5-ml aliquot of rehydration stock solution. Store in 2.5-ml aliquots at -20°C.



3. SDS equilibration buffer solution

Urea		72.1 g
Tris-HCL, pH 8.8		10 ml
Glycerol (87% w/w)		69 ml
SDS		4 g
Bromophenol blue stock solution		400 μ l
Double-distilled water	to	200 ml

Note: This is a stock solution. Just prior to use, add DTT or iodoacetamide as described in the protocol.

4. 10X Laemmli SDS electrophoresis buffer

Trisbase		303 g
Glycine		1441 g
SDS		100 g
Distilled water	to	10 liters

Note: The pH of this solution should not be adjusted.

5. 30% T, 2.6% C monomer stock solution

Acrylamide		300 g
N,N'-methylenebisacrylamide		8 g
Distilled water	to	1 liter

Note: Filter solution through a 0.45- μ m filter. Store at 4°C in the dark.

6. 4X resolving gel buffer solution

Tris base		181.7 g
Distilled water	to	750 ml
HCL		adjust to pH 8.8
Distilled water	to	1 liter

Note: Filter solution through a 0.45- μ m filter. Store at 4°C.

7. Bromophenol blue stock solution

Bromophenol blue		100 mg
Tris-base		60 mg
Distilled water	to	10 ml

8. 10% SDS

SDS, electrophoresis grade		5 g
Distilled water	to	50 ml

Note: Filter solution through a 0.45- μ m filter. Store at RT.

III. Reagents for Western blot**1. 5X Stock Transfer buffer**

Tris base		15.1425 g
Glycine		72.0672 g
Distilled water	to	1 liter
pH is adjusted at 8.4		

2. 1X Working Transfer buffer

5X Stock Transfer buffer		600 ml
Methanol		600 ml
Distilled water	to	3 liters

3. 1X phosphate-buffered saline with 0.05% Tween-20 (PBST)

10 X PBS		100 ml
Tween-20		500 μ l
Distilled water	to	1 liter

4. 5% skimmed milk in PBST

Instant nonfat dry milk		5 g
1X PBST		100 ml

5. 2% skimmed milk in PBST

Instant nonfat dry milk		2 g
1X PBST		100 ml

IV. Reagents for immunohistochemistry**1. 0.01M Citrate buffer pH 6.0**

Citric acid		4.2 g
Dissolve and adjust to pH 6.0 with NaOH		
Distilled water	to	2 liter

2. 0.05M Tris buffer pH 7.6 for DAB

Tris base		6.1 g
Distilled water	to	800 ml
Dissolve and adjust to pH 7.6 with NaOH		
Distilled water	to	1 liter

3. PBS for diluting with 0.05% NaN₃

NaN ₃		0.05 g
Distilled water	to	100 ml

4. Stock DAB (25 mg/ml)

DAB		1 g
Distilled water	to	40 ml
Aliquot 1 ml to each microtubes.		

5. 10X phosphate buffered saline (PBS)

NaCl		40 g
KCl		1 g
Na ₂ HPO ₄		7.2 g
KH ₂ PO ₄		1.2 g
Distilled water	to	500 ml
pH is adjusted at 7.4		

6. 1X PBS

10X PBS		100 ml
Distilled water	to	1 liter

Table A.1 *Opisthorchis viverrini*-induced cholangiocarcinoma cell lines and normal biliary cell line.

Cell lines	Description
OCA17	Well differentiated adenocarcinoma CCA cell line
M156	Moderately differentiated adenocarcinoma CCA cell line
K100	Poorly differentiated adenocarcinoma CCA cell line
M139	Squamous cell carcinoma CCA cell line
M213	Adenosquamous cell carcinoma CCA cell line
H69	Immortalized normal cholangiocyte cell line

Note: CCA cell lines were established at Division of Experimental Pathology, Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand. Immortalized cholangiocyte cell line H69 was kindly provided by Dr. Gregory J. Gores, Mayo clinic, Rochester, MN, USA.

Table A.2 Immunohistochemical grading of ANXA2, PRX1, EBP50 and ENO1 proteins in human CCAs.

No.	Code	Immunohistochemical Grading			
		ANXA2 ^a	PRX1 ^b	EBP50 ^b	ENO1 ^b
1	40-059	1	0	1	1
2	40-108	1	0	1	1
3	40-116	1	1	1	1
4	41-122	1	0	0	1
5	41-123	1	0	1	1
6	M005	0	1	0	1
7	M009	1	1	1	0
8	M030	1	1	0	0
9	M040	0	1	1	0
10	M037	1	0	1	1
11	M043	1	1	0	0
12	M055	1	0	0	1
13	M068	1	1	1	0
14	M071	1	0	0	1
15	M079	1	0	1	1
16	M081	1	0	1	1
17	M082	0	1	1	0
18	M090	1	0	1	1
19	M101	0	1	1	0
20	M102	1	0	1	1
21	M111	1	1	0	1
22	M119	0	1	1	0
23	M132	1	0	1	1
24	M136	0	1	1	0
25	M138	0	1	0	0
26	M140	0	0	1	1
27	M148	0	0	1	1
28	M152	0	1	1	0
29	M155	0	1	1	0
30	M156	0	1	1	0
31	M157	0	0	1	1
32	M171	1	0	1	1
33	M187	1	0	1	1
34	M209	0	1	1	0
35	M213	1	0	0	1
36	M214	1	0	1	1
37	M229	1	1	1	1
38	M240	0	1	1	0
39	M249	1	0	1	1
40	M250	0	1	0	1
41	M259	1	0	0	1
42	M265	1	0	1	1
43	M270	1	1	1	1
44	M274	1	1	0	0
45	M276	0	1	1	0
46	M279	0	0	1	1
47	M283	1	0	0	1
48	M286	1	0	0	1
49	M287	1	0	1	1

Table A.2 Immunohistochemical grading of ANXA2, PRX1, EBP50 and ENO1 proteins in human CCAs (Cont.)

No.	Code	Immunohistochemical Grading			
		ANXA2 ^a	PRX1 ^b	EBP50 ^b	ENO1 ^b
50	M294	0	1	1	0
51	M298	1	0	1	1
52	M303	1	0	1	1
53	M311	0	1	1	0
54	N013	1	0	0	1
55	N017	0	0	1	1
56	N019	0	1	1	0
57	N021	1	1	0	0
58	N022	1	0	1	1
59	N023	1	1	0	0
60	N024	0	1	0	0
61	N029	1	0	0	1
62	N044	1	0	1	1
63	N046	1	1	0	0
64	N048	1	0	1	1
65	N052	1	0	0	1
66	N069	1	0	1	1
67	N059	1	1	1	1
68	N082	1	0	1	1
69	N088	1	0	1	1
70	N091	1	0	0	1
71	N104	1	0	1	1
72	P066	0	1	1	0
73	N105	1	0	0	1
74	N108	1	0	0	1
75	N110	1	1	1	0
76	N112	1	0	0	1
77	N114	1	0	1	1
78	N127	0	1	0	0
79	N128	1	1	0	1
80	N136	1	0	1	1
81	N140	1	1	1	1
82	N146	1	1	1	1
83	N154	0	1	1	0
84	N157	1	0	1	1
85	N158	1	0	1	1
86	P006	0	1	0	0
87	P011	1	0	1	1
88	P020	1	0	1	1
89	P029	1	0	0	1
90	P031	1	0	1	1
91	P040	1	0	1	1
92	P048	1	0	1	1
93	P055	1	0	1	1
94	P056	0	1	1	0
95	P059	1	0	1	1
96	P060	1	0	0	1
97	P068	1	0	0	1
98	P071	0	1	0	0
99	P077	1	0	1	1
100	P086	1	0	1	1

Table A.2 Immunohistochemical grading of ANXA2, PRX1, EBP50 and ENO1 proteins in human CCAs (Cont.)

No.	Code	Immunohistochemical Grading			
		ANXA2 ^a	PRX1 ^b	EBP50 ^b	ENO1 ^b
101	P090	1	0	0	1
102	P091	0	1	1	0
103	P092	1	0	1	1
104	P093	1	0	1	1
105	P094	1	0	1	1
106	P095	0	1	1	0
107	P112	1	1	0	1
108	P126	0	1	0	0
109	P120	1	0	1	1
110	P127	1	0	1	1
111	N135	1	0	0	1
112	P145	1	0	0	1
113	P147	1	1	1	1
114	P149	1	0	0	1
115	P152	1	0	1	1
116	Q006	1	0	1	1
117	Q007	1	0	1	1
118	Q008	1	1	0	1
119	Q013	1	1	0	1
120	Q015	1	0	1	1
121	Q021	1	0	1	1
122	Q023	1	0	1	1
123	Q025	1	0	0	1
124	Q036	1	0	0	1
125	Q037	0	1	0	0
126	Q041	1	0	0	1
127	Q043	1	0	1	1
128	Q045	0	1	0	0
129	Q048	1	0	0	1
130	Q053	1	0	1	1
131	Q058	1	0	0	1
132	Q062	1	0	1	1
133	Q063	0	1	0	0
134	Q066	1	1	1	1
135	Q071	1	0	1	1
136	Q074	1	1	1	1
137	Q089	0	1	1	0
138	Q093	1	0	1	1
139	Q095	1	0	1	1
140	Q099	1	1	1	1
141	Q101	1	0	0	1
142	Q102	1	0	1	1
143	Q103	1	0	1	1
144	Q106	1	0	1	1
145	Q111	1	0	1	1
146	Q114	1	0	1	1
147	Q123	1	0	0	1
148	Q126	1	0	1	1
149	R001	0	1	0	0

Table A.2 Immunohistochemical grading of ANXA2, PRX1, EBP50 and ENO1 proteins in human CCAs (Cont.)

No.	Code	Immunohistochemical Grading			
		ANXA2 ^a	PRX1 ^b	EBP50 ^b	ENO1 ^b
150	R003	0	1	0	0
151	R004	1	0	0	1
152	R009	1	0	1	1
153	R012	1	0	0	1
154	R023	1	0	0	1
155	R025	0	1	0	0
156	R026	1	0	0	1
157	R031	0	1	0	0
158	R038	1	0	1	1
159	R043	1	0	1	1
160	R049	1	0	1	1
161	R50	1	0	0	1
162	R052	0	1	0	0
163	R053	1	0	0	1
164	R054	1	0	0	1
165	R061	1	0	0	1
166	R062	1	0	1	1
167	R066	1	0	0	1
168	R074	0	1	0	0
169	R077	1	0	0	1
170	R078	0	1	1	0
171	R081	1	0	0	1
172	R085	0	1	1	0
173	R086	1	0	0	1
174	R087	1	0	1	1
175	R092	1	0	0	1
176	R093	1	0	1	1
177	R096	0	1	0	0
178	R100	1	0	1	1
179	R101	1	0	0	1
180	R103	1	1	0	1
181	R104	1	0	1	1
182	R105	1	0	0	1
183	R106	0	1	1	0
184	R118	1	0	1	1
185	R121	1	0	1	1
186	R123	0	1	1	0
187	R128	1	0	0	1
188	R133	1	1	1	1
189	R134	1	0	0	1
190	R137	1	0	0	1
191	R138	1	0	0	1
192	R141	0	1	1	0
193	R144	1	0	1	1
194	R145	0	1	0	0
195	R149	1	0	1	1
196	R155	1	0	1	1
197	R156	1	0	1	1
198	T001	0	1	0	0
199	T003	0	1	0	0
200	T005	1	0	1	1

Table A.2 Immunohistochemical grading of ANXA2, PRX1, EBP50 and ENO1 proteins in human CCAs (Cont.)

No.	Code	Immunohistochemical Grading			
		ANXA2 ^a	PRX1 ^b	EBP50 ^b	ENO1 ^b
201	T006	0	1	1	0
202	T010	0	1	1	0
203	T015	1	0	0	1
204	T016	1	0	0	1
205	T017	0	1	1	0
206	T026	1	0	1	1
207	T036	1	0	0	1
208	T041	0	1	0	0
209	T042	1	0	0	1
210	T043	1	0	1	1
211	T046	1	0	1	1
212	T049	0	1	0	0
213	T051	1	0	1	1
214	T058	0	1	1	0
215	T068	1	0	1	1
216	T071	1	0	0	1
217	T077	0	1	0	0
218	T079	1	0	0	1
219	T090	1	0	1	1
220	T092	1	0	1	1
221	T102	1	0	0	1
222	T117	1	0	0	1
223	T124	1	0	0	1
224	T151	1	0	1	1
225	T160	0	1	1	0
226	T162	1	0	0	1
227	T165	1	0	1	1
228	T168	1	0	0	1
229	T174	1	0	1	1
230	U004	1	0	1	1
231	U005	1	0	0	1
232	U010	0	1	0	0
233	U018	1	1	1	1
234	U019	1	0	0	1
235	U027	1	0	1	1
236	U028	0	1	1	0
237	U030	1	0	0	1
238	U034	1	0	0	1
239	U041	1	0	1	1
240	U044	1	1	0	1
241	U051	0	1	0	0
242	U052	1	0	0	1
243	U054	0	1	1	0
244	U059	1	0	1	1
245	U090	1	0	1	1
246	U107	0	1	0	0
247	U109	1	0	1	1
248	U110	1	0	0	1
249	U120	1	0	1	1
250	U122	1	0	1	1
251	U126	1	1	1	1

Table A.2 Immunohistochemical grading of ANXA2, PRX1, EBP50 and ENO1 proteins in human CCAs (Cont.)

No.	Code	Immunohistochemical Grading			
		ANXA2 ^a	PRX1 ^b	EBP50 ^b	ENO1 ^b
252	U137	1	0	1	1
253	W002	0	1	1	0
254	W004	1	1	1	1
255	W005	1	0	1	1
256	W008	0	1	1	0
257	W011	0	1	0	0
258	W012	1	0	0	1
259	W016	1	1	0	1
260	W017	1	0	0	1
261	W022	1	0	1	1
262	W025	1	0	1	1
263	W027	1	1	1	1
264	W035	1	0	1	1
265	W039	1	1	1	1
266	W040	1	0	0	1
267	W048	1	0	1	1
268	W049	1	0	1	1
269	W050	1	1	0	1
270	W056	1	0	1	1
271	W060	1	0	1	1
272	W061	1	0	1	1
273	W063	0	1	0	0
274	W068	0	1	0	0
275	W069	1	0	1	1
276	W077	1	0	0	1
277	W080	0	1	1	0
278	W084	1	0	0	1
279	W085	1	0	0	1
280	W087	1	0	0	1
281	W090	1	0	1	1
282	W092	1	0	1	1
283	W093	1	0	1	1
284	W100	1	0	1	1
285	W128	0	1	1	0
286	W130	1	0	1	1
287	W132	1	0	0	1
288	W147	0	1	1	0
289	W149	1	0	1	1
290	W151	1	0	1	1
291	W153	1	0	1	1
292	W164	1	0	0	1
293	W169	1	0	1	1
294	W171	1	1	1	1
295	W176	1	0	1	1
296	W181	1	0	0	1
297	W187	1	0	1	1
298	W195	1	0	1	1
299	W196	1	0	1	1
300	OCA17	0	1	0	0

Note: '0' = low expression group (positive staining cells \leq 10%); '1' = high expression group (positive staining cells >10%). ^aMembrane protein, ^bCytosolic proteins.

RESEARCH PUBLICATIONS

1. **Yonglitthipagon P**, Pairojkul C, Chamgramol Y, Mulvenna J, Sripa B. Up-regulation of annexin A2 in cholangiocarcinoma caused by *Opisthorchis viverrini* and its implication as a prognostic marker. **Int J Parasitol** 2010; 40: 1203-1212.
2. **Yonglitthipagon P**, Pairojkul C, Chamgramol Y, Mulvenna J, Loukas A, Bethony J, Sripa B. Prognostic significance of peroxiredoxin 1 and ezrin-radixin-moesin-binding phosphoprotein 50 in cholangiocarcinoma. (*submitted to Human Pathol*).
3. **Yonglitthipagon P**, Pairojkul C, Chamgramol Y, Mulvenna J, Loukas A, Sripa B. Proteomics-based identification of enolase 1 as a potential prognostic marker in cholangiocarcinoma. (*submitted to Clin Biochem*).

CURRICULUM VITAE



Name: Mr. Ponlapat Yonglitthipagon

Date of birth: June 12, 1983

Academic background:

- 2002-2005 Bachelor of Science (Physiotherapy), Faculty of Associated Medical Science. Khon Kaen University, Khon Kaen, Thailand.
- 2006-2011 Doctor of Philosophy in Pathobiology, Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand.

Training and visiting scholar:

- 2009 Application of laser capture microdissection and proteomics in cholangiocarcinoma, Queensland Institute of Medical Research (Brisbane), Queensland, Australia.
- 2010 Quantitative proteomics analysis of cholangiocarcinoma cell lines using OFFGEL, iTRAQ labeling and mass spectrometry (LC-MS/MS), Queensland Tropical Health Alliance, James Cook University (Cairns), Australia.

Honors:

- 2005 First class honor (Physiotherapy), Khon Kaen University, Khon Kaen, Thailand.
- 2006-2011 Royal Golden Jubilee Ph.D. program (RGJ-PhD) scholarship from Thailand Research Fund.

Outstanding poster presentation awards:

- 2008 RGJ-Ph.D. Congress IX, April 4-6, 2008. Jomtien Palm Beach Resort Pattaya, Chonburi, Thailand.
- 2011 Oncology Biomarkers Conference. January 27-28, 2011. Paradise Point Resort and Spa, San Diego, CA, USA.

Fields of interest: Proteomics, Tumor markers, Pathology.

