



เอกสารอ้างอิง

- วัชร กุณกิตติ, จันทนา อารมย์ดี, สุวรรณ วรัตน์, ผดุงขวัญ จิตโรภาส. 2549. การพัฒนาตำรับไลชันกันแดดที่มีแป้งขนุนเป็นสารเพิ่มความหนืดและการประเมินความพึงพอใจของอาสาสมัครต่อผลิตภัณฑ์. *Songklanakarin Journal of Science and Technology*, 28, 1-8.
- วิชชา บัวชุม, ชีพร กงบังเกิด. 2545. การเตรียมฟิล์มอะมัยโลสที่รับประทานได้จากแป้งเมล็ดขนุนและแนวทางการใช้ประโยชน์. *เกษตรนเรศวร*, 5, 21-31.
- อรอนงค์ กิตติพงษ์พัฒนา, นิสิต กิตติพงษ์พัฒนา, สุคนธ์ จันทร์ดี และ กรรณิกา เทียรณินธิกุล. (2554) แป้งจากเมล็ดขนุน: การดัดแปร สมบัติทางเคมีกายภาพ และการประยุกต์ใช้ทางเภสัชกรรม. รายงานวิจัยฉบับสมบูรณ์. คณะเภสัชศาสตร์ มหาวิทยาลัยเชียงใหม่. 160 หน้า.
- Adebowale, K.O., Afolabi, T.A., & Olu-Owolabi, B.L. (2005). Hydrothermal treatments of finger millet (*Eleusine coracana*) starch. *Food Hydrocolloids*, 19, 974-983.
- Adebowale, K.O., Henle, T., Schwarzenbolz, U., & Doert, T. (2009). Modification and properties of African yam bean (*Sphenostylis stenocarpa* Hochst. Ex A. Rich) Harms starch I: Heat moisture treatments and annealing. *Food Hydrocolloids*, 23, 1947-1957.
- Ambigaipalan, P., Hoover, R., Donner, E., & Liu, Q. (2014) Starch chain interactions within amorphous and crystalline domains of pulse starches during heat-moisture treatment at different temperatures and their impact on physicochemical properties. *Food Chemistry*, 143, 175-184.
- Bobbio, F.O., El-Dash, A.A., Bobbio, P.A., & Rodrigues, L.R. (1978). Isolation and characterization of the physicochemical properties of the starch of jackfruit seeds (*Artocarpus heterophyllus*). *Cereal Chemistry*, 55, 505-11.
- Chen, L., Li, X., Pang, Y., Li, L., Zhang, X., & Yu, L. (2007). Resistant starch as a carrier for oral colon-targeting drug matrix system. *Journal of Material Sciences: Material Medicine*, 18, 2199-2203.
- Chen, F., Zhang, B., & Lu, H. (2009). Jackfruit seeds starch digestibility and its dextrin properties. *Shipin Yu Fajiao Gongye*, 35, 43-46.
- Chung, H.J., Hoover, R., & Liu, Q. (2009a). The impact of single and dual hydrothermal modifications on the molecular structure and physicochemical properties of normal corn starch. *International Journal of Biological Macromolecules*, 44, 203-210.



- Chung, H.J., Liu, Q., & Hoover, R. (2009b). Impact of annealing and heat-moisture treatment on rapidly digestible, slowly digestible and resistant starch levels in native and gelatinized corn, pea and lentil starches. *Carbohydrate Polymers*, 75, 436-447.
- Dutta, H., Paul, S.K., Kalita, D., & Mahanta, C.L. (2011). Effect of acid concentration and treatment on acid-alcohol modified jackfruit seed starch properties. *Food Chemistry*, 128, 284-291.
- Eerlingen, R.C., Jacobs, H., Van Win, H., & Delcour, J.A. (1996). Effects of hydrothermal treatment on the gelatinization properties of potato starch as measured by differential scanning calorimetry. *Journal of Thermal Analysis*, 47, 1229-1246.
- Fuentes-Zaragoza, E., Riquelme-Navarrere, M.J., Sanchez-Zapata, E., Perez-Alvarez, & J.A. (2010). Resistant starch as functional ingredient: A review. *Food Research International*, 43, 931-942.
- Gunaratne, A. & Hoover, R. (2002). Effect of heat-moisture treatment on the structure and physicochemical properties of tuber and root starches. *Carbohydrate Polymers*, 49, 425-437.
- Hasjim, J., Ai, Y., & Jane, J.L. (2013). Novel applications of amylose-lipid complex as resistant starch type 5, in *Resistant Starch: Sources, Applications and Health Benefits* (eds. Y.-C. Shi & C.C. Maningat), Chichester: John Wiley & Sons. DOI: 10.1002/9781118528723.ch04
- Heß, C., Hartmann, B., Lechner, M.D., Nierling, W., Seidel, C., & Kulicke, W.M. (2007). Influence of soluble polymer residues in crosslinked carboxymethyl starch on some physical properties of its hydrogels. *Starch/Stärke*, 59, 425-429.
- Hettiaratchi, U.P.K., Ekanayake, S., & Welihinda, J. (2011). Nutritional assessment of a jackfruit (*Artocarpus heterophyllus*) meal. *Ceylon Medical Journal*, 56, 54-58.
- Jacobs, H., & Delcour, J.A. (1998). Hydrothermal modifications of granular starch, with retention of the granular structure: A review. *Journal of Agricultural and Food Chemistry*, 46, 2895-2905.
- Jagtap, U.B., & Bapat, V.A. (2010). *Artocarpus*: A review of its traditional uses, phytochemistry and pharmacology. *Journal of Ethnopharmacology*, 129, 142-166.



- Jagtap, U.B., Panaskar, S.N., & Bapat, V.A.. (2010). Evaluation of antioxidant capacity and phenol content in Jackfruit (*Artocarpus heterophyllus* Lam.) fruit pulp. *Plant Foods Hum Nutr*, 65, 99–104.
- Juliano, B.O. (1971). A simplified assay for milled-rice amylose, *Cereal Science Today*, 16, 334-360.
- Kittipongpatana, O.S., & Kittipongpatana, N. (2011). Preparation and physicochemical properties of modified jackfruit starches, *LWT – Food Science and Technology*, 44, 1766-1773.
- Kittipongpatana, N., Janta, S., & Kittipongpatana, O.S. (2011). Preparation of cross-linked carboxymethyl jackfruit starch and evaluation as a tablet disintegrant. *Pakistan Journal of Pharmaceutical Sciences*, 24, 415-420.
- Klein, B., Pinto, V.Z., Vanier, N.L., Zavareze, E.R., Colussi, R., Evangelho, J.A., Gutkoski, L.C., & Dias, A.R.G. (2013). Effect of single and dual heat-moisture treatments on properties of rice, cassava, and pinhão starches. *Carbohydrate Polymers*, 98, 1578-1584.
- Lai, X., He, C., Huang, Y., Chen, H., & Xiao, S. (2006). Starch extraction and its characteristics of jack-fruit seeds. *Shipin Yu Fajiao Gongye*, 32, 132-136.
- Li, S., Ward, R., & Gao, Q. (2011). Effect of heat-moisture treatment on the formation and physicochemical properties of resistant starch from mung bean (*Phaseolus radiatus*) starch. *Food Hydrocolloids*, 25, 1702-1709.
- Li, X., & Zhong, M. (2004). Study on granular morphology of starch isolated from jack-fruit seeds. *Shipin Kexue (Beijing, China)*, 25, 69-74.
- Lima, B.N.B., Lima, F.F., Tavares, M.I.B., Costa, A.M.M., & Pierucci, A.P.T.R. (2014). Determination of the centesimal composition and characterization of flours from fruit seeds. *Food Chemistry*, 151, 293-299.
- Liu, J., Ming, J., Li, W., & Zhao, G. (2012). Synthesis, characterisation and *in vitro* digestibility of carboxymethyl potato starch rapidly prepared with microwave-assistance. *Food Chemistry*, 133, 1196-1205.
- Madrigal-Aldana, D.L., Tovar-Gómez, B., Mata-Montes de Oca, M., Sáyago-Ayerdi, S.G., Gutierrez-Meraz, F., & Bello-Pérez, L.A. (2011). Isolation and characterization of Mexican



- jackfruit (*Artocarpus heterophyllus* L) seeds starch in two mature stages. *Starch/Starke*, 63, 364–372.
- Madrugá, M.S., de Albuquerque, F.S.M., Silva, I.R.A., do Amaral, D.S., Magnani, M., & Neto, V.Q. (2014). Chemical, morphological and functional properties of Brazilian jackfruit (*Artocarpus heterophyllus* L.) seeds starch. *Food Chemistry*, 143, 440–445.
- Mitra, A., Bhattacharya, D., & Roy, S. (2007). Role of resistant starches particularly rice containing resistant starches in Type II diabetes. *J. Hum. Ecol.*, 21, 47–51.
- Morais, M.B., Feste, A., Miller, R.G., & Lifchitz, C.H. (1996). Effect of resistant starch and digestible starch on intestinal absorption of calcium, iron and zinc in infant pigs. *Paediatric Research*, 39, 872–876.
- Mukherjee, B.D., Hossain, M.A., Haque, M.N., Ahad, M.M., Rahman, M., & Khan, N.A. (1960). Cereals and foods-starches in common food items and their properties. *Pakistan Journal of Biological and Agricultural Sciences*, 3(1), 41–45.
- Mukprasirt, A., & Sajjaanantakul, K. (2004). Physico-chemical properties of flour and starch from jackfruit seeds (*Artocarpus heterophyllus* Lam.) compared with modified starches. *International Journal of Food Science and Technology*, 39, 271–276.
- Narkhede, S.B., Bendale, A.R., Jadhav, A.G., Patel, K., & Vidyasagar, G. (2011). Isolation and evaluation of starch of *Artocarpus heterophyllus* as a tablet binder. *International Journal of PharmTech Research*, 3, 836–840.
- Nogami, H., Nagai, T., Fukuoka, E., Sonobe, T. (1969). Disintegration of the aspirin tablets containing starch and microcrystalline cellulose in various concentrations. *Chemical and Pharmaceutical Bulletins*, 17, 1450–1455.
- Oates, C.G., Powell, A.D. (1996). Bioavailability of carbohydrate material stored in tropical fruit seeds. *Food Chemistry*, 56, 405–414.
- Ocloo, F.C.K., Bansa, D., Boatın, R., Adom, T., & Agbemavor, W.S. (2010). Physico-chemical, functional and pasting characteristics of flour produced from Jackfruits (*Artocarpus heterophyllus*) seeds. *Agriculture and Biology Journal of North America*, 1, 903–908.



- Odoemelam, S.A. (2005). Functional properties of raw and heat processed jackfruit (*Artocarpus heterophyllus*) flour. *Pakistan Journal of Nutrition*, 4, 366-370.
- Ovando-Martínez, M., Whitney, K., Reuhs, B.L., Doehlert, D.C., & Simsek S. (2013). Effect of hydrothermal treatment on physicochemical and digestibility properties of oat starch. *Food Research International*, 52, 17-25.
- Pu, H., Chen, L., Li, X., Xie, F., Yu, L., & Li, L. (2011). An oral colon-targeting controlled release system based on resistant starch acetate: Synthetization, characterization, and preparation of film-coating pellets. *Journal of Agricultural and Food Chemistry*, 59, 5738-5745.
- Puncha-amon, S., & Uttapap, D. (2013). Rice starch vs. rice flour: Differences in their properties when modified by heat-moisture treatment. *Carbohydrate Polymers*, 91, 85-91.
- Rahman, M.A., Nahar, N., Mian, A.J., & Mosihuzzaman, M. (1999). Variation of carbohydrate composition of two forms of fruit from jack tree (*Artocarpus heterophyllus* L.) with maturity and climatic conditions. *Food Chemistry*, 65, 91-97.
- Rengsutthi, K., & Charoenrein, S. (2011). Physico-chemical properties of jackfruit seed starch (*Artocarpus heterophyllus*) and its application as a thickener and stabilizer in chilli sauce, *LWT - Food Science and Technology*, 44, 1309-1313.
- Roy, N., & Mitra, A.K. (1970). Amylose content in starches from some unfamiliar sources of food. *Journal of Food Science and Technology*, 7, 164.
- Santos, L.S., Bonomo, R.C.F., Fontan, R.C.I., Santos, W.O., & Silva, A.A.L. (2009). Gelatinization temperature and acid resistance of jackfruit seed starch. *CyTA – Journal of Food*, 7, 1–5.
- Sajilata, M.G., Singhal, R.S., & Kulkarni, P.R. (2006). Resistant starch - A review. *Comprehensive Reviews in Food Science and Food Safety*, 5, 1-17.
- Shih, F., King, J., Daigle, K., An, H.-J., & Ali, R. (2007). Physicochemical properties of rice starch modified by hydrothermal treatments. *Cereal Chemistry*, 84, 527-531.
- Sieb, P.A., Woo, K. (1999). Food grade starch resistant to alpha amylase and method of preparing the same. US patent 5,855,946.



- Sun, Q., Han, Z., Wang, L., & Xiong, L. (2014). Physicochemical differences between sorghum starch and sorghum flour modified by heat-moisture treatment. *Food Chemistry*, 145, 756-764.
- Sun, Q., Wang, T., Xiong, L., & Zhao, Y. (2013). The effect of heat moisture treatment on physicochemical properties of early indica rice. *Food Chemistry*, 141, 853-857.
- Swami, S.B., Thakor, N.J., Haldankar, P.M., & Kalse, S.B. (2012). Jackfruit and its many functional components as related to human health: A review. *Comprehensive reviews in Food Science and Food Safety*, 11, 565-576.
- Thompson, D.B. (2000). Strategies for the manufacturing of resistant starch. *Trends in Food Science & Technology*, 11, 245-253.
- Tongdang, T. (2008). Some properties of starch extracted from three Thai aromatic fruit seeds. *Starch/Starke*, 60, 199-207.
- Tulyathan, V., Tananuwig, K., Songjinda, P., & Jaiboon, N. (2002). Some physicochemical properties of jackfruit (*Artocarpus heterophyllus* Lam) seed flour and starch. *Science Asia*, 28, 37-41.
- Vatanasuchart, N., Niyomwit, B., & Wongkrajang, K. (2012). Resistant starch content, *in vitro* starch digestibility and physico-chemical properties of flour and starch from Thai bananas. *Maejo International Journal of Science & Technology*, 6(2), 259-271.
- Zavareze, E.R., El Halal, S.L.M., Santos, D.G., Helbig, E., Pereira, J.M., & Guerra Dias, A.R. (2012). Resistant starch and thermal, morphological and textural properties of heat-moisture treated rice starches with high-, medium- and low-amylose content. *Starch/Starke*, 64, 45-54.
- Zhang, H., & Jin, Z. (2011). Preparation of products rich in resistant starch from maize starch by an enzymatic method. *Carbohydrate Polymers*, 86, 1610-1614.