Isolation of Endophytic Fungi from some Orchid Varieties

ชื่อโครงการ Isolation of Endophytic Fungi from some Orchid Varieties

แหล่งเงิน เงินงบประมาณ คณะเทคโนโลยีการเกษตร สถาบันเทคโนโลยีพระจอมเกล้าเจ้าคุณทหาร ลาดกระบัง

ประจำปังบประมาณ 2557 จำนวนเงินที่ได้รับการสนับสนุน 100,000 บาท ระยะเวลาทำการวิจัย 1 ปี ตั้งแต่ 1 ตุลาคม ถึง 30 กันยายน 2555 ชื่อ-สกุล หัวหน้าโครงการ ผศ.คร.ศรายุทธ ผลโพธิ์

ผู้ร่วมโครงการวิจัย นายวรรณะ สัว นักศึกษาระดับปริญญาโท สาขาเกษตรศาสตร์

Abstract

First study on endophytic fungi from orchids in Chantaburi province, Thailand. Endophytic fungi were isolated from 9 orchid varieties in green house condition. Leaves and roots of Grammatophyllum scriptum, Cymbidium dayanum, Dendrobium hercoglossum, Dendrobiumpalpebrae, Toreniafournieri, Doritispulcherrima, Dendrobium crumenatum, Dendrobium friedericksianum and Grammatophyllum specinocum. All samples were collected then brought to laboratory. Isolation was done by surface sterilization, removed the outer surface and got inner tissues then place on water agar and transferred culture to potato dextrose agar to be pure culture. The morphological identification based on colony and agar color, liquid drops, colony shapes, the growth characteristic Twelve speciesare identified as Chaetomium cochliodes, rate Chaetomiumcupreum, Xylaria sp., Colletotrichum sp., Nigrospora sp., Phoma sp. Curvularia sp., Fusarium sp., Pestalotiopsis sp., Corynascus sp. and Cladosporium-like fungi, were obtained. Most of fungi that isolated were familiar to endophytic that isolated from many plant as previous study. All of the fungi have different benefit up to using.

Keywords: Endophytic fungi, Orchids

Isolation of Endophytic Fungi from some Orchid Varieties

Acknowledgement

This research has got scholarship from King Mungkut's Instutute of Technology Ladkrabang. I would like to express my sincere to thank to Faculty of Agriculture Technology, King Mongkut's Institute of Technology Ladkrabang (KMITL), Bangkok, Thailand. I also thank to the teacher and stuff of Rajamangala University Technology Tawan-ok Chantaburi campus what allowed to collect orchid samples in this research.

> ผศ.คร.ศรายุทธ ผลโพธิ์ นายวรรณะ สัว

รายงานวิจัยฉบับสมบูรณ์ | iii Isolation of Endophytic Fungi from some Orchid Varieties |

Table of Content

Content	Page
Abstract	i
Acknowledgments	ii
Table of content	iii
List of figures	iv
Chapter 1 Introduction	1
Chapter 2 Review literature	3
Chapter 3 Materials and Methods	11
Chapter 4 Result and Discussion	13
Reference	20
Appendix	22

รายงานวิจัยฉบับสมบูรณ์ | iv Isolation of Endophytic Fungi from some Orchid Varieties |

List of figures

Figures	Page
Fig. 1: Chaetomium cochliodes A: The colony that culture on PDA 30 days, BThe spores	13
under microscops 10XCThe ascospores under microscops 40X	
Fig. 2: Chaetomium cupreum A: The colony cultured on PDA 30 daysB: The spores	13
under microscops 10X and C:Theascospores under microscops 40X	
Fig. 3:Xylaria sp.A: Colony culture on PDA 30 days B: Mycelia of xylaria sp. under	14
microscops 40X	
Fig. 4: Colletotrichum sp.A: The colony culture on PDA 30 days B: The spores under	14
microscope 40X	
Fig. 5: Nigrospora sp. A: The colony culture on PDA 30 daysB: The hyphea and spores	15
under microscope 40X	
Fig. 6:Phoma sp1. A: The colony culture on PDA 30 days and B: The spores and	15
mycelia under microscope 40X	
Fig. 7: Curvularia sp. A: The colony culture on PDA 30 days and B: The spores under	16
microscope 40X	
Fig. 9:Pestalotiopsis sp. A: The colony culture on PDA 30 days B:The conidia under	16
microscops 40X	
Fig. 10: Corynascus sp. A: The colony culture on PDA 30 days and B: The spores and	16
acospores under microscops40X	
Fig. 11: Cladosporium-like fungiA: The colony culture on PDA 30 days	17
and B: Thespores and mycelia under microscops 40X	
Fig. 12:Achaetomium sp.A: The colony culture on PDA 30 days B: The spores under	27
microscops 10X and C: The ascospores under microscops 40X	