

LIST OF TABLES

TABLE	PAGE
2.1 A basic drug list for cancer pain relief	16
2.2 Studies reporting the cancer pain.....	29
2.3 Cases and ASR of leading type of cancer in 1996	41
2.4 Availability of opioids analgesics in Thailand (update Dec. 2008)	41
2.5 The annually quantity of opioids notified in the Ministerial Notification of Ministry of Public Health No.197 B.E.2549 (2006)	42
2.6 The annually quantity of opioids notified in the Ministerial Notification of Ministry of Public Health No.121 B.E.2536 (1993)	42
3.1 Clinical setting and demographic factors	57
3.2 Distribution of primary cancer sites by gender	58
3.3 Clinical setting and oncological factors.....	58
3.4 The percentage of pain control, depression and quality of life score by clinical setting	59
3.5 Comparison of BQ-II between original version and modified version using in the main study with its reasons.....	61
3.6 Background characteristics of physicians by setting	69
3.7 Physicians' attitudes toward the optimal use of analgesic for cancer pain management.....	70
3.8 The percentage of physicians' knowledge of opioid prescribing.....	71
3.9 The percentage of physicians' attitudes toward opioid prescribing.....	72
3.10 Identified barriers to cancer pain management by setting.....	73
4.1 Distribution of the study sample by demographic characteristics.....	83
4.2 Distribution of primary cancer sites	86
4.3 Distribution of primary cancer sites by gender	87
4.4 Cancer characteristics and treatment	88
4.5 Cancer pain characteristics	89
4.6 Mean, standard deviation, median and range score on the pain profile (n=228)	92

TABLE	PAGE
4.7 Patients' evaluation of their pain profile on different severity (n=228).....	92
4.8 Patients' evaluation of their pain profile by gender	92
4.9 Patients' evaluation of pain impact on every living activity and symptom by clinical setting	93
4.10 Pain Management Index (PMI) by clinical setting.....	93
4.11 PMI and corresponding pain severity	94
4.12 Prevalence of inadequate pain control, inappropriate analgesic prescription and depression by clinical setting	94
4.13 Mean scores of global quality of life and subscales by cancer staging.....	94
4.14 Mean scores of global quality of life and subscales by clinical setting.....	95
4.15 The patients' characteristics between the clinical settings	99
4.16 Distribution of primary cancer sites by clinical setting.....	100
4.17 Patients' pain characteristics by clinical setting.....	100
4.18 Cancer treatment within the previous month by clinical setting	101
4.19 Pain medications by clinical setting	102
4.20 Average dosage of opioid medication in patients with inadequate pain control by clinical setting	103
4.21 The equianalgesic oral morphine dosage (mg/day) by clinical setting	103
4.22 Co-analgesic usage in patients with neuropathic pain by clinical setting ...	104
4.23 Number of analgesic items prescribed by clinical setting	105
4.24 Mean scores subscale of Modified BQ-II by clinical setting	105
4.25 The association between clinical setting and patients' characteristics.....	108
4.26 The association between clinical setting and analgesic prescription.....	108
4.27 ORs for the association between inadequate pain control, clinical setting and potential confounders.....	109
4.28 The Odds ratio following logistic regression for the association between clinical setting and inadequate pain control, controlling for potential confounders.....	112

TABLE	PAGE
4.29 Final ‘best fit’ multivariate model (logistic regression), based on clinical setting and other variables with pain control as the dependent variable	114
4.30 The association between depression and potential confounders	116
4.31 Final ‘best fit’ multivariate model (logistic regression), based on clinical setting and other variables with depression as the dependent variable.....	117
4.32 The association between quality of life and potential confounders.....	119
4.33 Final ‘best fit’ multivariate model (logistic regression), based on clinical setting and other variables with poor QOL as the dependent variable	120