

References

Book

- Chan, P.K. & Hayes, A.W. (1989). Principles and methods for acute toxicity and eye irritancy. In Principles and methods of toxicology. 2nd ed. New York Raven Press.
- Juliano, B.O., & Bechtel, D.B., (1985). The rice grain and its gross composition. In: Juliano, B.O. (Eds.), Rice Chemists and Technology, second ed. The American Association of Cereal Chemists, St Paul, MI, USA, pp. 17–57.
- Nicolosi, R.J., Rogers, E.J., Ausman, LM., & Orthoefer, F.T. (1993). Rice bran oil and its health benefits. In Rice science and Technology. Marshall WE, Wadsworth JI., (Eds.), Dekker: New York, p421.
- Rukmini, C. (2000). Bioactive in rice bran and rice bran oil, In: Bidlack,W.R., Omaye, S.T., Meskin,M.S., Topham, D.K.W. (Eds), Phytochemicals as Bioactive Agents, Technomic Publishing Company, Inc.851 New Hooland Avenue, Box3535, Lancaster, Pennsylvania, pp.213-240.
- Shuman, C.R, (1988). Diabetes mellitus: definition, classification, and diagnosis. In J.A.Galloway, J.H. Potvin,& C.R. Shuman (Eds.), Diabetes mellitus (9th ed.) (pp.2-13). Indianapolic,Indiana: Eli Lilly and Company Lilly Corporate Center.
- Williams, G., & Pickup, JC. (1999). Hand book of diabetes mellitus. Second edition. Cambridge, Great Britain at the University Press.

Articles

- Abbasi, F., Carantoni, M., Chen, Y.D., & Reaven G.M. Further evidence for a central role of adipose tissue in the antihyperglycemic effect of metformin. *Diabetes Care*, 21, 1301-1305, 1998.
- Adlerberth, A.M., Rosengren, A., & Wilhelmsen, L. (1998). Diabetes and long-term risk of mortality from coronary and other causes in middle-aged Swedish men. *Diabetes Care*, 21, 539-545.
- Aekplakorn, W., Stolk, R.P., Neal, B., Suriyawongpaisal, P., Chongsuvivatwong, V., Cheepudomwit, S., et al. (2003). The prevalence and management of diabetes in Thai adults. *Diabetes Care*, 26, 2758-63.
- Ardiansyah, Shirakawa, H., Koseki, T., Ohinata, K., Hashizume, K., & Komai, M. (2006). Rice bran fractions improve blood pressure, lipid profile, and glucose metabolism in stroke-prone spontaneously hypertensive rats. *Journal of agricultural and food chemistry*, 54(5), 1914-1920.
- Ardiansyah, Shirakawa, H., Koseki, T., Hashizume, K., & Komai, M. (2007). The Driselase-treated fraction of rice bran is a more effective dietary factor to improve hypertension, glucose and lipid metabolism in stroke-prone spontaneously hypertensive rats compared to ferulic acid. *British Journal of Nutrition*, 97, 67-76.
- American Diabetes Association.(2002). Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care*, 25, S5-S20.
- American Diabetes Association. (2004). Screening for type 2 diabetes. *Diabetes Care*, 27 (Suppl.1), S11– S14.
- Behall, K.M., Scholfield, D.J., & Hallfrisch, J. (2004). Diet containing barley significantly reduce lipids in mildly hypercholesterolemic men and women. *The American Journal of Clinical Nutrition*, 106(9), 1185-1193.
- Bergman, R.N., Kim, S.P., Katalano, K.J., Hsu, I.S., Chiu, J.D., Kabir, M., et al. (2006). Why visceral fat is bad: mechanisms of the metabolic syndrome. *Obesity*, 14 (Suppl.1), 16S-19S.

- Bowman, S.A., & Vinyard, B.T. (2004). Fast Food Consumption of U.S. Adults: Impact on Energy and Nutrient Intakes and Overweight Status. *Journal of the American College of Nutrition*, 23(2), 163–168.
- Bray, G.A., & Popkin, B.M. (1998). Dietary fat intake does affect obesity!. *American Journal of Clinical Nutrition*, 68, 1157–73.
- Buettner, B., Parhofer, K.G., Woenckhaus, M., Wrede, C.E., Kunz-Schughart, L.A., Schölmerich, J. et al. (2006). Defining high-fat-diet rat models: metabolic and molecular effects of different fat types. *Journal of Molecular Endocrinology*, 36, 485–501.
- Chotimarkorn, C., Benjakul, S., & Silalai, N. (2008). Antioxidant components and properties of five long-grained rice bran extracts from commercial available cultivars in Thailand. *Food Chemistry*, 111, 636–641.
- Claret, M., Corominola, H., Canals, I., Nadal, B., Chavanieu, A., Pfeiffer, B., et al. (2004). S 23521 decreases food intake and body weight gain in diet-induced obese rats. *Obesity Research*, 12, 1596-603.
- Cicero, AFG., & Derosa, G. (2005). Rice bran and its main components: Potential role in the management of coronary risk factors. *Current Topics in Nutraceutical Research*, 3(1), 29-46.
- Eitenmiller, R.R. (1997). Vitamin E Content of Fats and Oils: Nutritional Implications. *Food Technology*, 51(5), 78-81.
- Freemark, M. & Bursey, D. (2001). The effects of metformin on body mass index and glucose tolerance in obese adolescents with fasting hyperinsulinemia and a family history of type 2 diabetes. *Pediatrics*, 107, E55.
- Fuh, W-S., & Chiang, B-H. (2001). Dephytinisation of rice bran and manufacturing a new food ingredient. *Journal of the Science of Food and Agriculture*, 81, 1419-25.
- Gabir, M.M., Hanson, R.L., Dabelea, D., Imperatore, G., Roumain, J., Bennett, PH., et al. (2000). The 1997 American Diabetes Association and 1999 World Health Organization Criteria for Hyperglycemia in the Diagnosis and Prediction of Diabetes. *Diabetes Care*, 23, 1108-1112.
- Gerhardt, A.L., & Gallo, N.B. (1998). Full-fat rice bran and oat bran similarly reduce hypercholesterolemia in humans. *The journal of Nutrition*, 128, 865-9.

- Golay, A., & Bobbioni, E. (1997). The role of dietary fat in obesity. *International Journal of Obesity*, 21 (Suppl.3), S2-11.
- Grundy, S.M., Brewer, H.B. Jr, Cleeman, Jr, Smith, S.C. Jr, Lenfant, C. and for the Conference Participants.(2004). Definition of Metabolic Syndrome: Report of the National Heart, Lung, and Blood Institute/American Heart Association Conference on Science Issues Related to Definition. *Circulation*, 109:433-8.
- Hallfrisch, J., FACN., & Behall, KM. (2000). Mechanisms of the effects of grains on insulin and glucose responses. *Journal of the American College of Nutrition*, 19(3), 320S–325S.
- Hiramatsu, S., & Grill, V. (2001). Influence of a high-fat diet during chronic hyperglycemia on β -cell function in pancreatic islet transplants to streptozotocin-diabetic rats. *European Journal Endocrinology*, 144:521–527
- Hu, F.B., Dam, R.M.van, & Liu, S. (2001) Diet and the risk of Type-II diabetes. *Diabetologia*, 4, 805–817.
- Hu, F.B., Rimm, E.B., Stampfer, M.J., Ascherio, A., Spiegelman, D., & Willett, W.C. (2000). Prospective study of major dietary patterns and risk of coronary heart disease in men. *The American Journal of Clinical Nutrition*, 72, 912–21.
- Hu, W., Wels, JH., Shin, TS., & Godber, JS. (1996). Comparison of isopropanal and hexane for extraction of vitamin E and oryzanols from stabilized rice bran. *Journal of the American Oil chemists Society*, 73, 1653-1656.
- Isganaitis, E., & Lustig, RH. (2005). Fast Food, Central Nervous System Insulin Resistance, and Obesity. *Arteriosclerosis, Thrombosis, Vascular Biology*, 25, 2451-62.
- Juliano, C., Cossu, M., Alamanni, MC., & Piu,L. (2005). Antioxidant activity of gamma-oryzanol: Mechanism of action and its effect on oxidative stability of pharmaceutical oils. *International Journal of Pharmaceutics*, 299, 146-154.

- Kahlon, T.S., Chow, F.I., Sayre, R.N., & Betschart, A. (1992). Cholesterol-lowering in hamsters fed rice bran at various level, defatted rice bran and rice bran oil. *The journal of nutrition*, 122, 513-519.
- Kahlon, T.S., Saunders R.M., Sayre R.N., Choe, F.I., Chiu, M.M., & Betschart, AA. (1992). Cholesterol-lowering effect of rice bran and rice bran oil fractions in hypercholesterolemia hamsters. *Cereal Chemistry*, 69(5): 485-489.
- Kahlon, T.S., Chow, F.I., Chiu, M.M., Hudson, CA., & Sayre, R.N. (1996). Cholesterol-lowering by rice bran and rice bran oil unsaponifiable matter in hamsters. . *Cereal Chemistry*, 73, 69-74.
- Kanaya, Y., Doi, T., Sasaki, H., Fujita, A., Matsuno, S., Okamoto, K., et al. (2004). Rice bran extract prevents the elevation of plasma peroxylipid in KKAY diabetic mice. *Diabetes research and Clinical Practice*, 66S, S157-S160.
- Katz, A., Nambi, SS., Mather, K., Baron, AD., Follmann, DA., Sullivan, G., et al. (2000). Quantitative insulin sensitivity check index: a simple, accurate method for assessing insulin sensitivity in humans. *The Journal of Clinical Endocrinology & Metabolism*, 85, 2402-2410.
- Kay, J.P., Alemzadeh, R., Langley, G., D'Angelo, L., & Holshouser S. (2001). Beneficial effects of metformin in normoglycemic morbidly obese adolescents. *Metabolism*, 50 (12), 1457-1461.
- Kennedy, G., & Burlingame, B. (2003). Analysis of food composition data on rice from a plant genetic resources perspective. *Food Chemistry*, 80, 589–596.
- Kestin, M., Moss, R., Clifton, P.M., & Nestel, P.J. (1990). Comparative effects of three cereal brans on plasma lipids, blood pressure, and glucose metabolism in mildly hypercholesterolemic men. *The American Journal of Clinical Nutrition*, 52, 661-6.
- Kharroubi, I., Ladriere, L., Cardozo, A. K., Dogusan, Z., Cnop, M., & Eizirik, D. L. (2004). Free fatty acids and cytokines induce pancreatic beta-cell apoptosis by different mechanisms: role of nuclear factor-kappaB and endoplasmic reticulum stress. *Endocrinology*, 145(11), 5087-5096.

- King, H., Aubert, R.E., & Herman, W.H. (1998). Global burden of diabetes 1995-2025; Prevalence, numerical estimates, and projections. *Diabetes care*, 21, 1414-1431.
- Lloyd, B.J., Siebenmorgen, T.J., & Beers, K.W. (2000). Effects of commercial processing on antioxidant in rice bran. *Cereal Chemistry*, 77, 551-555.
- Lichtenstein, A.H., Ausman, L.M., Carrasco, W., Gualtieri, L.J., Jenner, J.L., Ordovas, J.M., et al. (1994). Rice bran oil consumption and plasma lipid levels in moderately hypercholesterolemic human. *Arteriosclerosis, Thrombosis, Vascular Biology*, 14, 549-556.
- Lichtenstein, A.H., & Schwab, U.S. (2000). Relationship of dietary fat to glucose metabolism. *Atherosclerosis*, 150, 227-243.
- Lorenzo, C., Williams, K., Hunt, K.J., & Haffner, S.M. (2007). The National Cholesterol Education Program—Adult Treatment Panel III, International Diabetes Federation, and World Health Organization Definitions of the Metabolic Syndrome as Predictors of Incident Cardiovascular Disease and Diabetes. *Diabetes Care*, 30:8–13.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L., & Randall, R. J. (1951). Protein measurement with the Folinphenol reagents. *Journal of Biological Chemistry*, 193, 265–275 (1951).
- Matthews, D.R., Hosker, P.J., Rudenski, A.S., Naylor, B.A., Treacher, D.F., & Turner, R.C. (1985). Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*, 28, 412-419.
- Minhajuddin, M., Beg, Z.H., & Iqbal, J. (2005). Hypolipidemic and antioxidant properties of tocotrienol rich fraction isolated from rice bran oil in experimentally. *Food and Chemical Toxicology*, 43, 747-753.
- Mooradian, A.D. (2003). Cardiovascular disease in type 2 diabetes mellitus: Current management Guideline. *Archives Internal Medicine*, 163, 33-40.
- Most, M.M., Tulley, R., Morales, S., & Lefevre, M. (2005). Rice bran oil, not fiber, cholesterol in humans. *The American Journal of Clinical Nutrition*, 81, 64-8.

- Newman R.K., Betschart A.A., Newman C.W., & Hofer, P.J. (1992). Effect of full-fat or defatted rice bran on serum cholesterol. *Plant Foods for Human Nutrition*, 42: 37-43.
- Nicolosi, R.J., Ausman, L.M., & Hegsted, D.M. (1991). Rice bran oil lowers serum total and LDL lipoprotein cholesterol and apo B levels in non human primates. *Atherosclerosis*, 88, 133-142.
- Nicolosi, R.J., Ausman, L.M., & Hegsted, D.M. (1989). Lipoprotein level in monkeys fed a diet containing rice oil. *Circulation* 80, A86 (Abstract)
- Nishida, C., Uauy, R., Kumanyika, S., & Shetty, P. (2004). The joint WHO/FAO expert consultation on diet, nutrition and the prevention of chronic diseases: process, product and policy implications. *Public Health Nutrition*, 7(1A), 245-250.
- Pagliassotti M.J., Gayles, E.C., Podolin, D.A., Wei,Y., & Morin, C.L. (2000). Developmental stage modifies diet-induce peripheral insulin resistance in rats. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 278, 66-73.
- Prakash, J. (1996). Rice bran protein: properties and food uses. *Critical Review in Food Science and Nutrition*, 36(6), 537-552.
- Rocchini, A.P., Marker, P., & Cervenka, T. (1997). Time course of insulin resistance associated with feeding dogs a high-fat diet. *American Journal of Physiology*, 272, E147-E154.
- Qureshi, A.A., Sami, S.A., & Khan, F.A.(2002). Effects of stabilized rice bran, its soluble and fiber fractions on blood glucose levels and serum lipid parameters in humans with diabetes mellitus Types I and II. *The Journal of Nutritional Biochemistry*, 13, 175-187.
- Qureshi, A.A., Sami, S.A., Salser, W.A, & Khan, F.A. (2002). Dose-dependent suppression of serum cholesterol by tocotrienol-rich fraction (TRF25) of rice bran in hypercholesterolemic humans. *Atherosclerosis*, 161,199-207.
- Rawdaree, P., Ngarmukos, C., Deerochanawong, C., Suwanwalaikorn ,S., Chetthakul, T., Krittiyawong, S., et al. (2006). Thailand Diabetes Registry (TDR) Project. Clinical status and long term vascular complications in diabetic patients. *Journal Medicine Associate Thailand*, 89, S1-9.

- Reagan-Shaw, S., Nihal, M., & Ahmad, N. (2007). Dose translation from animal to human studies revisited. *The Journal of the Federation of American Societies for Experimental Biology*, 22, 659-661.
- Richmond, W. (1992). Analytical reviews in clinical biochemistry: the quantitative analysis of cholesterol. *Annals of Clinical Biochemistry*. 577–597.
- Rimm, EB., Ascherio, A., Giovannucci, E., Spiegelman, D., Stampfer, MJ., & Willett WC. (1996). Vegetable, fruit, and cereal fiber intake and risk of coronary heart disease among men. *JAMA*, 275, 447–51.
- Rocchini, A.P., Marker, P., & Cervenka, T. (1997). Time course of insulin resistance associated with feeding dogs a high-fat diet. *American Journal of Physiology*, 272, E147-E154.
- Rukmini, C., & Raghuram, T.C. (1991). Nutritional and biochemical aspects of the hypolipidemic action of rice bran oil: A review. *Journal of the American College Nutrition*, 10(6), 593-601.
- Sharma, R.D., & Rukmini, C. (1986). Rice bran oil and hypocholesterolemia in rat. *Lipids*, 21, 715-717.
- Shin, T.G., Godber, J.S., Martin, D.E., & Wells, J.H. (1997). Hydrolytic stability and changes in E vitamins and oryzanol of extruded rice bran during storage. *Journal of food science*, 62, 704-708.
- Srinivasan, S., Ambler, G.R., Baur, L.A., Garnett, S.P., Tepsa, M., Yap, F., et al. (2006). Randomized, controlled trial of metformin for obesity and insulin resistance in children and adolescents: Improvement in body composition and fasting insulin. *The Journal of Clinical Endocrinology & Metabolism*, 91(6), 2074–2080.
- Storlien, L.H., Oakes, N.D., Pan, D.A., Kusunoki, M., & Jenkins, AB., (1993). Syndrome of insulin resistance in the rat: inducement by diet and amelioration with benfluorex. *Diabetes*, 42: 457-462.
- Storlein, L.H., Baur, L.A., Kriketos, A.D., Pan, D.A., Cooney, G.J., Jenkins, A.B., et al. (1996) Dietary fats and insulin action. *Diabetologia*, 39:621–631.
- Sugano, M., & Tsuji, E. (1997). Rice bran oil and cholesterol metabolism. *The Journal of nutrition*, 127, 521S-524S.

- Tripathy, D., Carlsson, M., Almgren, P., Isomaa, B., Taskinen, M.R., Tuomi, T., et al. (2000). Insulin secretion and insulin sensitivity in relation to glucose tolerance. Lessons from the Botnia Study. *Diabetes*, 49, 975-80.
- Wang, M., Hettiarachchy, N.S., Qi, M., Burks, W., & Siebenmorgen, T. (1999). Preparation and Functional Properties of Rice Bran Protein Isolate. *Journal of agricultural and food chemistry*. 1999, 47, 411-416
- Weickert, M. O., Mohlig, M., Schofl, C., Arafat, A. M., Otto, B., Viehoff, H., et al. (2006). Cereal fiber improves whole-body insulin sensitivity in overweight and obese women. *Diabetes Care*, 29(4), 775-780.
- Weststrate, J.A., & Meijer, G.W. (1998). Plant sterol enriched margarines and reduction of plasma total, LDL-cholesterol concentrations in normocholesterolemic and mildly hypercholesterolemic subjects. *European Journal of Clinical Nutrition*, 52(5), 334-343.
- WHO/IDF, World Health Organization / International Diabetes Federation. (2006). Definition and diagnosis of diabetes mellitus and intermediate hyperglycemia: report of a WHO/IDF consultation. Geneva, Switzerland: World Health Organization.
- Wilson, T.A., Nicolosi, R.J., Woolfrey, B., & Kritchevsky, D. (2007). Rice bran oil and oryzanol reduce plasma lipid and lipoprotein cholesterol concentrations and aortic cholesterol ester accumulation to a greater extent than ferulic acid in hypercholesterolemic hamsters. *Journal of Nutritional Biochemistry*, 18, 105-112.
- Wild, S., Roglic, G., Green, A., Sicree, R., King, H. (2004). Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care*, 27, 1047-53.
- Wilkes, J.J., Bonen, A., & Bell, R.C. (1998). A modified high-fat diet induce insulin resistance in rat skeletal muscle but not adipocytes. *American Journal Physiology*, 275, E679-686.
- Wolever TMS, Jenkins DJA, Jenkins AL, Josse RG. (1991). The glycemic index: Methodology and clinical implications. *The American Journal of Clinical Nutrition*, 54, 846-854.

- World Health Organization. (1999). Definition, Diagnosis and Classification of Diabetes Mellitus and Its Complications: Report of a WHO Consultation. Geneva: World Health Organization.
- Xu, Z., & Godber, JS. (1999). Purification and identification of component of γ -oryzanol in rice bran oil. *Journal of Agricultural and Food Chemistry*, 47, 2724-2728.
- Yu, L., Haley, S., Perret, J., Harris, M., Wilson, J., & Qian, M. (2002). Free Radical Scavenging Properties of Wheat Extracts. *Journal of Agricultural and Food Chemistry*, 50, 1619–1624.
- Zhou, Y-P., & Grill, V., (1995). Long term exposure to fatty acids and ketones inhibits B-cell functions in human pancreatic islets of langerhans. *Journal of Clinical Endocrinology and Metabolism*, 80, 1584-1590.
- Zuberi A.(2008). Strategies for assessment of botanical action on metabolic syndrome in the mouse and evidence for a genotype-specific effect of Russian tarragon in the regulation of insulin sensitivity. *Metabolism Clinical and Experimental*, 57(1), S10-S15.