

LIST OF FIGURES

FIGURE	PAGE
2.1 <i>Piper chaba</i> Linn. or <i>Piper longum</i> Linn. (Piperaceae)	9
2.2 Fruit of <i>Piper chaba</i> Linn. (Piperaceae).....	9
2.3 <i>Piper sarmentosum</i> Roxb.(Piperaceae)	10
2.4 Fruit of <i>Piper sarmentosum</i> Roxb. (Piperaceae).....	10
2.5 <i>Piper interruptum</i> Opiz. (Piperaceae)	11
2.6 Fruit of <i>Piper interruptum</i> Opiz. (Piperaceae).....	12
2.7 <i>Plumbago indica</i> Linn. (Plumbaginaceae)	13
2.8 Flower of <i>Plumbago indica</i> Linn. (Plumbaginaceae)	13
2.9 The upper stem of <i>Zingiber officinale</i> Roscoe (Zingiberaceae)	15
2.10 The Rhizome of <i>Zingiber officinale</i> Roscoe (Zingiberaceae).....	15
2.11 Structures of some chemical constituents found in <i>P. chaba</i>	20
2.12 Structures of some chemical constituents found in <i>P. sarmentosum</i>	24
2.13 Structures of some chemical constituents found in <i>P. interruptum</i> ..	27
2.14 Structures of some chemical constituents found in <i>P. indica</i>	28
2.15 Structures of some chemical constituents found in <i>Z. officinale</i>	35
3.1 Dried fruit of <i>Piper longum</i> Linn.	65
3.2 Dried root of <i>Piper sarmentosum</i> Roxb.	65
3.3 Dried stem of <i>Piper interruptum</i> Opiz.	65
3.4 Dried root of <i>Plumbago indica</i> Linn.	66
3.5 Dried rhizome of <i>Zingiber officinale</i> Roscoe.	66
3.6 Bioassay-guided fractionation of the ethanolic extract of Benjakul preparation.....	71
4.1 Cytotoxic activity [IC ₅₀ (µg/ml)] of five plants and Benjakul preparation on five types of cell lines exposure time 72 hr (n=3) using student t-test from prism to compare the significant difference between normal cell (MRC-5) and each cancer cell (Hela, HepG2, MCF-7 and COR-L23) (* p<0.05, **p<0.0001).	79
4.2 Histogram comparing IC ₅₀ of each fraction against COR-L23 at exposure time 72 h.....	81

LIST OF FIGURES (continued)

FIGURE	PAGE
4.3 Structure of piperine	82
4.4 ¹ H NMR spectrum of BENS1 in CDCl ₃ and CD ₃ OD	84
4.5 EIMS spectrum of BENS1.....	85
4.6 Structure of plumbagin	86
4.7 ¹ H NMR spectrum of BENS2 in CDCl ₃ and CD ₃ OD	87
4.8 EIMS spectrum of BENS2.....	88
4.9 Structure of 6-gingerol.....	89
4.10 ¹ H NMR spectrum of BENS3 in CDCl ₃	91
4.11 EIMS spectrum of BENS3.....	92
4.12 The chemical structure of three compounds isolated from the ethanolic extract of Benjakul preparation.	93
4.13 Histogram comparing IC ₅₀ values of isolated compounds of Benjakul preparation against four cancer cell lines (Hela, HepG2, MCF-7 and COR-L23) and one normal cell lines (MRC-5) at exposure time 72 hrs (n=3).....	96
4.14 HPLC chromatogram of ethanolic extract of Benjakul preparation (10 mg/ml). Mobile phase: water-acetonitrile with gradient elution as follows: 0 min, 60:40; 30 min, 50:50; 50 min, 5:95; 60 min, 0:100; Flow rate 1.0 ml/min; UV detector at 256 nm.	98
4.15 The specificity validation for the HPLC analytical method for piperine and plumbagin: (A) blank solution; (B) piperine and plumbagin standards solution and (C) ethanolic extract of Benjakul preparation sample solution.....	100
4.16 Calibration curve of standard piperine, concentrations range from 50-400 µg/ml.	101
4.17 Calibration curve of standard plumbagin, concentrations range from 10-200 µg/ml.	102

LIST OF FIGURES (continued)

FIGURE		PAGE
4.18	The stability of piperine and plumbagin (% content) in the ethanolic extract of Benjakul preparation under accelerated condition ($45 \pm 2^{\circ}\text{C}$ with $75 \pm 5\%$ RH).....	106