

List of Figures

Figure		Page
2.1	Plants called Hua-Khao-Yen	4-5
2.2	<i>Dioscorea membranacea</i> Pierre A and B inflorescence	7
	C flower D stem	
2.3	Chemical structures of isolated compounds from <i>D. membranacea</i> ...	9
2.4	Loss of normal cell growth control	14
2.5	The effects of oncogenes and tumorsuppressor genes	17
	on cellular function	
2.6	The breast	24
2.7	MCF-7 human breast cancer cell line	27
2.8	Sulphorhodamine B	31
3.1	Conceptual framework	42
3.2	Rhizome of <i>D. membranacea</i> in dried powder	43
3.3	Extraction of <i>D. membranacea</i>	45
3.4	Isolation of dioscorealide B from <i>D. membranacea</i> by column	47
	chromatography	
4.1	Starch grains of <i>D. membranacea</i> rhizome under microscope	57
4.2	Calcium oxalate crystals of <i>D. membranacea</i> rhizome	58
	under microscope	
4.3	TLC chromatograms of isolated compound and dioscorealide B	62
	standard	
4.4	Chemical structure of dioscorealide B	62
4.5	¹ HNMR spectrum of isolated dioscorealide B	64
4.6	HPLC chromatogram of dioscorealide B	67
4.7	Calibration curve of dioscorealide B	68
4.8	HPLC fingerprints of <i>D. membranacea</i> extracts from different	71-73
	extraction methods	
4.9	Total content of dioscorealide B (µg/g dried plant) in crude extracts ...	77
	of <i>D. membranacea</i> from different extraction methods	

List of Figures (continued)

Figure		Page
4.10	Concentration of dioscorealide B in crude extracts at 10 mg/ml and IC ₅₀ of cytotoxicity against MCF-7 by SRB assay of crude extracts from different extraction methods	80
4.11	Percentage of dioscorealide B remaining in the ethanolic extracts of <i>D. membranacea</i> in stability test under accelerated conditions at 45°C and 75% RH for 4 months	82
4.12	Percentage of dioscorealide B remaining in the ethanolic extracts of <i>D. membranacea</i> in stability test under accelerated conditions at 45°C, 60°C, 70°C and 80°C with 75% for 1 month	84
4.13	Stability test of the ethanolic extract of <i>D. membranacea</i> under accelerated condition at 45°C, 75% RH for 4 months exhibited as %dioscorealide B remaining and %inhibition of cytotoxic activity against MCF-7 by SRB assay at 10 µg/ml	87
4.14	Stability test of the ethanolic extract of <i>D. membranacea</i> under accelerated condition at 60°C, 75% RH for 1 month exhibited as %dioscorealide B remaining and %inhibition of cytotoxic activity against MCF-7 by SRB assay at 10 µg/ml	87
4.15	Stability test of the ethanolic extract of <i>D. membranacea</i> under accelerated condition at 70°C, 75% RH for 1 month exhibited as %dioscorealide B remaining and %inhibition of cytotoxic activity against MCF-7 by SRB assay at 10 µg/ml	88
4.16	Stability test of the ethanolic extract of <i>D. membranacea</i> under accelerated condition at 80°C, 75% RH for 1 month exhibited as %dioscorealide B remaining and %inhibition of cytotoxic activity against MCF-7 by SRB assay at 10 µg/ml	88