

รายการอ้างอิง

หนังสือ

- Abraham, G. & Hron, R.J. (1992). *Oilseeds and Their Oils*. New York: John Wiley & Sons, Inc.
- Apiratkamon, S.(2004). *Principles of Statistics*. Bangkok: University of Thai Chamber of Commerce Publishers.
- Gunstone, F.D., Harwood, J.L. & Padley, F.B. (1994). *The Lipid Handbook* (2nd ed). Cambridge: Chapman & Hall.
- Gunstone, F.D. & Hamilton, R.J. (Eds.) (2001). *Basic oleochemicals, oleochemical products and new industrial oils*. Sheffield: Sheffield Academic Press.
- Liu, K. (1999). *Soybeans: Chemistry, Technology, and Utilization*. Maryland: Aspen Publishers, Inc.
- Nelson, D.L. & Cox, M.M. (2000). *Lehninger Principles of Biochemistry* (3rd ed). New York: Worth Publishers.

บทความในวารสารวิชาการ

- Alderman, D.J. & Jones, E.B.G. (1971). Physiological requirements of two marine phycomycetes, *Althornia crouchii* and *Ostracoblabe implexa*. *Transactions of the British Mycological Society*, 57, 213-255.
- Bahnweg, G. (1979). Studies on the physiology of Thraustochytriales I: Growth requirements and nutrition of *Thraustochytrium* spp., *Schizochytrium* sp., *Japonochytrium* sp., *Ulkenia* spp. and *Labyrinthuloides* spp. *Veröffentlichen Institute Meeresforschung Bremerhaven*, 17, 245-268.
- Bajpai, P.K., Bajpai, P. & Ward, O.P. (1991). Optimization of Production of Docosahexaenoic Acid (DHA) by *Thraustochytrium aureum* ATCC 34304. *Journal of the American Oil Chemists Society*, 68(7), 509-514.
- Barclay, W.R., Meager, K.M. & Abril, J.R. (1994). Heterotrophic production of long chain omega-3 fatty acids utilizing algae and algae-like organisms. *Journal of Applied Phycology*, 6, 123-129.

- Belarbi, E.H., Molina, E. & Chisti, Y. (2000). A process for high yield and scaleable recovery of high purity eicosapentaenoic acid esters from microalgae and fish oil. *Enzyme Microbiology and Technology*, 26, 516-529.
- Bower, S.M. (1987). Artificial culture of *Labyrinthuloides haliotidis* (Protozoa: Labyrinthomorpha), a pathogenic parasite of abalone. *Canadian Journal of Zoology*, 65, 2013-2020.
- Burja, A.M., Radianingtyas, H., Windust, A. & Barrow, C.J. (2006). Isolation and characterization of polyunsaturated fatty acid producing *Thraustochytrium* species: screening of strains and optimization of omega-3 production. *Apply Microbial Biotechnol*, 10, 253-261.
- Certik, M. & Shimizu, S. (1999). Biosynthesis and regulation of microbial polyunsaturated fatty acid production. *Journal of Bioscience and Bioengineering*, 87(1), 1-14.
- Chen, F. (1996). High cell density culture of microalgae in heterotrophic growth. *Trends in Biotechnology*, 14, 421-426.
- Chen, X., Du, W. & Liu, D. (2008). Response surface optimization of biocatalytic biodiesel production with acid oil. *Biochemical Engineering Journal*, 40, 423-429.
- Cheng, M.H., Walker, T.H., Hulbert, G.J. & Raman, D.R. (1999). Fungal production of eicosapentaenoic and arachidonic acids from industrial waste streams and crude soybean oil. *Bioresource Technology*, 67, 101-110.
- Crawford, M.A., Costeloe, K., Ghebremeskel, K., Phlactos, A., Akirvin, L. & Stacey, F. (1997). Are deficits of arachidonic and docosahexaenoic acids responsible for the neutral and vascular complications of pre-term babies?. *American Journal Clinical Nutrition*, 66, 1032-1041.
- Darley, W.M., Porter, D. & Fuller, M.S. (1973). Cell wall composition and synthesis via golgi-directed scale formation in the marine eucaryote, *Schizochytrium aggregatum*, with a note on *Thraustochytrium* sp. *Architech Mikrobiol*, 90, 89-106

- de Swaaf, M.E., Pronk, J.T. & Sitjtsma, L. (2003). Fed-batch cultivation of the docosahexaenoic acid producing marine alga *Cryptocodinium cohnii* on ethanol. *Applied Microbiology and Biotechnology*, 61, 40-43.
- Fan, K.W., Chen, F., Jones, E.B.G. & Vrijmod, L.L.P. (2000). Utilization of food processing waste by Thraustochytrids. *Fungal Diversity*, 5, 185-194.
- Fan, K.W., Vrijmoed, L.L.P. & Jones, E.B.G. (2002). Physiological Studies of Subtropical Mangrove Thraustochytrids. *Botanica Marina*, 45, 50-57.
- Fan, K.W., Jiang, Y., Faan, Y.W. & Chen, F. (2007). Lipid characterization of mangrove Thraustochytrid - *Schizochytrium mangrovei*. *Journal of Agricultural and Food Chemistry*, 55, 2906-2910.
- Gill, I. & Valivety, R. (1997). Polyunsaturated fatty acids, part1: occurrence, biological activities and applications. *Trends in Biotechnology*, 15, 401-409.
- Giusto, N.M., Pasquaré, S.J., Salvador, G.A., Castagnet, P.I., Roque, M.E. & Ilincheta de Boschero, M.G. (2000). Lipid metabolism invertebrate retinal rod outer segments. *Progress in Lipid Research*, 39, 315-391.
- Goldstein, S. (1963a). Development and nutrition of a new species of *Thraustochytrium*. *American Journal of Botany*, 50, 271-279.
- Goldstein, S. (1963b). Studies of a new species of *Thraustochytrium* that displays light stimulated growth. *Mycologia*, 55, 799-811.
- Goldstein, S. (1963c). Morphological variation and nutrition of a new monocentric marine fungus. *Archives of Microbiology*, 45, 101-110.
- Goldstein, S. (1973). Zoosporic marine fungi- Thraustochytriaceae and Dermocystidiaceae. *Annual Review of Microbiology*, 161-163.
- Harrington, G.W. & Holz, G.G. (1968). The monoenoic and docosahexaenoic fatty acids of a heterotrophic dinoflagellate. *Biochimica Biophysica Acta*, 164, 137-139.
- Horrocks, L.A. & Yeo, Y.K. (1999). Health benefits of docosahexaenoic acid (DHA). *Pharmacological Research*, 40, 211-215.

- Iida, I., Nakahara, T., Yokochi, T., Kamisaka, Y., Yagi, H., Yamaoka, M., et al. (1996). Improvement of docosahexaenoic acid production in a culture of *Thraustochytrium aureum* by medium optimization. *Journal of Fermentation and Bioengineering*, 81, 76-78.
- Jiang, Y. & Chen, F. (2000). Effect of medium glucose concentration and pH on docosahexaenoic acid content of heterotrophic *Cryptocodinium cohnii*. *Process Biochemistry*, 35, 1205-1209.
- Jiang, Y., Fan, K.W., Wong, R.T.Y. & Chen, F. (2004). Fatty acid composition and squalene content of the marine microalga *Schizochytrium mangrovei*. *Journal of Agricultural and Food Chemistry*, 52, 1196-1200.
- Kamlangdee, N. & Fan, K.W. (2003). Polyunsaturated fatty acids production by *Schizochytrium* sp. Isolated from mangrove. *Songklanakarin Journal of Science and Technology*, 25(5), 643-650.
- Kumon, Y., Yokochi, T. & Nakahara, T. (2005). High yield of long-chain polyunsaturated fatty acids by labyrinthulids on soybean lecithin-dispersed agar medium. *Biotechnological Products and Process Engineering*, 69, 253-258.
- Leano, E.M., Gapasin, R.S.J., Polohan, B. & Vrijmoed, L.L.P. (2003). Growth and fatty acid production of thraustochytrids from Panay mangroves, Philippines. *Fungal Diversity*, 12, 111-122.
- Leonard, A.E., Pereira, S.L., Sprecher, H. & Huang, Y.S. (2004). Elongation of long-chain fatty acids. *Lipid Research*, 43, 36-54.
- Lewis, T.E., Nichols, P.D. & McMeekin, T.A. (1999). The biotechnological potential of Thraustochytrids. *Marine Biotechnology*, 1, 580-587.
- Martinetz, M. (1992). Tissue levels of polyunsaturated fatty acids during early human development. *Journal of Podiatry*, 120, 129-138.
- Metz, J.G., Roessler, P., Facciotti, D., Levering, C., Dittrich, F., Lassner, M., et al. (2001). Production of polyunsaturated fatty acids by polyketide synthases in both prokaryotes and eukaryotes. *Science*, 293, 290-293.
- Mukherjee, K.D. (1999). Production and use of microbial oils. *Inform*, 10, 308-313.

- Nichols, D.S. (2003). Prokaryotes and the input of polyunsaturated fatty acids to the marine food web. *FEMS Microbiology*, 219, 1-7.
- Pereira, S.L., Leonard, A.E., Huang, Y.S., Chuang, L.T. & Mukerji, P. (2004). Identification of two novel microalgal enzymes involved in the conversion of the ω -3-fatty acid, eicosapentaenoic acid, into docosahexaenoic acid. *Biochemistry Journal*, 384, 357-366.
- Qiu, X. (2003). Biosynthesis of docosahexaenoic acid (DHA, 22:6-4, 7, 10, 13, 16, 19): two distinct pathways. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 68, 181-186.
- Ratlidge, C. (2004). Fatty acid biosynthesis in microorganisms being used for single cell oil production. *Biochimie*, 86, 807-815.
- Rees, J.F., Cure, K., Piyatiratitivorakul, S., Sorgeloos, P. & Menasveta, P. (1994). Highly unsaturated fatty acid requirements of *Penaeus monodon* postlarvae-and experimental approach based on *Artemia* enrichment. *Aquaculture*, 122, 193-207.
- Shabala, L., Shabala, S., Ross, T. & McMeekin, T. (2001). Membrane transport activity and ultradian ion flux oscillation associated with cell cycle of *Thraustochytrium* sp. *Australian Journal of Plant Physiology*, 28, 87-99.
- Simopoulos, A.P. (1999). Essential fatty acids in health and chronic disease. *American Journal of Clinical and Nutrition*, 70, 560-569.
- Singh, A. & Ward, O.P. (1996). Production of high yields of docosahexaenoic acid by *Thraustochytrium roseum* ATCC 20810. *Journal of Industrial Microbiology*, 16, 630-373.
- Spector, A.A. (1999). Essentiality of fatty acids. *Lipids*, 34, 1-3.
- Uauy, R., Hoffman, D.R., Peirano, P., Birch, B.G. & Birch, E.E. (2001). Essential fatty acids in visual and brain development. *Lipids*, 36, 885-895.
- Viljoen, B.C., Kock, J.L.F. & Lategan, P.M. (1986). The influence of culture age on the cellular fatty acid composition of four selected yeasts. *Journal of General Microbiology*, 132, 1895-1898.

- Ward, O.P. & Singh, A. (2005). Omega3/6 fatty acids: Alternative sources of production. *Process Biochemistry*, 40, 3627-3652.
- Wu, S.T., Yu, S.T. & Lin, L. (2005). Effect of culture conditions on docosahexaenoic acid production by *Schizochytrium* sp. S31. *Process Biochemistry*, 40, 3103-3108.
- Yamasaki, T., Aki, T., Shinozaki, M., Taguchi, M., Kawamoto, S. & Ono, K. (2006). Utilization of *Shochu* Distillery wastewater for production of polyunsaturated fatty acids and xanthophylls using thraustochytrid. *Journal of Bioscience and Bioengineering*, 102(4), 323-327.
- Yaguchi, T., Tanaka, S., Yokochi, T., Nakahara, T. & Higashihara, T. (1997). Production of high yields of docosahexaenoic acid by *Schizochytrium* sp. strain SR21. *Journal of the American Oil Chemists Society*, 74(11), 1431-1434.
- Yokochi, T., Honda, D., Higashihara, T. & Nakahara, T. (1998). Optimization of docosahexaenoic acid production by *Schizochytrium limacinum* SR21. *Applied Microbiology and Biotechnology*, 49, 72-76.
- Yoshida, H., Saiki, M., Yoshida, N., Tomiyama, Y. & Mizushima, Y. (2009). Fatty acid distribution in triacylglycerols and phospholipids of broad beans (*Vicia faba*). *Food Chemistry*, 112, 924-928.
- Yongmanitchai, W. & Ward, P. (1989). Omega-3 fatty acids: alternative sources of production. *Process Biochemistry*, 24, 117-125.

วิทยานิพนธ์และเอกสารอื่นๆ

- Gawrisch, K. & Eldho, N. (2002). Docosahexaenoic vs docosapentaenoic acid: the difference that the loss of a single double bond makes. *Fifth Congress of the International Society for the Study of Fatty Acids and Lipids*, Montreal.
- Krongkaew Ramwong. (2007). *Growth and physiology of Schizochytrium mangrovei Sk-02 grown on oils and/or free fatty acids*. Unpublished master's thesis, Mahidol University, Faculty of graduate studies.

Panida Unagul. (2006). *Production of docosahexaenoic acid by Schizochytrium mangrovei Sk-02*. Unpublished doctoral dissertation, King Mongkut's University of Technology Thonburi, School of Bioresources and Technology.

โรงงานอาหารผลิตภัณฑ์น้ำมันพืช. สืบค้นเมื่อวันที่ 10 มกราคม 2009, http://www.cook.co.th/cook_process_flow.htm