

VIII. REFERENCES

- Ajello L. Occurrence of *Cryptococcus neoformans* in soils. Am J Hyg. 1958;67:72-77.
- Baro T, Torres-Rodriguez JM, Morera Y, Alia C, Lopez O, Mendez R. Serotyping of *Cryptococcus neoformans* isolates from clinical and environmental sources in Spain. J Clin Microbiol. 1999; 37:1170-1172.
- Bauwens L, Swinne D, DeVroey C, DeMeurichy W. Isolation of *Cryptococcus neoformans* var. *neoformans* in the aviaries of the Antwerp Zoological Gardens. Mykosen. 1986;29:291-4.
- Benham RW. Cryptococci, their identification by morphology and serology. J Infect Dis. 1935;57:255-274.
- Benham RW. Cryptococcosis and blastomycosis. Ann NY Acad Sci. 1950;50:1299-1314.
- Bennett JE, Bailey JW. Control for rheumatoid factor in the latex test for cryptococcosis. Am J Clin Pathol. 1971;56:360-365.
- Bennett JE, Kwon-Chung KJ, Howard D. Epidemiologic differences among serotypes of *Cryptococcus neoformans*. Am J Epidemiol. 1984;105:582-586.
- Beyt BE, Waltman SR. Cryptococcal endophthalmitis after corneal transplantation. N Engl J Med. 1978;298:825-826.
- Bhattacharjee AK, Bennett JE, Glaudemans CP. Capsular polysaccharides of *Cryptococcus neoformans*. Rev Infect Dis. 1984;6:619-24.

- Bloomfield N, Gordon MA, Elmendorf DF, Jr. Detection of *Cryptococcus neoformans* antigen in body fluids by latex particle agglutination. Proc Soc Exp Biol Med. 1963;114:64-67.
- Brandt ME, Bragg SL, Pinner RW. Multilocus enzyme typing of *Cryptococcus neoformans*. J Clin Microbiol. 1993;31:2819-2823.
- Brandt ME, Hutwagner LC, Klug LA, Baughman WS, Rimland D, Graviss ED, et al. Molecular subtypes distribution of *Cryptococcus neoformans* in four areas of the United States. J Clin Microbiol. 1996;34:912-917
- Brandt ME, Hutwagner LC, Kuykendall RJ, Pinner RW, The Cryptococcal Disease Active Surveillance Group. Comparison of multilocus enzyme electrophoresis and random amplified polymorphic DNA analysis for molecular subtyping of *Cryptococcus neoformans*. J Clin Microbiol. 1995;33:1890-1895.
- Bruatto M, Vidotto V, Maina AM. Growth of *Cryptococcus neoformans* in thiamine-free media. Mycopathologia. 1992;119:129-132.
- Buchanan KL, Murphy JW. What makes *Cryptococcus neoformans* a pathogen? Emerg Infect Dis. 1998;4:71-83.
- Bulmer GS. Twenty-five years with *Cryptococcus neoformans*. Mycopathology. 1990;109:111-122.
- Buschke A. Ueber eine durch Coccidien hervorgerufene Krankheit des Menschen. Dtsch Med Wochenschr. 1895;21:14.
- Busse O. Ueber parasitaere Zelleinschlusse and ihre Zuchtung. Int J Med Microbiol. 1894;16:175-180.
- Callejas A, Ordonez N, Rodriguez MC, Castoneda E. First isolation of *Cryptococcus*

neoformans var. *gattii* serotype C from the environment in Colombia. Med Mycol. 1998;36:341-344.

Casadevall A, Freundlich LF, Marsh L, Scharff MD. Extensive allelic variation in *Cryptococcus neoformans*. J Clin Microbiol. 1992;30:1080-1084.

Casadevall A, Perfect R. *Cryptococcus neoformans*. Washington, D.C.: American Society for Microbiology, 1998.

Chakrabarti A, Jatana M, Kumar P, Chatha L, Kaushal A, Padhye AA. Isolation of *Cryptococcus neoformans* var. *gattii* from *Eucalyptus camaldulensis* in India. J Clin Microbiol. 1997;35:3340-3342.

Chang YC, Kwon-Chung KJ. Complementation of a capsule-deficient mutation of *Cryptococcus neoformans* restores its virulence. Mol Cell Biol. 1994;14:4912-4919.

Chang YC, Kwon-Chung KJ. Isolation of the third capsule-associated gene, *CAP60*, required for virulence in *Cryptococcus neoformans*. Infect Immun. 1998; 66:2230-2236.

Chang YC, Kwon-Chung. Isolation, Characterization and Localization of a Capsule-Associated Gene, *CAP10*, of *Cryptococcus neoformans*. J Bacteriol. 1999;181:5636-5643.

Chang YC, Penoyer LA, Kwon-Chung KJ. The second capsule gene of *Cryptococcus neoformans*, *CAP64*, is essential for virulence. Infect Immun. 1996; 64:1977-1983.

Chasakes S, Tyndall RL. Pigment production by *Cryptococcus neoformans* from para- and ortho- di-phenols: effect of the nitrogen source. J Clin Microbiol. 1975; 1:509-514.

- Chen SCA, Brownlee AG, Sorrell TC, Ruma P, Ellis DH, Pfeiffer TJ, et al.
Identification by random amplification of polymorphic DNA of a common molecular type of *Cryptococcus neoformans* var. *neoformans* in patients with AIDS or other immunosuppressive conditions. *J Infect Dis.* 1996;173:754-758.
- Cherniak R, Reiss E, Slodki ME, Plattner RD, Blumer SO. Structure and antigenic activity of the capsular polysaccharide of *Cryptococcus neoformans*. *Mol Immunol.* 1980;17:1025-32.
- Cherniak R, Sundstrom JB. Polysaccharide antigens of the capsule of *Cryptococcus neoformans*. *Infect Immun.* 1994;62:1507-12.
- Clancy MN, Fleischmann J, Howard DH, Kwon-Chung KJ, Shimizu RY. Isolation of *Cryptococcus neoformans* var. *gattii* from a patient with AIDS in Southern California. *J Infect Dis.* 1990;161:809.
- Cooper BH, Silva-Hutner M. Yeasts of medical importance. In: Lennette EH, ed. *Manual of clinical microbiology*. 4th ed. Washington, DC: American Society for Microbiology, 1985;526-41.
- Crampin AC, Matthews RC, Hall D, Evans EGV. PCR fingerprinting *Cryptococcus neoformans* by random amplification of polymorphic DNA. *J Med Vet Mycol.* 1993;31:463-465.
- Currie BP, Freundlich LF, Casadevall A. Restriction fragment length polymorphism analysis of *Cryptococcus neoformans* isolates from environmental (pigeon excreta) and clinical sources in New York City. *J Clin Microbiol.* 1994;32:1188-1192.
- DeVroey C, Swinne D. Isolements de *Cryptococcus neoformans* a l'occasion de concours de chant de canaris (+). *Bull Soc Fr Mycol Med.* 1986;15:353-6.

- Division of Epidemiology, Ministry of Public Health (Thailand). Annual epidemiological surveillance report 1999. Bangkok. The Division;1999.
- Dromer F, Gueho E, Ronin O, Dupont B. Serotyping of *Cryptococcus neoformans* by using a monoclonal antibody specific for capsular polysaccharide. J Clin Microbiol. 1993;31:359-63.
- Dromer F, Mathoulin S, Dupont B, Laporte A, Ronin O, the French Cryptococcosis Study Group. Individual and environmental factors associated with infection due to *Cryptococcus neoformans* serotype D. Clin Infect Dis. 1996b;23:91-96.
- Dromer F, Mathoulin S, Dupont B, Laporte A, the French Cryptococcosis Study Group. Epidemiology of cryptococcosis in France: a 9-year survey (1985-1993). Clin Infect Dis. 1996a;23:82-90.
- Dromer F, Varma A, Ronin A, Mathoulin S, Dupont B. Molecular typing of *Cryptococcus neoformans* serotype D clinical isolates. J Clin Microbiol. 1994;32:2364-2371.
- Ellis D, Pfeiffer TJ. The ecology of *Cryptococcus neoformans*. Eur J Epidemiol. 1992;8:321-325.
- Ellis DH, Pfeiffer TJ. Cryptococcosis and the ecology of *Cryptococcus neoformans*. Jpn J Med Mycol. 1994;35:111-122.
- Ellis DH, Pfeiffer TJ. Natural habitat of *Cryptococcus neoformans* var. *gattii*. J Clin Microbiol. 1990;28:1642-1644.
- Emmons CW. Isolation of *Cryptococcus neoformans* from soil. J Bacteriol. 1951;62:685-90.
- Emmons CW. Saprophytic sources of *Cryptococcus neoformans* associated with the

- pigeon (*Columba livia*). *Am J Hyg.* 1955;62:227-32.
- Erke KH, Schneidau JD. Relationship of some *Cryptococcus neoformans* hypha-forming strains to standard strains and to other species of yeasts as determined by deoxyribonucleic acid base ratios and homologies. *Infect Immun.* 1973;7:941-948.
- Fan M, Currie BP, GutellRR, Ragen MA, Casadevall A. The 16s-like, 5.8s and 23s-like rRNA of the two varieties of *Cryptococcus neoformans*: sequence secondary structure, phylogenetic analysis, and restriction fragment polymorphisms. *J Med Vet Mycol.* 1994;32:163-180.
- Fragner P. The findings of Cryptococci in excrements of birds. *Czech Epidem Micro Immun.* 1962;11:135-139.
- Frank UK, Nishimura SL, Li NC, Sugai K, Yajko DM, Hadley WK, Ng VL. Evaluation of an enzyme immunoassay for detection of cryptococcal capsular polysaccharide antigen in serum and cerebrospinal fluid. *J Clin Microbiol.* 1993;31:97-101.
- Franzot SP, Fries BC, Cleare W, Casadevall A. Genetic Relationship between *Cryptococcus neoformans* var. *neoformans* Strains of Serotypes A and D. *J Clin Microbiol.* 1998; 36:2200-2204.
- Franzot SP, Hamdan JS, Currie BP, Casadevall A. Molecular epidemiology of *Cryptococcus neoformans* in Brazil and the United States: evidence for both local genetic differences and a global clonal population structure. *J Clin Microbiol.* 1997;35:2243-2251.
- Franzot SP, Salkin IF, Casadevall A. *Cryptococcus neoformans* var. *grubii*: Separate varietal status for *Cryptococcus neoformans* serotype A isolates. *J Clin Microbiol.* 1999;37:838-840.

- Fromtling RA, Shadomy HJ, Jacobson ES. Decreased virulence in stable, acapsular mutants of *Cryptococcus neoformans*. *Mycopathologia*. 1982;79:23-9.
- Gade W, Hinnefeld SW, Babcock LS, Gilligan P, Kelly W, Wait K, et al. Comparison of the premier cryptococcal antigen enzyme immunoassay and the latex agglutination assay for detection of cryptococcal antigens. *J Clin Microbiol*. 1991;29:1616-1619.
- Garcia-Hermoso D, Janbon G, Dromer F. Epidemiological evidence for dormant *Cryptococcus neoformans* infection. *J Clin Microbiol*. 1999;37:3204-3209.
- Gezuele E, Calegari L, Sanabria D, Davel G, Civilia E. Isolation of *Cryptococcus neoformans* var. *gattii* from nest of the wasp. *Polybia occidentalis*. *Rev Iberoam Micol*. 1993;10:5-6.
- Gorgon MA, Devine J. Filamentation and endogenous sporulation in *Cryptococcus neoformans*. *Sabouraudia*. 1970;8:227-234.
- Granger DL, Perfect JR, Durack DT. Virulence of *Cryptococcus neoformans*: regulation of capsule synthesis by carbon dioxide. *J Clin Microbiol*. 1985;76:508-516.
- Griseo G, Bolignano MS, De Leo F, Staib F. Evidence of canary droppings as an important reservoir of *Cryptococcus neoformans*. *Zentralbl Bakteriol*. 1995;282:244-254.
- Gueho E, Improvisi L, Christen R, de Hoog GS. Phylogenetic relationships of *Cryptococcus neoformans* and some related basidiomycetous yeasts determined from partial large subunit rRNA sequences. *Antonie van Leeuwenhoek*. 1993;63:175-189.
- Haynes KA, Sullivan DJ, Coleman DC, Clarke JCK, Emilianus R, Atkinson J, Cann

- KJ. Involvement of multiple *Cryptococcus neoformans* strains in a single episode of cryptococcosis and reinfection with novel strains in recurrent infection demonstrated by random amplification of polymorphic DNA and DNA fingerprinting. *J Clin Microbiol.* 1995;33:99-102.
- Ikeda RT, Shinoda T, Morita T, Jacobson ES. Characterization of a phenol oxidase from *Cryptococcus neoformans* var. *neoformans*. *Microbiol Immunol.* 1993;37:759-764.
- Ishaq CM, Bulmer GS, Felton FG. An evaluation of various environmental factors affecting the propagation of *Cryptococcus neoformans*. *Mycopathol Mycol Appl.* 1968;35:81-90.
- Iwata K, Yamashita T, Oshumi M, Baba M, Naito N, Taki A, Yamada N. Comparative morphological and biological studies on the itraconazole- and ketoconazole-resistant mutants of *Cryptococcus neoformans*. *J Med Vet Mycol.* 1990;28:77-90.
- Kersulyte D, Woods JP, Keath EJ, Boldman WE, Berg DE. Diversity among clinical isolates of *Histoplasma capsulatum* detected by polymerase chain reaction with arbitrary primer. *J Bacteriol.* 1992;174:7075-7079.
- Klein E. Pathogenic microbes in milk. *J Hyg (Cambridge)* 1901;1:78-95.
- Khayhan S. Serotyping and DNA analysis by PCR-fingerprinting of clinical and environmental isolates of *Cryptococcus neoformans* in Chiang Mai. M.S. Thesis in microbiology, Department of Microbiology, Faculty of Medicine, Chiang Mai University, 2001.
- Kozel TR, Cazin J Jr. Non-encapsulated variant of *Cryptococcus neoformans* I. Virulence studies and characterization of soluble polysaccharide. *Infect Immun.* 1971;3:287-94.

- Kwon-Chung KJ, Bennett JE, Rhodes JC. Taxonomic studies on *Filobasidiella* species and their anamorphs. *Antonie van Leeuwenhoek*. 1982a;48:25-38.
- Kwon-Chung KJ, Bennett JE. Epidemiologic differences between the two varieties of *Cryptococcus neoformans*. *Am J Epidemiol*. 1984;120:123-130.
- Kwon-Chung KJ, Bennett JE. *Medical Mycology*. Pennsylvania: Lea and Febiger, 1992.
- Kwon-Chung KJ, Hill WB, Bennett JE. New special stain for histopathological diagnosis of cryptococcosis. *J Clin Microbiol*. 1981;13:383-387.
- Kwon-Chung KJ, Polacheck I, Bennett JE. Improved diagnostic medium for separation of *Cryptococcus neoformans* var. *neoformans* (serotype A and D) and *Cryptococcus neoformans* var. *gattii* (serotype B and C). *J Clin Microbiol*. 1982b;15:535-537.
- Kwon-Chung KJ, Polacheck I, Popkin TJ. Melanin-lacking mutants of *Cryptococcus neoformans* and their virulence for mice. *J Bacteriol*. 1982c;150:1414-1421.
- Kwon-Chung KJ, Rhodes JC. Encapsulation and melanin formation as indicators of virulence in *Cryptococcus neoformans*. *Infect Immun*. 1986;51:218-23.
- Kwon-Chung KJ, Varma A. *Cryptococcus neoformans*-ecology and epidemiology. In: 3rd symposium topics in mycology, mycoses in AIDS patients. Paris: Pasteur Institute, November, 1989;55-56 (abstr).
- Kwon-Chung KJ, Wickes BL, Stockman L, Roberts GD, Ellis D, Howard DH. Virulence, serotype, and molecular characteristics of environmental strains of *Cryptococcus neoformans* var. *gattii*. *Infect Immun*. 1992;60:1869-1874.

- Kwon-Chung KJ. A new genus, *Filobasidiella*, the perfect state of *Cryptococcus neoformans*. *Mycologia*. 1975;67:1197-1200.
- Kwon-Chung KJ. A new species of *Filobasidiella*, the sexual state of *Cryptococcus neoformans* B and C serotypes. *Mycologia*. 1976b;68:942-946.
- Kwon-Chung KJ. Cryptococcosis. In: Kwon-Chung KJ, Bennett JE, editors. *Medical mycology*. Philadelphia: Lea & Febiger, 1992:397-446.
- Kwon-Chung KJ. Morphogenesis of *Filobasidiella neoformans*, the sexual state of *Cryptococcus neoformans*. *Mycologia*. 1976a;68:821-833.
- Larsen RA, Leal MAE, Chan LS. Fluconazole compared with amphotericin B plus flucytosine for cryptococcal meningitis in AIDS; a randomized trial. *Ann Intern Med*. 1990;113:183-187.
- Laurenson IF, Lalloo DG, Naraqi S, et al. *Cryptococcus neoformans* in Papua New Guinea: a common pathogen but an elusive source. *J Med Vet Mycol*. 1997;35:437-440.
- Lazera MS, Pires FDA, Camillo-Coura L, et al. Natural habitat of *Cryptococcus neoformans* var. *neoformans* in decaying wood forming hollows in living trees. *J Med Vet Mycol*. 1996;34:127-131.
- Lazera MS, Salmito Cavalcanti MA, Londero AT, Trilles L, Nishikawa MM, Wanke B. Possible primary ecological niche of *Cryptococcus neoformans*. *Med Mycol*. 2000;38:379-383.
- Lazera MS, Wanke B, Nishikawa MM. Isolation of both varieties of *Cryptococcus neoformans* from saprophytic sources in the city of Rio de Janeiro, Brazil. *J Med Vet Mycol*. 1993;31:449-454.

- Levitz SM. The ecology of *Cryptococcus neoformans* and the epidemiology of cryptococcosis. *Rev Infect Dis.* 1991;13:1163-1169.
- Littman ML, Walter JE. Cryptococcosis: current status. *Am J Med.* 1968;45:922-32.
- Littman ML. Capsule synthesis by *Cryptococcus neoformans*. *Trans NY Acad Sci.* 1958;20:623-648.
- Lopez-Martinez R, Castanon-Olivares LR. Isolation of *Cryptococcus neoformans* var. *neoformans* from bird droppings, fruits and vegetables in Mexico City. *Mycopathologia.* 1995;129:25-28.
- McDonough ES, Ajello L, Ausherman RJ, Balows A, McClellan JT, Brinkman S. Human pathogenic fungi recovered from soil in an area endemic for North American blastomycosis. *Am J Hyg.* 1961;73:75-83.
- Meyer W, Lieckfeldt E, Kuhls K, Freedman EZ, Borner T, Mitchell TG. DNA- and PCR-fingerprinting in fungi. 1993a, p. 311-320. *In* Pena SDJ, Chakraborty R, Epplen JT, Jeffreys AJ (ed.), *DNA Fingerprinting: State of the Science.* Birkhauser Verlag, Basel.
- Meyer W, Mitchell TG, Freedman EZ, Vilgalys R. Hybridization probes for conventional DNA fingerprinting used as single primers in the polymerase chain reaction to distinguish strains of *Cryptococcus neoformans*. *J Clin Microbiol.* 1993b;31:2274-2280.
- Meyer W, Mitchell TG. PCR fingerprinting to distinguish species and strains of yeasts. 1994, p. 293-302. *In* Maresca B and Kobayashi GS (ed.), *Molecular Biology of Pathogenic Fungi: A Laboratory Manual.* Telos Press, New York.
- Meyer W, Mitchell TG. Polymerase chain reaction fingerprinting in fungi using single primers specific to minisatellites and simple repetitive DNA sequences: strain

- variation in *Cryptococcus neoformans*. Electrophoresis. 1995;16:1648-1656.
- Miller MF, Mitchell TG, Storkus WJ, Dawson JR. Human natural killer cells do not inhibit growth of *Cryptococcus neoformans* in the absence of antibody. Infect Immun. 1990;58:639-645.
- Mitchell DH, Sorrell TC, Allworth AM, et al. Cryptococcal disease of the CNS in immunocompetent hosts: influence of cryptococcal variety on clinical manifestations and outcome. Clin Infect Dis. 1995;20:611-6.
- Mitchell TG, Freedman EZ, White TJ, Taylor JW. Unique oligonucleotide primers in PCR for identification of *Cryptococcus neoformans*. J Clin Microbiol. 1994a;32:253-255.
- Mitchell TG, Perfect JR. Cryptococcosis in the ere of AIDS--- 100 years after the discovery of *Cryptococcus neoformans*. Clin Microbiol Rev. 1995;8:515-548.
- Mitchell TG, Sandin RL, Bowman BH, Meyer W, Merz WG. Molecular mycology: DNA probes and applications of PCR technology. J Med Vet Mycol. 1994b;32 (suppl. 1):351-366.
- Mody CE, Toews GB, Lipscomb MF. Cyclosporin A inhibits the growth of *Cryptococcus neoformans* in a murine model. Infect Immun. 1988;56:7-12.
- Montagna MT, Viviani MA, Pulito A, et al. *Cryptococcus neoformans* var. *gattii* in Italy, Environmental investigation related to an autochthonous clinical case in Apulia. J Mycol Med. 1997;7:93-96.
- Montenegro H, Paula CR. Environmental isolation of *Cryptococcus neoformans* var. *gattii* and *Cryptococcus neoformans* var. *neoformans* in the city of Sao Paulo, Brazil. Med Mycol. 2000;38:385-390.

- Muchmore RE. Occurrence of *Cryptococcus neoformans* in the environment of three geographically associated cases of cryptococcal meningitis. *N Engl J Med*. 1963;268:1112-4.
- Murray PR, Rosenthal KS, Kobayashi Gs, Pfaller MA. *Medical Microbiology*. 4th ed. Missouri: Mosby, 2002.
- Neilson JB, Fromtling RA, Bulmer GS. *Cryptococcus neoformans*: size range of infectious particles from aerosolized soil. *Infect Immun*. 1977;17:634-638.
- Odom A, Muir S, Lim E, Toffaletti DL, Perfect J, Heitman J. Calcineurin is required for virulence of *Cryptococcus neoformans*. *EMBO J*. 1997;16:2576-89.
- Pal M, Mehrotra BS. Studies on the isolation of *Cryptococcus neoformans* from fruits and vegetables. *Mykosen*. 1984;28:200-5.
- Passo CL, Pernice I, Gallo M, Barbara C, Luck ET, Criseo G, Pernice A. Genetic relatedness and diversity of *Cryptococcus neoformans* strains in the Maltese Islands. *J Clin Microbiol*. 1997;35:751-755.
- Perfect JR, Durack DT. Effects of cyclosporin in experimental cryptococcal meningitis. *Infect Immun*. 1985;50:22-26.
- Perfect JR, Ketabchi NK, Cox GM, Ingram CI, Beiser C. Karyotyping of *Cryptococcus neoformans* as an epidemiological tool. *J Clin Microbiol*. 1993;31:3305-3309.
- Perfect JR, Magee BB, Magee PT. Separation of chromosomes of *Cryptococcus neoformans* by pulsed field gel electrophoresis. *Infect Immun*. 1989;57:2624-2627.
- Perfect JR, Wong T, Chang YC, Kwon-Chung KJ, Williamson PR. *Cryptococcus*

neoformans: virulence and host defences. Med Mycol. 1998;36:79-86.

Persing DH, Smith TF, Tenover FC, White TJ. Diagnostic molecular microbiology: principles and applications. 1993; p. 1-641. American Society for Microbiology, Washington, D.C.

Pfeiffer TJ, Ellis DH. Additional eucalyptus hosts for *Cryptococcus neoformans* var. *gattii*. In: Abstracts of the 13th Congress of the International Society for Human and Animal Mycology (ISHAM), Parma, Italy: Jun 8-13, 1997. Parma: University of Parma, 1997: Abstract P111.

Pfeiffer TJ, Ellis DH. Environmental isolation of *Cryptococcus neoformans* var. *gattii* from *Eucalyptus tereticornis*. J Med Vet Mycol. 1992;30:407-408.

Polacheck I, Kwon-Chung KJ. Melanogenesis in *Cryptococcus neoformans*. J Gen Microbiol. 1988;134:1037-41.

Polacheck I, Lebens G, Hicks JB. Development of DNA probes for early diagnosis and epidemiological study of cryptococcosis in AIDS patients. J Clin Microbiol. 1992;30:925-930.

Polacheck I, Lebens GA. Electrophoretic karyotype of the pathogenic yeast *Cryptococcus neoformans*. J Gen Microbiol. 1989;135:67-71.

Polacheck I, Platt Y, Aronovitch J. Catecholamines and virulence of *Cryptococcus neoformans*. Infect Immun. 1990;58:2919-2922.

Poonwan N, Mikami Y, Poonswan S, et al. Serotyping of *Cryptococcus neoformans* strains isolated from clinical specimens in Thailand and their susceptibility to various antifungal agents. Eur J Epidemiol. 1997;13:335-340.

Powell KE, Dahl BA, Weeks RJ, Tosh FE. Airborne *Cryptococcus neoformans* :

particles from pigeon excreta compatible with alveolar deposition. J Infect Dis. 1972;125:412-415.

Prariyachatigul C, Chaiprasert A, Meevootisom V, Pattanakitsakul S. Assessment of a PCR technique for the detection and identification of *Cryptococcus neoformans*. J Med Vet Mycol. 1996; 34:251-258.

Randhawa HS, Paliwal DK. Occurrence and significance of *Cryptococcus neoformans* in the oropharynx and on the skin of a healthy human population. J Clin Microbiol. 1977;6:325-327.

Restrepo BI, Barbour AG. Cloning of the 18S and 25S ribosomal DNA from the pathogenic fungus *Cryptococcus neoformans*. J Bacteriol. 1989;171:5596-5600.

Rippon JW. Medical mycology. The pathogenic fungi and the pathogenic actinomycetes, 3rd ed. Philadelphia: WB Saunders, 1998;582-609.

Rippon JW. Medical mycology. The pathogenic fungi and the pathogenic actinomycetes, 2nd ed. Philadelphia: WB Saunders, 1982;542-58.

Ro JY, Lee SS, Ayala AG. Advantage of Fontana-Masson stain in capsule-deficient cryptococcal infection. Arch Pathol Lab Med. 1987;111:53-57.

Sanfelice F. Contributo alla morfologia e biologia dei blastomiceti che si sviluppano nei succhi di alcuni frutti. Ann Igiene. 1894;4:463-495.

Sanfelice F. Über einen neuen pathogenen Blastomyceten, welcher innerhalb des Gewebe unter Bildung Kalkartig aussenhender Massendegeneration. Int J Med Microbiol. 1895;18:521-6.

Satitnimarnkarn T, Na Nakorn S, Rigunti M. Cryptococcosis report of first case in Thailand. J Med Assoc. Thailand. 1960; 43:78-80.

- Segal E, Baum GL. Pathogenic yeasts and yeast infections. 1994.
- Sekhon AS, Garg AK, Kaufman L, Kobayashi GS, Hamir Z, Jalbert M, Moledina N. Evaluation of a commercial enzyme immunoassay for the detection of a cryptococcal antigen. *Mycoses*. 1993;36:31-34.
- Shaw CE, Kapica L. Production of diagnostic pigment by phenoloxidase activity of *Cryptococcus neoformans*. *Applied Microbiology*. 1972;24:824-30.
- Soll D. DNA fingerprinting the infectious fungi. *Clin Microbiol Rev*. 2000;13:331-370.
- Sorrell TC, Chen SCA, Ruma P, Meyer N, Pfeiffer TJ, Ellis DH, Brownlee AC. Concordance of clinical and environmental isolates of *Cryptococcus neoformans* var. *gattii* by random amplification of polymorphic DNA analysis and PCR fingerprinting. *J Clin Microbiol*. 1996;34:1253-1260.
- Spitzer ED, Spitzer SG. Use of a dispersed repetitive DNA element to distinguish clinical isolates of *Cryptococcus neoformans*. *J Clin Microbiol*. 1992;30:1094-1097.
- Spitzer SG, Spitzer ED, Freundlich LF, Casadevall A. Persistence of initial infection in recurrent *Cryptococcus neoformans* meningitis. *Lancet*. 1993;341:595-596.
- Spitzer SG, Spitzer ED. Characterization of the CNRE-1 family of repetitive DNA elements in *Cryptococcus neoformans*. *Gene*. 1994;144:103-106.
- Staib F, Seibold M, Antweiler E, Frohlich B. Staib agar supplemented with a triple antibiotic combination for the detection of *Cryptococcus neoformans* in clinical specimens. *Mycoses* 1989;32:448-54.
- Staib F, Seibold M. Use of the membrane filtration technique and Staib agar for the

detection of *Cryptococcus neoformans* in the urine of AIDS patients---a contribution to diagnosis, therapy and pathogenesis of cryptococcosis. *Mycoses*. 1989;32:63-72.

Staib F. *Cryptococcus neoformans* and *Guizottia abyssinica*, (Syn. *G. oleifera*)
Farbreaktion fur *C. neoformans*. *Z Hyg*. 1962;148:466-475.

Staib F. New concepts in the occurrence and identification of *Cryptococcus neoformans*. *Mycopathologia*. 1963;19:143-145.

Sugar AM, Stern JJ, Dupont B. Overview: treatment of cryptococcal meningitis. *Rev Infect Dis*. 1990;12(Suppl. 3):368-373.

Sukroongreung S, Nilakul C, Ruangsomboon O, Chuakul W, Eampokalap B.
Serotypes of *Cryptococcus neoformans* isolated from patients prior to and during the AIDS era in Thailand. *Mycopathologia*. 1996;135:75-78.

Swinne D, Kayembe K, Niyimi M. Isolation of saprophytic *Cryptococcus neoformans* var. *neoformans* in Kinshasa, Zaire. *Ann Soc Belg Med Trop*. 1986;66:57-61.

Swinne D, Taelman H, Batungwanayo J, Bigirankana A, Bogaerts J. Ecology of *Cryptococcus neoformans* in central Africa. *Med Trop*. 1994;54:53-55.

Tanner DC, Weinstein MP, Fedorciw B, Joho KL, Thorpe JJ, Reller LB. Comparison of commercial kits for detection of cryptococcal antigen. *J Clin Microbiol*. 1994;32:1680-1684.

Tarrand JJ, Guillot C, Wenglar M, Jackson J, Lajeunesse JD, Rolston KVI. Clinical comparison of the resin-containing BATEC 26 Plus and the Isolator 10 blood culturing systems. *J Clin Microbiol*. 1991;29:2245-2249.

Tharavichitkul P, Kanjanasthiti P, Panasampol K. Occurrence of *Cryptococcus*

- neoformans* in dove excreta. Chiang Mai Medical Bulletin. 1973;12:91-97.
- Torres-Guerrero H, Edman JC. Melanin-deficient mutants of *Cryptococcus neoformans*. J Med Vet Mycol. 1994;32:303-13.
- Vanbreuseghem R, Takashio M. An atypical strain of *Cryptococcus neoformans* (San Felice) Vuillemin 1894. Part II. *Cryptococcus neoformans* var. *gattii* var. nov. Ann Soc Belg Med Trop. 1970;50:695-702.
- Varma A, Kwon-Chung KJ. DNA probe for strain typing of *Cryptococcus neoformans*. J Clin Microbiol. 1992;30:2960-2967.
- Varma A, Kwon-Chung KJ. Restriction fragment polymorphism in mitochondrial DNA of *Cryptococcus neoformans*. J Gen Microbiol. 1989;135:3353-3362.
- Varma A, Swinne D, Staib F, Bennett JE, Kwon-Chung KJ. Diversity of DNA fingerprints in *Cryptococcus neoformans*. J Clin Microbiol. 1995;33:1807-1814.
- Vogel RA. The indirect fluorescent antibody test for the detection of antibody in human cryptococcal disease. J Infect Dis. 1966;116:573-580.
- Walter JE, Coffee EG. Control of *Cryptococcus neoformans* in pigeon coops by alkalization. Am J Epidemiol. 1968;87:173-178.
- Walter JE, Yee RB. Factors that determine the growth of *Cryptococcus neoformans* in avian excreta. Am J Epidemiol. 1968;88:445-450.
- Wang Y, Casadevall A. Decreased susceptibility of melanized *Cryptococcus neoformans* to UV light. Appl Environ Microbiol. 1994;60:3864-3866.
- Wang Y, Aisen P, Casadevall A. *Cryptococcus neoformans* melanin and virulence: mechanism of action. Infect Immun. 1995;63:3131-3136.

- Weinstein L, Jacoby I. Successful treatment of cerebral cryptococcoma and meningitis with miconazole. *Ann Intern Med.* 1980;93:569-571.
- Wickes BL, Moore TDE, Kwon-Chung KJ. Comparison of the electrophoretic karyotypes and chromosomal location of ten genes in the two varieties of *Cryptococcus neoformans*. *Microbiology.* 1994;140:543-550.
- Wong B, Perfect JR, Beggs S, Wright KA. Production of the hexitol D-mannitol by *Cryptococcus neoformans* in vitro and in rabbits with experimental meningitis. *Infect Immun.* 1990;58:1664-1670.
- Yamamoto Y, Kohno S, Koga H, Kakeya H, Tomono K, Kaku M, et al. Random amplification of polymorphic DNA analysis of clinical and environmental isolated *Cryptococcus neoformans* in Nagasaki. *J Clin Microbiol.* 1995;33:3328-3332.
- Zerpa R, Huicho L, Guillen A. Modified India ink preparation for *Cryptococcus neoformans* in cerebrospinal fluid specimens. *J Clin Microbiol.* 1996;34:2290-2291.