

References

- Anderson, J. L., Ding, R., Ellern, A., & Armstrong, D. W. (2004). Structure and properties of high stability geminal dicationic ionic liquids. *Journal of the American Chemical Society*, 127(2), 593-604.
- Armenta, S., Garrigues, S., & de la Guardia, M. (2008). Green analytical chemistry. *Trends in Analytical Chemistry*, 27(6), 497-511.
- Armstrong, D. W., & Henry, S. J. (1980). Use of an aqueous micellar mobile phase for separation of phenols and polynuclear aromatic hydrocarbons via HPLC. *Journal of Liquid Chromatography*, 3(5), 657-662.
- Arunyanart, M., & Cline-Love, L. J. (1984). Model for micellar effects on liquid chromatography capacity factors and for determination of micelle-solute equilibrium constants. *Analytical Chemistry*, 56(9), 1557-1561.
- Baker, D. (1995). *Capillary Electrophoreses*. New York: Wiley Interscience.
- Ballesta Claver, J., Valencia, M. C., & Capitán-Vallvey, L. F. (2009). Analysis of parabens in cosmetics by low pressure liquid chromatography with monolithic column and chemiluminescent detection. *Talanta*, 79(2), 499-506.
- Barron, L., & Paull, B. (2006). Simultaneous determination of trace oxyhalides and haloacetic acids using suppressed ion chromatography-electrospray mass spectrometry. *Talanta*, 69(3), 621-630.
- Belgaied, J. E., & Trabelsi, H. (2003). Determination of cisapride, its oxidation product, propyl and butyl parabens in pharmaceutical dosage form by reversed-phase liquid chromatography. *Journal of Pharmaceutical and Biomedical Analysis*, 33(5), 991-998.
- Beltran, A., Marcé, R. M., Cormack, P. A. G., & Borrull, F. (2010). Synthetic approaches to parabens molecularly imprinted polymers and their applications to the solid-phase extraction of river water samples. *Analytica Chimica Acta*, 677(1), 72-78.
- Bianco Prevot, A., Pramauro, E., Gallarate, M., Carlotti, M. E., & Orio, G. (2000). Determination of micelle/water partition coefficients of cosmetic

- preservatives: Optimisation of the capillary electrophoretic method. *Analytica Chimica Acta*, 412(1-2), 141-148.
- Bidlingmeyer, B. A. (1979). Retention mechanism for reversed-phase ion-pair liquid chromatography. *Journal of Chromatography*, 186, 419-434.
- Bidlingmeyer, B. A. (1980). Separation of ionic compounds by reversed-phase liquid chromatography: an update of ion-pairing techniques. *Journal of Chromatography*, 18, 525-539.
- Blanco, E., Casais, M. d. C., Mejuto, M. d. C., & Cela, R. (2009). Combination of off-line solid-phase extraction and on-column sample stacking for sensitive determination of parabens and p-hydroxybenzoic acid in waters by non-aqueous capillary electrophoresis. *Analytica Chimica Acta*, 647(1), 104-111.
- Blount, B. C., & Valentin-Blasini, L. (2006). Analysis of perchlorate, thiocyanate, nitrate and iodide in human amniotic fluid using ion chromatography and electrospray tandem mass spectrometry. *Analytica Chimica Acta*, 567(1), 87-93.
- Boyd, R. K., Basic, C., & Bethem, R. A. (2008). *Trace quantitative analysis by mass spectrometry*. West Sussex: John Wiley & Sons, Ltd.
- Breadmore, M. C., Haddad, P. R., & Fritz, J. S. (2001). Optimisation of the separation of anions by ion chromatography-capillary electrophoresis using indirect UV detection. *Journal of Chromatography A*, 920(1-2), 31-40.
- Bruzzoniti, M. C., Mentasti, E., & Sarzanini, C. (1997). Divalent pairing ion for ion-interaction chromatography of sulfonates and carboxylates. *Journal of Chromatography A*, 770(1), 51-57.
- Burns, D. T., & Tungkananuruk, N. (1987). Spectrophotometric determination of perchlorate after extraction of its brilliant green ion-pair with microcrystalline benzophenone. *Analytica Chimica Acta*, 199, 237-240.
- Calafat, A. M., Xiaoyun, Y., Lee-Yang, W., Bishop, A. M., & Needham, L. L. (2010). Urinary concentrations of four parabens in the U.S. Population: NHANES 2005-2006. *Environmental Health Perspectives*, 118(5), 679-685.
- Casoni, D., Kot-Wasik, A., Namieśnik, J., & Sârbu, C. (2009). Lipophilicity data for some preservatives estimated by reversed-phase liquid chromatography and

- different computation methods. *Journal of Chromatography A*, 1216(12), 2456-2465.
- Cecchi, T. (2010). *Ion-pair chromatography and related techniques*. Boca Raton: Taylor and Francis Group, LLC.
- Cecchi, T., Pucciarelli, F., & Passamonti, P. (2004). Ion interaction chromatography of zwitterions: the fractional charge approach to model the influence of mobile phase concentration of the ion-interaction reagent. *Analyst*, 129, 1037-1042.
- Cecchi, T., Pucciarelli, F., Passamonti, P., & Cecchi, P. (2001). The dipole approach in ion-interaction chromatography of zwitterions. *Chromatographia*, 54(1), 38-44.
- Charnley, G. (2008). Perchlorate: Overview of risks and regulation. *Food and Chemical Toxicology*, 46(7), 2307-2315.
- Chen, H.-C., Chen, W.-T., & Ding, W.-H. (2009). Determination of perchlorate in river by ion-pair hollow-fiber liquid-phase microextraction coupled with electrospray ionization tandem mass spectrometry. *Talanta*, 79(2), 442-445.
- Chu, Q., Wang, J., Zhang, D., & Ye, J. (2010). Sensitive determination of parabens in soy sauces by capillary zone electrophoresis with amperometric detection. *European Food Research and Technology*, 231(6), 891-897.
- Darbre, P. D., Aljarrah, A., Miller, W. R., Coldham, N. G., Sauer, M. J., & Pope, G. S. (2004). Concentrations of parabens in human breast tumours. *Journal of Applied Toxicology*, 24, 5-13.
- Dasgupta, P. K., & Martinelango, P. K. (2007). Dicationic ion-pairing agents for the mass spectrometric determination of perchlorate. *Analytical Chemistry*, 79(18), 7198-7200.
- De Hoffmann, E., & Stroobant, V. (2007). *Mass spectrometry: Principles and applications* (3rd ed.). West Sussex: John Wiley & Sons, Ltd.
- Driouich, R., Takayanagi, T., Oshima, M., & Motomizu, S. (2000). Separation and determination of haloperidol, parabens and some of their degradation products by micellar electrokinetic chromatography. *Journal of Chromatography A*, 903(1-2), 271-278.

- Environmental Working Group (2007). Perchlorate. Retrieved May 5, 2011, from <http://www.ewg.org/chemindex/chemicals/perchlorate>
- Eriksson, E., Andersen, H. R., & Ledin, A. (2008). Substance flow analysis of parabens in Denmark complemented with a survey of presence and frequency in various commodities. *Journal of Hazardous Materials*, 156(1-3), 240-259.
- EU (1998). EU's drinking water standards: Council directive 98/83/EC on the quality of water intended for human consumption. Retrieved February 9, 2011, from <http://www.lenntech.com/who-eu-water-standards.htm>
- EU (2002). Council Directive 96/23/EC concerning the performance of analytical methods and the interpretation of results. Retrieved February 9, 2011, from http://ec.europa.eu/food/food/chemicalsafety/residues/control_en.htm
- Farajzadeh, M. A., Djozan, D., & Bakhtiyari, R. F. (2010). Use of a capillary tube for collecting an extraction solvent lighter than water after dispersive liquid-liquid microextraction and its application in the determination of parabens in different samples by gas chromatography--Flame ionization detection. *Talanta*, 81(4-5), 1360-1367.
- Fischer, J., & Jandera, P. (1996). Chromatographic behaviour in reversed-phase high-performance liquid chromatography with micellar and submicellar mobile phases: effects of the organic modifier. *Journal of Chromatography B*, 681(1), 3-19.
- Gaona-Galdos, A. A., García, P. L., Aurora-Prado, M. S., Santoro, M. I. R. M., & Kedor-Hackmann, É. R. M. (2008). Simultaneous determination of econazole nitrate, main impurities and preservatives in cream formulation by high performance liquid chromatography. *Talanta*, 77(2), 673-678.
- García-Jiménez, J. F., Valencia, M. C., & Capitán-Vallvey, L. F. (2007). Simultaneous determination of antioxidants, preservatives and sweetener additives in food and cosmetics by flow injection analysis coupled to a monolithic column. *Analytica Chimica Acta*, 594(2), 226-233.
- González-Mariño, I., Quintana, J. B., Rodríguez, I., & Cela, R. (2009). Simultaneous determination of parabens, triclosan and triclocarban in water by liquid

- chromatography/electrospray ionisation tandem mass spectrometry. *Rapid Communications in Mass Spectrometry*, 23(12), 1756-1766.
- Grabitech Solutions AB (n.d.). The simplex optimization methods. Retrieved November 23, 2010, from <http://www.grabitech.com/algorithm.htm>
- Hamoudová, R., Pospíšilová, M., Kavalířová, A., Solich, P., & Šícha, J. (2006). Separation and determination of clotrimazole, methylparaben and propylparaben in pharmaceutical preparation by micellar electrokinetic chromatography. *Journal of Pharmaceutical and Biomedical Analysis*, 40(1), 215-219.
- Han, F., He, Y.-Z., & Yu, C.-Z. (2008). On-line pretreatment and determination of parabens in cosmetic products by combination of flow injection analysis, solid-phase extraction and micellar electrokinetic chromatography. *Talanta*, 74(5), 1371-1377.
- Han, Y., Jia, X., Liu, X., Duan, T., & Chen, H. (2010). DLLME Combined with GC-MS for the Determination of Methylparaben, Ethylparaben, Propylparaben and Butylparaben in Beverage Samples. *Chromatographia*, 72(3), 351-355.
- Harvey, P. W., & Everett, D. J. (2004). Significance of the detection of esters of p-hydroxybenzoic acid (parabens) in human breast tumours. *Journal of Applied Toxicology*, 24, 1-4.
- Haumann, I., Boden, J., Mainka, A., & Jegle, U. (2000). Simultaneous determination of inorganic anions and cations by capillary electrophoresis with indirect UV detection. *Journal of Chromatography A*, 895(1-2), 269-277.
- He, M., Phillips, J., Wang, P., Xiang, Y., Qi, Y., & Sun, H. (2008). Profiling and characterization of polyphenol polymers from cinnamon using an ion trap mass spectrometer. Retrieved February 20, 2011, from <http://chromatographyonline.findanalytichem.com/lcgc/App+Notes+Medical/Profiling-and-Characterization-of-Polyphenol-Polym/ArticleStandard/Article/detail/503135?contextCategoryId=36341&ref=25>
- He, S., Zhao, Y., Zhu, Z., Liu, H., Li, M., Shao, Y., et al. (2006). Comparative study for the analysis of parabens by micellar electrokinetic capillary chromatography with and without large-volume sample stacking technique. *Talanta*, 69(1), 166-171.

- Henriksen, T., Juhler, R. K., Svensmark, B., & Cech, N. B. (2005). The relative influences of acidity and polarity on responsiveness of small organic molecules to analysis with negative ion electrospray ionization mass spectrometry (ESI-MS). *Journal of the American Society for Mass Spectrometry*, 16(4), 446-455.
- Huang, H.-Y., Lai, Y.-C., Chiu, C.-W., & Yeh, J.-M. (2003). Comparing micellar electrokinetic chromatography and microemulsion electrokinetic chromatography for the analysis of preservatives in pharmaceutical and cosmetic products. *Journal of Chromatography A*, 993(1-2), 153-164.
- Jackson, P. E., Gokhale, S., Streib, T., Rohrer, J. S., & Pohl, C. A. (2000). Improved method for the determination of trace perchlorate in ground and drinking waters by ion chromatography. *Journal of Chromatography A*, 888(1-2), 151-158.
- Jackson, P. E., Laikhtman, M., & Rohrer, J. S. (1999). Determination of trace level perchlorate in drinking water and ground water by ion chromatography. *Journal of Chromatography A*, 850(1-2), 131-135.
- James-Walke, N. L., Williams, H. L., Taylor, D. A., & McMillen, B. A. (2006). The effect of oral consumption of perchlorate, alone and in combination with ethanol, on plasma thyroid hormone and brain catecholamine concentrations in the rat. *Basic & Clinical Pharmacology & Toxicology*, 99, 340-345.
- Jandera, P., & Fischer, J. (1996). Chromatographic behaviour in reversed-phase high-performance liquid chromatography with micellar and submicellar mobile phases. *Journal of Chromatography A*, 728(1-2), 279-298.
- Jinno, K., Watanabe, H., & Kiso, Y. (2001). Fibrous stationary phase in capillary electrochromatography. *Journal of Biochemical and Biophysical Methods*, 48(3), 209-218.
- Khaledi, M. G. (1997). Micelles as separation media in high-performance liquid chromatography and high-performance capillary electrophoresis: overview and perspective. *Journal of Chromatography A*, 780(1), 3-40.
- Khandpur, R. S. (2006). *Analytical instruments* (2nd ed.). New Delhi: Tata McGraw-Hill Publishing Company Limited.

- Kojima, H., Inagaki, M., Tomita, T., Watanabe, T., & Uchida, S. (2010). Improved separation and characterization of lipopolysaccharide related compounds by reverse phase ion pairing-HPLC/electrospray ionization-quadrupole-mass spectrometry (RPIP-HPLC/ESI-Q-MS). *Journal of Chromatography B*, 878(3-4), 442-448.
- Kulikov, A., & Verushkin, A. (2008). Simultaneous Determination of Paracetamol, Caffeine, Guaifenesin and Preservatives in Syrups by Micellar LC. *Chromatographia*, 67(5), 347-355.
- Kyriacos, D. (n.d.). PVC chemistry and production. Retrieved March 29, 2011, from <http://www.gem-chem.net/pvcprod.html>
- Labat, L., Kummer, E., Dallet, P., & Dubost, J. P. (2000). Comparison of high-performance liquid chromatography and capillary zone electrophoresis for the determination of parabens in a cosmetic product. *Journal of Pharmaceutical and Biomedical Analysis*, 23(4), 763-769.
- Labbozzetta, S., Valvo, L., Bertocchi, P., Alimonti, S., Gaudiano, M., & Manna, L. (2009). Focused microwave-assisted extraction and LC determination of ketoprofen in the presence of preservatives in a pharmaceutical cream formulation. *Chromatographia*, 69(3), 365-368.
- Lambertson, C. (n.d.). Simulation addresses band-broadening in HPLC systems. Retrieved March 29, 2011, from http://www.comsol.com/stories/waters_corp_hplc_systems/full/
- Lee, M.-R., Lin, C.-Y., Li, Z.-G., & Tsai, T.-F. (2006). Simultaneous analysis of antioxidants and preservatives in cosmetics by supercritical fluid extraction combined with liquid chromatography-mass spectrometry. *Journal of Chromatography A*, 1120(1-2), 244-251.
- Li, Y., & George, E. J. (2005). Analysis of perchlorate in water by reversed-phase LC/ESI-MS/MS using an internal standard technique. *Analytical Chemistry*, 77(14), 4453-4458.
- Li, Y., & George, E. J. (2006). Reversed-phase liquid chromatography/electrospray ionization tandem mass spectrometry for analysis of perchlorate in water. *Journal of Chromatography A*, 1133(1-2), 215-220.

- Loginova, L. P., Samokhina, L. V., Boichenko, A. P., & Kulikov, A. U. (2006). Micellar liquid chromatography retention model based on mass-action concept of micelle formation. *Journal of Chromatography A*, 1104(1-2), 190-197.
- Magnuson, M. L., Urbansky, E. T., & Kelty, C. A. (2000). Determination of perchlorate at trace levels in drinking water by ion-pair extraction with electrospray ionization mass spectrometry. *Analytical Chemistry*, 72(1), 25-29.
- Mahuzier, P. E., Altria, K. D., & Clark, B. J. (2001). Selective and quantitative analysis of 4-hydroxybenzoate preservatives by microemulsion electrokinetic chromatography. *Journal of Chromatography A*, 924(1-2), 465-470.
- Manuela, M., Simona, S., Irina, R., Ioan, T., & Vasile, O. (2010). Development and application of ultra performance liquid chromatography method to the quantification of the biotransformation of methyl paraben in *eisenia foetida*. *Chinese Journal of Chemistry*, 28(2), 320-324.
- Márquez-Sillero, I., Aguilera-Herrador, E., Cárdenas, S., & Valcárcel, M. (2010). Determination of parabens in cosmetic products using multi-walled carbon nanotubes as solid phase extraction sorbent and corona-charged aerosol detection system. *Journal of Chromatography A*, 1217(1), 1-6.
- Martinelango, P. K., Anderson, J. L., Dasgupta, P. K., Armstrong, D. W., Al-Horr, R. S., & Slingsby, R. W. (2005). Gas-phase ion association provides increased selectivity and sensitivity for measuring perchlorate by mass spectrometry. *Analytical Chemistry*, 77(15), 4829-4835.
- Martinelango, P. K., Gümüş, G., & Dasgupta, P. K. (2006). Matrix interference free determination of perchlorate in urine by ion association-ion chromatography-mass spectrometry. *Analytica Chimica Acta*, 567(1), 79-86.
- Martinelango, P. K., Tian, K., & Dasgupta, P. K. (2006). Perchlorate in seawater: Bioconcentration of iodide and perchlorate by various seaweed species. *Analytica Chimica Acta*, 567(1), 100-107.

- McMahon, G. (2007). *Analytical instrumentation: A guide to laboratory, portable and miniaturized instruments* (1st ed.). West Sussex: John Wiley & Sons, Ltd.
- Melo, L. P., & Queiroz, M. E. C. (2010). Simultaneous analysis of parabens in cosmetic products by stir bar sorptive extraction and liquid chromatography. *Journal of Separation Science*, 33(12), 1849-1855.
- Memon, N., Bhanger, M. I., & Khuhawer, M. Y. (2005). Determination of preservatives in cosmetics and food samples by micellar liquid chromatography. *Journal of Separation Science*, 28(7), 635-638.
- Nabar, G. M., & Ramachandran, C. R. (1959). Quantitative determination of perchlorate ion in solution. *Analytical Chemistry* 31(2), 263-265.
- NC State University (1976). Water resource characterization DSS-Phosphorus. Retrieved February 9, 2011, from <http://www.water.ncsu.edu/watershedss/info/phos.htm>
- Niessen, W. M. A. (1999). *Liquid chromatography-Mass spectrometry* (2nd ed.). New York: Marcel Dekker, Inc.
- Niessen, W. M. A. (2006). *Liquid Chromatography-Mass Spectrometry* (3rd ed.). Boca Raton: Taylor and Francis Group, LLC.
- Nishi, H. (1997). Pharmaceutical applications of micelles in chromatography and electrophoresis. *Journal of Chromatography A*, 780(1-2), 243-264.
- Noguera-Orti, J. F., Villanueva-Camanas, R. M., & Ramis-Ramos, G. (1999). Determination of parabens in cosmetics without previous extraction by micellar liquid chromatography. *Journal of Chromatographic Science*, 37(3), 83-87.
- Núñez, L., Tadeo, J. L., García-Valcárcel, A. I., & Turiel, E. (2008). Determination of parabens in environmental solid samples by ultrasonic-assisted extraction and liquid chromatography with triple quadrupole mass spectrometry. *Journal of Chromatography A*, 1214(1-2), 178-182.
- Núñez, L., Turiel, E., Martín-Esteban, A., & Tadeo, J. L. (2010). Molecularly imprinted polymer for the extraction of parabens from environmental solid samples prior to their determination by high performance liquid chromatography-ultraviolet detection. *Talanta*, 80(5), 1782-1788.

- Papoušková, B., Bednář, P., Fryšová, I., Stýskala, J., Hlaváč, J., Barták, P., et al. (2007). Mass spectrometric study of selected precursors and degradation products of chemical warfare agents. *Journal of Mass Spectrometry*, 42(12), 1550-1561.
- Pedrouzo, M., Borrull, F., Marcé, R. M., & Pocurull, E. (2009). Ultra-high-performance liquid chromatography-tandem mass spectrometry for determining the presence of eleven personal care products in surface and wastewaters. *Journal of Chromatography A*, 1216(42), 6994-7000.
- Ramírez, N., Marcé, R. M., & Borrull, F. (2010). Development of a thermal desorption-gas chromatography-mass spectrometry method for determining personal care products in air. *Journal of Chromatography A*, 1217(26), 4430-4438.
- Rouessac, F., & Rouessac, A. (1994). *Chemical analysis: Modern instrumentation methods and techniques* (2nd ed.). West Sussex: John Wiley & Sons, Ltd.
- Routledge, E. J., Parker, J., Odum, J., Ashby, J., & Sumpter, J. P. (1998). Some alkyl hydroxy benzoate preservatives (parabens) are estrogenic. *Toxicology and Applied Pharmacology*, 153(1), 12-19.
- Rozenski, J. (1999). MolE-Molecular mass calculator. Retrieved April 29, 2011, from <http://www.library.med.utah.edu/masspec/mole.htm>
- Ruiz-Ángel, M. J., Carda-Broch, S., Torres-Lapasió, J. R., & García-Álvarez-Coque, M. C. (2009). Retention mechanisms in micellar liquid chromatography. *Journal of Chromatography A*, 1216(10), 1798-1814.
- Saad, B., Bari, M. F., Saleh, M. I., Ahmad, K., & Talib, M. K. M. (2005). Simultaneous determination of preservatives (benzoic acid, sorbic acid, methylparaben and propylparaben) in foodstuffs using high-performance liquid chromatography. *Journal of Chromatography A*, 1073(1-2), 393-397.
- Šafra, J., & Pospíšilová, M. (2008). Separation and determination of ketoprofen, methylparaben and propylparaben in pharmaceutical preparation by micellar electrokinetic chromatography. *Journal of Pharmaceutical and Biomedical Analysis*, 48(2), 452-455.

- Segui, M. J., Lizondo-Sabater, J., Martínez-Máñez, R., Sancenón, F., Soto, J., García-Breijo, E., et al. (2006). An ion-selective electrode for anion perchlorate in thick-film technology. *Sensors*, 6(5), 480-491.
- Shabir, G. A. (2010). Development and validation of a stability-indicating LC method for the determination of domperidone, sorbic acid, and propylparaben in pharmaceutical formulations. *Journal of Liquid Chromatography & Related Technologies*, 33(20), 1802-1813.
- Shabir, G. A. (2010). A new validated HPLC method for the simultaneous determination of 2-phenoxyethanol, methylparaben, ethylparaben and propylparaben in a pharmaceutical gel. *Indian Journal of Pharmaceutical Sciences*, 72(4), 421-425.
- Shanmugam, G., Ramaswamy, B. R., Radhakrishnan, V., & Tao, H. (2010). GC-MS method for the determination of paraben preservatives in the human breast cancerous tissue. *Microchemical Journal*, 96(2), 391-396.
- Shimoaki, T. (n.d.). Surfactant. Retrieved February 28, 2011, from http://www.ach.nitech.ac.jp/~physchem/taga/1tagag_e.html
- Snyder, S. A., Pleus, R. C., Vanderford, B. J., & Holady, J. C. (2006). Perchlorate and chlorate in dietary supplements and flavor enhancing ingredients. *Analytica Chimica Acta*, 567(1), 26-32.
- Snyder, S. A., Vanderford, B. J., & Rexing, D. J. (2005). Trace analysis of bromate, chlorate, iodate, and perchlorate in natural and bottled waters. *Environmental Science & Technology*, 39(12), 4586-4593.
- Soukup-Hein, R. J., Remsburg, J. W., Dasgupta, P. K., & Armstrong, D. W. (2007). A general, positive ion mode ESI-MS approach for the analysis of singly charged inorganic and organic anions using a dicationic reagent. *Analytical Chemistry*, 79(19), 7346-7352.
- Tavares, R. S., Martins, F. C., Oliveira, P. J., Ramalho-Santos, J., & Peixoto, F. P. (2009). Parabens in male infertility--Is there a mitochondrial connection? *Reproductive Toxicology*, 27(1), 1-7.
- Tayama, S., Nakagawa, Y., & Tayama, K. (2008). Genotoxic effects of environmental estrogen-like compounds in CHO-K1 cells. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 649(1-2), 114-125.

- Tian, K., Dasgupta, P. K., & Anderson, T. A. (2003). Determination of trace perchlorate in high-salinity water samples by ion chromatography with on-line preconcentration and preelution. *Analytical Chemistry*, 75(3), 701-706.
- U.S. Environmental Protection Agency (2005). EPA Method 314.1: Determination of perchlorate in drinking water using inline column concentration/matrix elimination ion chromatography with suppressed conductivity detection. Retrieved February 25, 2011, from <http://yosemite.epa.gov/water/owrrccatalog.nsf/e673c95b11602f2385256ae1007279fe/afc152d00cae6b648525707400565c2a!OpenDocument>
- Vidović, S., Stojanović, B., Veljković, J., Pražić-Arsić, L., Roglić, G., & Manojlović, D. (2008). Simultaneous determination of some water-soluble vitamins and preservatives in multivitamin syrup by validated stability-indicating high-performance liquid chromatography method. *Journal of Chromatography A*, 1202(2), 155-162.
- Vlasenko, A. S., Loginova, L. P., & Iwashchenko, E. L. (2009). Dissociation constants and micelle-water partition coefficients of hydroxybenzoic acids and parabens in surfactant micellar solutions. *Journal of Molecular Liquids*, 145(3), 182-187.
- Waichigo, M. M., & Danielson, N. D. (2006). Ethylammonium formate as an organic solvent replacement for ion-pair reversed-phase liquid chromatography. *Journal of Chromatographic Science*, 44(10), 607-614.
- Waldman, P. (2002). Perchlorate runoff flows to water supply of millions: A fuel of cold war defenses now ignites health controversy. Retrieved February 22, 2010, from <http://www.organicconsumers.org/Toxic/wj-perchlorate.cfm>
- Wang, Y., Cao, Y., Fang, C., & Gong, Q. (2010). Electrochemical sensor for parabens based on molecular imprinting polymers with dual-templates. *Analytica Chimica Acta*, 673(2), 145-150.
- Washington University (2004). Molecular representations and tables of common polyatomic ions. Retrieved February 28, 2011, from <http://www.chemistry.wustl.edu/~courses/genchem/Tutorials/Ions/ions.htm>

- Watson, J. T., & Sparkman, O. D. (2007). *Introduction to mass spectrometry: Instrumentation, applications and strategies for data interpretation* (4th ed.). West Sussex: John Wiley & Sons, Ltd.
- Winkler, P., Minter, M., & Willey, J. (2004). Analysis of perchlorate in water and soil by electrospray LC/MS/MS. *Analytical Chemistry*, 76(2), 469-473.
- Woollard, D. C., & Indyk-Harvey, E. (2002). Rapid determination of thiamine, riboflavin, pyridoxine, and niacinamide in infant formulas by liquid chromatography. *Journal of AOAC International*, 85, 945-951.
- Wu, T., Wang, C., Wang, X., & Ma, Q. (2008). Simultaneous determination of 21 preservatives in cosmetics by ultra performance liquid chromatography. *International Journal of Cosmetic Science*, 30, 367-372.
- Yang, T.-J., Tsai, F.-J., Chen, C.-Y., Yang, T. C.-C., & Lee, M.-R. (2010). Determination of additives in cosmetics by supercritical fluid extraction on-line headspace solid-phase microextraction combined with gas chromatography-mass spectrometry. *Analytica Chimica Acta*, 668(2), 188-194.
- Youngvise, N., Liawruangrath, B., & Liawruangrath, S. (2003). Simultaneous micellar LC determination of lidocaine and tolperisone. *Journal of Pharmaceutical and Biomedical Analysis*, 31(4), 629-638.
- Zhang, Q., Lian, M., Liu, L., & Cui, H. (2005). High-performance liquid chromatographic assay of parabens in wash-off cosmetic products and foods using chemiluminescence detection. *Analytica Chimica Acta*, 537(1-2), 31-39.
- Zotou, A., Sakla, I., & Tzanavaras, P. D. (2010). LC-determination of five paraben preservatives in saliva and toothpaste samples using UV detection and a short monolithic column. *Journal of Pharmaceutical and Biomedical Analysis*, 53(3), 785-789.