

รายการอ้างอิง

- Abrami, L. and van der Goot, F.G. (1999). Plasma membrane microdomains act as concentration platforms to facilitate intoxication by aerolysin. *Journal of Cell Biology*, 147, 175-184.
- Abrami, L., Fivaz, M. and van der Goot, F.G. (2000). Adventures of a pore-forming toxin at a target cell surface. *Trends in Microbiology*, 8, 168-172.
- Akiba, T., Abe, Y., Kitada, S., Kasuka, Y., Ito, A., Ichimatsu, T., Katayama, H., Akao, T., Higuchi, K., Misuki, E., Ohba, M., Kanai, R. and Harata, K. (2009). Crystal structure of the parasporin-2 *Bacillus thuringiensis* toxin that recognizes cancer cells. *Journal of Molecular Biology*, 386, 121-133.
- Alouf, J.E. and Popoff, M.R. (2006). *The comprehensive sourcebook of bacterial protein toxins*. USA: Elsevier Ltd.
- Amersham Biosciences. *GST gene fusion system: handbook*.
- Aquino de Muro, M., Mitchell, W.J. and Priest, F.G. (1992). Differentiation of mosquito pathogenic strains of *Bacillus sphaericus* from non-toxic varieties by ribosomal RNA gene restriction pattern. *Journal of General Microbiology*, 138, 1159-1166.
- Baumann, P. and Broadwell, A.H. (1987). Proteolysis in the gut of mosquito larvae results in further activation of the *Bacillus sphaericus* toxin. *Applied and Environmental Microbiology*, 53, 1333-1337.
- Baumann, P., Clark, M.A., Baumann, L. and Broadwell, A.H. (1991). *Bacillus sphaericus* as a mosquito pathogen: Properties of the organism and its toxins. *Microbiological Reviews*, 55, 425-436.
- Bayyareddy, K., Andacht, T.M., Abdullah, M.A. and Adang, M.J. (2009). Proteomic identification of *Bacillus thuringiensis* subsp. *israelensis* toxin Cry4Ba binding protein in midgut membranes from *Aedes (Stegomyia) aegypti* Linnaeus (Diptera, Culicidae) larvae. *Insect Biochemistry and Molecular Biology*, 1-8.

- Beyenbach, K.W. and Wieczorek, H. (2006). The V-type H⁺ ATPase: molecular structure and function, physiological roles and regulation. *The Journal of Experimental Biology*, 209, 577-589.
- Charles, J.F. and Nielsen-LeRoux, C. (1992). Binding of *Bacillus sphaericus* binary toxin to a specific receptor on midgut brush-border membranes from mosquito larvae. *European Journal of Biochemistry*, 210, 585-590.
- Charles, J.F., Nielsen-LeRoux, C. and Delecluse, A. (1996). *Bacillus sphaericus* Toxins: Molecular biology and mode of action. *Annual Review of Entomology*, 41, 451-472.
- Davidson, E.W. (1988). Binding of the *Bacillus sphaericus* (Eubacteriales: Bacillaceae) toxin to midgut cells of mosquito (Diptera: Culicidae) larvae: relationship to host range. *Journal of Medical Entomology*, 25, 151-157.
- Diep, D.B., Nelson, K.L., Raja, S.M., Pleshak, E.N. and Buckley, J.T. (1988b). Glycosylphosphatidylinositol anchors of membrane glycoproteins are binding determinants for the channel-forming toxin aerolysin. *The Journal of Biological chemistry*, 273, 2355-2360.
- Federici, B.A., Park, H.-W., Bideshi, D.K., Wirth, M.C. and Johnson, J.J. (2003). Recombinant bacteria for mosquito control. *The Journal of Experimental Biology*, 206, 3877-3885.
- GE Healthcare. (2007). *Restriction site for pGEX-4T-1*. USA: General Electric Company.
- Gomez, I., Pardo-Lopez, L., Munoz-Garay, C., Fernandez, L.E., Perez, C., Sanchez, J., Soberon, M. and Bravo, A. (2007). Role of receptor interaction in the mode of action of insecticidal Cry and Cyt toxins produced by *Bacillus thuringiensis*. *Peptide*, 28, 169-173.
- Ito, A., Sasaguri, Y., Kitada, S., Kusaka, Y., Kuwano, K., Matsutomi, K. et al. (2004). A *Bacillus thuringiensis* crystal protein with selective cytotoxic action to human cells. *The Journal of Biological chemistry*, 279, 21282-21286.
- Janda, J.M. and Abbott, S.L. (1998). Evolving concepts regarding the genus *Aeromonas*: an expanding panorama of species, disease presentations,

- and unanswered questions. *Journal of Clinical and Infectious Diseases*, 27, 332-344.
- Katayama, H., Yokota, H., Akao, T., Nakamura, O., Ohba, M., Mekada, E. and Mizuki, E. (2005). Parasporin-1, a novel cytotoxic protein to human cells from non-insecticidal parasporal inclusions of *Bacillus thuringiensis*. *The Journal of Biological chemistry*, 137, 17-25.
- Kirov, S.M. (1997). *Aeromonas* and *Plesiomonas* species. In Doyle, M., Beuchat, L. and Montville, T. (Eds). (1997). *Food Microbiology: Fundamental and Frontiers*, 265-287.
- Kitada, S., Abe, Y., Shimada, H., Kusaka, Y., Matsuo, Y., Katayama, H., Okumura, S., Akao, T., Mizuki, E., Kuge, O., Sasaguri, Y., Ohba, M. and Ito, A. (2006). Cytocidal actions of parasporin-2, an anti-tumor crystal toxin from *Bacillus thuringiensis*. *The Journal of Biological chemistry*, 281, 26350-26360.
- Lee, D.W., Akao, T., Yamashita, S., Katayama, H., Maeda, M., Saito, H. et al. (2000). Noninsecticidal parasporal proteins of a *Bacillus thuringiensis* serova *shandongiensis* isolate exhibit a preferential cytotoxicity against human leukemic T. cell. *Biochemical and Biophysical Research Communications*. 272, 218-223.
- Liu, J.W., Porter, A.G., Wee, B.Y. and Thanabalu, T. (1996). New gene from nine *Bacillus sphaericus* strains encoding highly conserved 35.8-kilodalton mosquitocidal toxins. *Applied and Environmental Microbiology*, 62, 2174-2176.
- Melton-Witt, J.A., Bentsen, L.M. and Tweten, R.K. (2006). Identification of functional domains of *Clostridium septicum* alpha toxin. *Biochemistry*, 45, 14347-14354.
- Nelson, K.L., Raja, S.M. and Buckley, J.T. (1997). The glycosylphosphatidylinositol-anchored surface glycoprotein Thy-1 is a receptor for the channel-forming toxin aerolysin. *The Journal of Biological Chemistry*, 272, 12170-12174.
- Nishiwaki, H., Nakashima, K., Ishida, C., Kawamura, T. and Matsuda, K. (2007). Cloning, functional characterization, and mode of action of a novel insecticidal pore-

- forming toxin, Sphaericolysin, produced by *Bacillus sphaericus*. *Applied and Environmental Microbiology*, 73, 3404-3411.
- Parker, M.W., Buckley, J.T., Postma, J.P.M., Tucker, A.D., Leonard, K. and Pattus, F. et al. (1994). Structure of the *Aeromonas* toxin proaerolysin in its water-soluble and membrane-channel states. *Nature*, 367, 292-295.
- Payne, D. and Oysten, P. (1997). The *Clostridium perfringens* ϵ -toxin. In: *The Clostridia: Molecular Biology and Pathogenesis* (eds. Rood, J.I., McClane, B.A., Songer, J.G. and Titball, R.W.), Academic Press.
- Popova-Butler, A. and Dean, D.H. (2009). Proteomic analysis of the mosquito *Aedes aegypti* midgut brush border membrane vesicles. *Journal of Insect Physiology*, 55, 264-272.
- Regis, L.R., Silva-Filha, M.H., Nielsen-LeRoux, C. and Charles, J.F. (2001). Bacteriological larvicides of dipteran disease vectors. *TRENDS in Parasitology*, 17, 377-380.
- Sakurai, J. and Nagahama, M. (1985). Tryptophan content of *Clostridium perfringens* epsilon toxin. *Infection and Immunity*, 47, 260-263.
- Sakurai, J. and Nagahama, M. (1987a). Histidine residues in *Clostridium perfringens* epsilon toxin. *FEMS Microbiology. Lett.*, 41, 317-319.
- Sakurai, J. and Nagahama, M. (1987b). The inactivation of *Clostridium perfringens* epsilon toxin by treatment with tetranitromethane and N-acetylimidazole. *Toxicon*, 25, 279-284.
- Sambrook, J. and Russell, D.W. (2001). *Molecular cloning: a laboratory manual* (3rd ed.). New York: Cold Spring Harbor Laboratory Press.
- Silva-Filha, M.H., Nielsen-LeRoux, C. and Charles, J.F. (1997). Binding kinetics of *Bacillus sphaericus* binary toxin to midgut brush-border membranes of *Anopheles* and *Culex* sp. mosquito larvae. *European Journal of Biochemistry*, 247, 754-761.
- Silva-Filha, M.H., Nielsen-LeRoux, C. and Charles, J.F. (1999). Identification of the receptor for *Bacillus sphaericus* crystal toxin in the brush border membrane

- of the mosquito *Culex pipiens* (Diptera: Culicidae). *Insect Biochemistry and Molecular Biology*, 29, 711-721.
- Thanabalu, T., Hindley, J., Jackson-Yap, J. and Berry, C. (1991). Cloning, Sequencing, and expression of a gene encoding a 100-kilodalton mosquitocidal toxin from *Bacillus sphaericus* SSII-1. *Journal of Bacteriology*, 173, 2776-2785.
- Thanabalu, T., Berry, C. and Hindley, J. (1993). Cytotoxicity and ADP-ribosylating activity of the mosquitocidal toxin from *Bacillus sphaericus* SSII-1: possible roles of the 27- and 70-kilodalton peptides. *Journal of Bacteriology*, 175, 2314-2320.
- Thanabalu, T. and Porter, A.G. (1996). A *Bacillus sphaericus* gene encoding a novel type of mosquitocidal toxin of 31.8 kDa. *Gene*, 170, 85-89.
- Treiber, N., Reinert, D.J., Carpusca, I., Aktories, K. and Schulz, G.E. (2008). Structure and mode of action of a mosquitocidal holotoxin. *Journal of Molecular Biology*, 381, 150-159.
- Yamashita, S., Akao, T., Mizuki, E., Saitoh, H., Higuchi, K., Park, Y.S. et al. (2000). Characterization of the anti-cancer-cell parasporal proteins of a *Bacillus thuringiensis* isolate. *Canadian Journal of Microbiology*, 46, 913-919.