

## สารบัญ

	หน้า
บทคัดย่อ	1
Executive Summary	2
บทที่ 1 บทนำ	4
บทที่ 2 วัสดุและวิธีการ	17
บทที่ 3 ผลการทดลองและวิจารณ์ผลการทดลอง	24
บทที่ 4 สรุปผลการศึกษา	47
เอกสารอ้างอิง	48
Output ที่ได้จากโครงการ	55
ภาคผนวก	56

## สารบัญตาราง

	หน้า
<b>Table 1.1</b> สมบัติและองค์ประกอบทางเคมีของซีโอไลต์ ไครน็อฟติโลไลต์ ภูเขาไฟ และสเม็กไทต์	16
<b>Table 2.1</b> Summary of substrate formulations	19
<b>Table 3.1</b> The time period of different phases of oyster mushroom cultivation, yield and percentage BE of <i>Pleurotus ostreatus</i> grown on substrates non- supplemented and supplemented with gypsum, pumice sulfate and pumice.	26
<b>Table 3.2</b> Size classes (small, medium, large) expressed as a percentage of the total mushroom production per treatment. Quantiles 33.3% (6.70 g) and 66.6% (12.28 g) were used to delimit mushroom size classes (small, medium, large), mushroom cap diameter (cm) and firmness (kgf) of <i>Pleurotus ostreatus</i> grown on substrates non-supplemented and supplemented with gypsum, pumice sulfate and pumice.	30
<b>Table 3.3</b> The calcium content (mg/g dry wt.) and silicon content (mg/g dry wt.) of <i>Pleurotus ostreatus</i> grown on substrates non-supplemented and supplemented with gypsum, pumice sulfate and pumice.	31
<b>Table 3.4</b> The calcium content (mg/g dry wt.) and silicon content (mg/g dry wt.) in mycelia of <i>Pleurotus ostreatus</i> grown on WPDY medium	39
<b>Table 3.5</b> Percentage of weight gained and size loss and firmness (kgf) of canned mushrooms. Mushrooms were cultivated in substrates supplemented with gypsum, pumice sulfate and pumice.	41
<b>Table 3.6</b> Percentage of weight loss and size loss and firmness (kgf) of frozen mushrooms. Mushrooms were cultivated in substrates supplemented with gypsum, pumice sulfate and pumice.	42

## สารบัญรูป

## หน้า

- Figure 3.1** a.) A histogram showing the distribution of individual (2,848) mushroom weights. Quantiles 33.3% (6.70 g) and 66.6% (12.28 g) were used to delimit mushroom size classes (small, medium, large) b.) A histogram showing the distribution of individual (2,848) mushroom cap diameter showing normal distribution curve with mean of 7.3 cm and standard deviation of 2.26 cm. The caps with diameter of  $7.3 \pm 2.26$  cm were used to produce canned mushrooms. 29
- Figure 3.2** The percentage drip loss and solid content of the caps and stalks of oyster mushrooms grown on substrates non-supplemented and supplemented with gypsum, pumice sulfate and pumice. Means followed by the same letter in the same column in the same set of supplement are not significantly different according to the Duncan test ( $P \geq 0.05$ ) 40
- Figure 3.3A** Scanning electron micrographs showing a cross section of stalks of *P. ostreatus* grown on substrates non-supplemented (a) supplemented with gypsum (b) pumice sulfate (c) and pumice (d) after thermal treatment. (Canned mushroom; 500x magnification) For each sample, the image was the representative image from 3 different samples randomly collected from at least 10 canned mushrooms. Only mushroom caps with diameter of 7 cm and mushroom stalks with 1 cm thickness were chosen. 43
- Figure 3.3B** Scanning electron micrographs showing a cross section of stalks of *P. ostreatus* grown on substrates non-supplemented (a) supplemented with gypsum (b) pumice sulfate (c) and pumice (d) after thermal treatment. (Canned mushroom; 100x magnification) 44
- Figure 3.3C** Scanning electron micrographs showing a cross section of caps of *P. ostreatus* grown on substrates non-supplemented (a) supplemented with gypsum (b) pumice sulfate (c) and pumice (d) after thermal treatment. (Canned mushroom; 500x magnification) For each sample, the image was the representative image from 3 different samples randomly collected from at least 10 canned mushrooms. Only mushroom caps with diameter of 7 cm and mushroom stalks with 1 cm thickness were chosen. 45
- Figure 3.3D** Scanning electron micrographs showing a cross section of caps of *P. ostreatus* grown on substrates non-supplemented (a) supplemented with gypsum (b) pumice sulfate (c) and pumice (d) after thermal treatment. (Canned mushroom; 100x magnification) 46