

LIST OF CONTENTS

Chapter	Pages
I INTRODUCTION	1
Rationale for the study.....	1
Review methods for Thomas-Fermi model	2
Review methods for Adomian Polynomials	2
Purpose of the Study	4
Significance of the Study	4
Scope of the Study	4
 II THOMAS-FERMI MODEL	 5
Fermi statistics	5
The monatomic ideal gas	7
Thomas-Fermi equation.....	10
 III RESEARCH METHODOLOGY	 13
Modified Adomian Decomposition Method.....	13
The Padé approximants.....	15
Padé approximants for extends the truncated seriessolution	15
Padé approximants and Boundary Value Problems.....	17
Electron distribution for Mercury atom	20
 IV ORDER-BY-ORDER EXTRACTION TECHNIQUE	 21
 V RESULTS AND DISCUSSION.....	 25
Adomian Polynomials for the nonlinear term of Thomas - Fermi equation via OBOET	25
Thomas-Fermi equation solved by MADM via OBOET	27
The initial slope for each Padé approximants order [n/n].....	29
The solutions of Thomas-Fermi equation for each order [n/n].....	31
The results of electron distribution for Mercury atom.....	33

LIST OF CONTENTS (CONT.)

Chapter	Pages
VI CONCLUSIONS	35
REFERENCES	36
APPENDIX	39
BIOGRAPHY	74

LIST OF TABLES

Table	Page
1 Several values of initial slope $B = y'(0)$ calculated by Padé approximants.....	31
2 The solution of the Thomas-Fermi equation due to the Differential analyzer in form of $(x, y(x))$ and $(\eta, y(\eta))$	43

LIST OF FIGURES

Figures	Pages
1 Non-interacting quantum gas of fermions periodic boundary conditions in a large box $V = L^3$	7
2 The coordinates in number of state are n_x, n_y, n_z . The number of state available to a particle with a state whose magnitude is between n and $n + dn$ is proportional to the volume of a spherical shell in state space of radius n and thickness dn	8
3 Fermi-Dirac distribution at $T = 0$	9
4 Flowchart for computing A_n in case $F(y) = y^{3/2}$, via OBOET	24
5 Comparison of solutions of Thomas-Fermi equation obtained by the Differential Analyzer [1], MADM via OBOET, MADM via OBOET incorporated Padé approximants orders [4/4], [7/7], and [10/10]	31
6 Electron distributions obtained by the Differential Analyzer [1,3], MADM via OBOET, MADM via OBOET incorporated Padé approximants orders [4/4], [7/7], and [10/10]	33
7 Comparison of the electron distributions of an atom of mercury; using MADM via OBOET incorporated Padé approximants orders [10/10], the Differential Analyzer [1], and Hartree [3]	34