

## LIST OF TABLES

<b>Table</b>	<b>Page</b>
<b>LITERATURE REVIEW</b>	
1 Combinatorial antibody diversity in humans	35
2 Genes and gene products of the $\phi$ 1 bacteriophage	48
<b>RESULTS</b>	
3 Demographic data of the 50 blood donors and the concentrations and purity of their total RNA preparations.	104
4 Details of the 10 buffy coat samples obtained from Thai blood bank, Thai Red Cross, Bangkok, Thailand.	107
5 Degenerate and non-degenerate forward primer sequences that were designed from VBASE with <i>Sfi</i> I restriction endonuclease site (underlined alphabets) at 5'-end of the primers and were used for amplification of human <i>VH</i> by PCR.	114
6 Degenerate and non-degenerate reverse primer sequences that were designed from VBASE with polynucleotide linkers (underlined alphabets) at 5'-end of the primers and were used for amplification of human <i>VH</i> by PCR.	116
7 Degenerate and non-degenerate forward primer sequences were designed from VBASE with polynucleotide linker (underlined alphabets) at 5'-end of primers and were used to amplify human <i>V<math>\kappa</math></i> in PCR.	125
8 Degenerate and non-degenerate reverse primer sequences were designed from VBASE with <i>Not</i> I restriction endonuclease site (underlined alphabets) at 5'-end of primers and were used to amplify human <i>V<math>\kappa</math></i> in PCR.	126

## LIST OF TABLES (Cont.)

<b>Table</b>	<b>Page</b>
9 Orthologous proteins of the database matched with the peptide sequences generated from tryptic digestion of proteins in circles no. 1-7, 11, 16-22 and 24 of <i>N. kaouthia</i> venom components separated by 2DE at non-linear pH 3-10.	147
10 Numbers of alive and dead mice at 24 hours after intra-peritoneal injection with various amounts of crude <i>N. kaouthia</i> venom	158
11 Number of alive and dead mice at 24 hours after intra-muscular injection of various amounts of crude <i>N. kaouthia</i> .	159
12 Numbers of alive and dead mice at 24 hours after intra-peritoneal injection with 2.18 µg of column-fractionated crude <i>N. kaouthia</i> venom, P1 to P11 (0.5x LD50 by means of intra-peritoneal route).	160
13 Orthologs of the venom proteins in P3, P4, P5, and P8 identified by 2D-LC/MS-MS after orthologous search	163
14 Results of phage bio-panning using P1, P3, P4, P5, and P8 as antigens	168
15 Survival rates of different groups of mice after receiving crude <i>N. kaouthia</i> venom pre-incubated with horse anti-venom or HuScFv from different phage clones compared to the group that received only venom.	220
16 Survival rates of different groups of mice after receiving crude <i>N. kaouthia</i> venom only or the venom pre-incubated with horse anti-venom or HuScFv from different phage clones.	221