

LIST OF FIGURES

Figure	Page
LITERATURE REVIEW	
1 Morphology of <i>Naja kaouthia</i> or Thai cobra	5
2 Reported cases of poisonous snake bites per 100,000 populations by year, Thailand, 1997-2006.	7
3 Reported cases of poisonous snake bites by months, Thailand, 2002-2006.	8
4 Proportion (%) of snake bites by snake types in Thailand, 2005.	9
5 Proportion (%) of snake bites by snake types in Thailand, 2006.	10
6 The three dimensional structure of five representatives of the three-finger toxins.	13
7 Two dimensional structure of long α -neurotoxin of <i>N. naja siamensis</i> (presently, <i>N. kaouthia</i>).	14
8 Three dimensional structure and secondary structure of long neurotoxin of <i>N. naja siamensis</i> (gi 229777 pdb 1CTX).	15
9 Three dimensional model of $\alpha 7$ receptor- α -cobratoxin (α -Cbtx) complex.	17
10 The three dimensional structure of Cbtx-AChBP complex.	18
11 Three-dimensional structure of short α -neurotoxin (A) and the protein interaction between short α -neurotoxin (NTII) and α/γ -subunits of the muscular type-nicotinic acetylcholine receptor (nAChR) (B) of <i>Torpedo californica</i> .	20
12 (A) Amino acid residues involved in binding with α/γ interface of the muscular type-nicotinic acetylcholine receptor (nAChR). (B) Multiple alignments of three short α -neurotoxins are shown.	21
13 Mechanism of phospholipase A2 in hydrolysis of phospholipids (phosphatidylserine).	24

LIST OF FIGURES (Cont.)

Figure	Page
14 The protein structure of (human IgG1) immunoglobulin molecules.	30
15 Multigene organization and V(D)J recombination of heavy chain of immunoglobulin gene segments.	31
16 Molecular mechanism of immunoglobulin gene rearrangement.	33
17 Sequential steps in mouse monoclonal antibody production.	39
18 Chimeric antibody and CDR grafting of humanized-MAb.	42
19 Gene organization of filamentous bacteriophage M13 genome.	46
20 Schematic diagram of a filamentous bacteriophage M13.	47
21 Structure of intact immunoglobulin G (IgG) molecule, a minibody, and monomeric antibody fragments: Fab (antibody fragment), Fv (variable fragment), and scFv (single chain variable fragment).	51
22 Schematic diagram of a ScFv-displaying filamentous bacteriophage M13.	54
23 Schematic diagrams for constructing ScFv repertoire by using phage display technology.	55
24 Schematic process of phage bio-panning against immobilized antigen	58
MATERIALS AND METHODS	
25 Diagram showing binding of forward (P1r) and reverse (Pr2) primers to the immunoglobulin DNA template and positions of oligonucleotide linker and endonuclease restriction sites for PCR amplification of <i>VH</i> sequences.	68
26 Diagram showing binding of forward (Pr3) and reverse (Pr4) primers to the immunoglobulin DNA template and positions of oligonucleotide linkers and endonuclease restriction sites for PCR amplification of <i>Vκ</i> sequences.	69

LIST OF FIGURES (Cont.)

Figure	Page
27 Sequential steps in SOE-PCR for constructing <i>huscFv</i> sequences.	72
28 Equipment of the Mini Trans-Blot® Electrophoretic Transfer Cell	92
29 The gel sandwich preparation for trans-blotting.	93
RESULTS	
30 PBMC were prepared by using VACUTAINER® CPT™ Cell Preparation tubes containing sodium citrate (A) and Ficoll-Paque (B).	109
31 Agarose gel electrophoretic patterns of total RNA.	112
32 PCR amplicons of human β -actin gene segments (~250 bp) using cDNA of the volunteers and buffy coat samples as templates.	113
33 PCR products of VH-coding DNA sequences of some immunoglobulin subclasses (<i>VH1a, 1b, 1c, 1d, 2a, 2b, and 3a</i> family-specific forward primers were combined with <i>JH1245</i> reverse primer).	117
34 PCR products of VH-coding DNA sequences of some immunoglobulin subclasses (<i>VH3b, 3c, 4a, 4b, 5a, 6a, and 7a</i> family-specific forward primers were combined with <i>JH1245</i> reverse primer).	118
35 PCR products of VH-coding DNA sequences of some immunoglobulin subclasses (<i>VH1b, 1c, 2a, 2b, 4a, 4b, 6a, and 7a</i> family-specific forward primers were combined with <i>JH1245</i> reverse primers).	119
36 PCR products of VH-coding DNA sequences of some immunoglobulin (Ig) subclasses (<i>VH1a, 1b, 1c, 1d, 2a, 2b, and 3a</i> family-specific forward primers were combined with <i>JH3</i> reverse primer).	121

LIST OF FIGURES (Cont.)

Figure	Page
37 PCR products of VH-coding DNA sequences of some immunoglobulin (Ig) subclasses (<i>VH3b, 3c, 4a, 4b, 5a, 6a, and 7a</i> family-specific forward primers were combined with <i>JH3</i> reverse primer).	122
38 PCR products of VH-coding DNA sequences of some immunoglobulin (Ig) subclasses (<i>VH1a, 1b, 1c, 1d, 2a, 2b, and 3a</i> family-specific forward primers were combined with <i>JH6</i> reverse primer).	123
39 PCR products of VH-coding DNA sequences of some immunoglobulin (Ig) subclasses (<i>VH3b, 3c, 4a, 4b, 5a, 6a, and 7a</i> family-specific forward primers were combined with <i>JH6</i> reverse primer).	124
40 PCR products of human V κ -coding DNA sequences (~400 bp).	128
41 <i>VH</i> and <i>Vκ</i> DNA pools were analyzed on 1.5% agarose gel.	129
42 Optimization of annealing temperature for polynucleotide linkers.	131
43 Optimization of annealing temperature for outer primers.	132
44 Influence of outer primer adding step for generation of <i>huscFv</i> .	133
45 Large scale preparation of <i>huscFv</i> .	134
46 Quantification of <i>huscFv</i> .	135
47 The transformation efficiency of electro-competent TG1 <i>E. coli</i> .	137
48 TG1 <i>E. coli</i> transformants on SOBAG agar plate.	138
49 Phage titer of human antibody phage display library.	139
50 <i>huscFv</i> amplification of 20 TG1 <i>E. coli</i> clones derived from the transformation.	141
51 Patterns of <i>MvaI</i> RFLP of <i>huscFv</i> sequences.	142
52 SDS-PAGE-patterns of <i>N. kaouthia</i> venom.	145

LIST OF FIGURES (Cont.)

Figure	Page
53 Proteomic pattern of <i>N. kaouthia</i> venom.	146
54 Locations of tryptic peptides generated from proteins in circles no. 1 and no. 3-6 within cobra venom factor (accession no. I51018).	152
55 Locations of tryptic peptides generated from proteins in circles no. 16-21 within phospholipase A2 (accession no. P15445, P25498, PSNJ2K and 1OWSA).	153
56 Elution profiles of crude <i>N. kaouthia</i> venom from ion exchange column chromatography.	155
57 SDS-PAGE patterns of proteins in peaks numbered 1 to 11.	156
58 Mass spectrum of the P1 proteins (Figure 57A) which was identified as cobra venom factor by MALDI-TOF (accession no. 54035743).	161
59 Amplicons of <i>huscFv</i> of the 11 randomly selected transformed HB2151 <i>E. coli</i> colonies from a total of 252 transformed HB2151 <i>E. coli</i> colonies infected with phages that bound to P1 from the single round of bio-panning.	169
60 Results of HuScFv expressions and Western blot analysis of 10 <i>huscFv</i> -pCANTAB5E-transformed HB2151 <i>E. coli</i> colonies derived from single round of bio-panning with P1 which predominantly contained the cobra venom factor.	170
61 Results of <i>huscFv</i> amplification of the 38 transformed HB2151 <i>E. coli</i> clones selected from single round of bio-panning with P3 which principally contained phospholipase A2.	171
62 Results of HuScFv expression and Western blot analysis of the 20 <i>huscFv</i> -pCANTAB5E-transformed HB2151 <i>E. coli</i> colonies derived from single round of bio-panning with P3 .	172

LIST OF FIGURES (Cont.)

Figure	Page
63 Results of <i>huscFv</i> amplification of the 22 transformed HB2151 <i>E. coli</i> clones selected from single round of bio-panning with P4 which principally contained natrin toxin.	173
64 Results of HuScFv expression and Western blot analysis of the 12 <i>huscFv</i> -pCANTAB5E-transformed HB2151 <i>E. coli</i> colonies derived from single round of bio-panning with P4 .	174
65 Results of <i>huscFv</i> amplification of the 18 transformed HB2151 <i>E. coli</i> colonies derived from the third round of bio-panning with P4 (principally contained natrin).	175
66 Results of HuScFv expression and Western blot analysis of the 7 <i>huscFv</i> -pCANTAB5E-transformed HB2151 <i>E. coli</i> colonies derived from the third round of bio-panning with P4 .	176
67 Results of the <i>huscFv</i> amplification of the 20 transformed HB2151 <i>E. coli</i> colonies derived from the single round of bio-panning with P5 which principally contained phospholipase A2.	177
68 Results of the HuScFv expression and Western blot analysis of the 19 <i>huscFv</i> -pCANTAB5E-transformed HB2151 <i>E. coli</i> colonies derived from single round of bio-panning with P5 which principally contained phospholipase A2.	178
69 Results of the <i>huscFv</i> amplification of the 26 transformed HB2151 <i>E. coli</i> colonies derived from the first single round of bio-panning with P8 (principally contained long neurotoxin).	179
70 Results of the HuScFv expression and Western blot analysis of the 21 <i>huscFv</i> -pCANTAB5E-transformed HB2151 <i>E. coli</i> colonies derived from single round of bio-panning with P8 .	180
71 Results of the <i>huscFv</i> amplification of the 20 transformed HB2151 <i>E. coli</i> colonies derived from the second single round of bio-panning with P8 .	181

LIST OF FIGURES (Cont.)

Figure	Page
72 Results of the HuScFv expression and Western blot analysis of the 12 <i>huscFv</i> -pCANTAB5E-transformed HB2151 <i>E. coli</i> colonies derived from single round of bio-panning (2 nd time) with P8 .	182
73 Results of the <i>huscFv</i> amplification of the 31 transformed HB2151 <i>E. coli</i> colonies derived from the third round of bio-panning with P8 .	183
74 Results of the HuScFv expression and Western blot analysis of the 12 <i>huscFv</i> -pCANTAB5E-transformed HB2151 <i>E. coli</i> colonies derived from third round of bio-panning with P8 .	184
75 Results of the indirect ELISA of the HuScFv of the <i>E. coli</i> clones derived from single round of bio-panning with P1 (cobra venom factor).	188
76 Results of the indirect ELISA of HuScFv of <i>E. coli</i> clones derived from the single round bio-panning with P3 (phospholipase A2).	189
77 Result of the indirect ELISA of HuScFv of <i>E. coli</i> clones derived from single round of bio-panning with P4 (natrin).	190
78 Results of the indirect ELISA of HuScFv of the <i>E. coli</i> clones derived from the third rounds bio-panning with P4 (natrin).	191
79 Results of the indirect ELISA of HuScFv derived from the single round bio-panning with P5 (phospholipase A2).	192
80 Results of the indirect ELISA of HuScFv derived from the first single round of bio-panning with P8 (neurotoxin).	193
81 Indirect ELISA of HuScFv derived from the second time of single round of bio-panning with P8 (neurotoxin).	194
82 Indirect ELISA of HuScFv derived from the third rounds bio-panning with P8 (neurotoxin).	195

LIST OF FIGURES (Cont.)

Figure	Page
83 Dot-ELISA results of soluble HuScFv prepared from <i>E. coli</i> HB2151 colonies derived from bio-panning with cobra venom factor in P1 of Thai cobra venom, namely clones P1/1, P1/2, P1/3, P1/5, and P1/9.	198
84 Dot-ELISA results of soluble HuScFv prepared from <i>E. coli</i> HB2151 colonies derived from bio-panning with phospholipase in P3 of Thai cobra venom namely P3/2, P3/5, P3/6, P3/7, and P3/18 and natrin in P4 of Thai cobra venom, namely P4/8 and P4/21.	199
85 Dot-ELISA results of soluble HuScFv prepared from <i>E. coli</i> HB2151 derived from bio-panning with phospholipase A2 in P5 of Thai cobra venom, namely P5/1, P5/2, P5/3, P5/4, P5/5, P5/6, P5/7, P5/9, P5/10, and P5/11.	200
86 Dot-ELISA results of soluble HuScFv prepared from <i>E. coli</i> HB2151 derived from bio-panning with neurotoxin in P8 of Thai cobra venom, namely P8/0/1, P8/9/1, P8/19/1, P8/7/2, P8/10/2, P8/22/3, and P8/31/3.	201
87 Western blot analysis of HuScFv prepared from HB2151 <i>E. coli</i> clones derived from bio-panning with cobra venom factor (P1).	204
88 Western blot analysis of HuScFv prepared from HB2151 <i>E. coli</i> clones derived from bio-panning with phospholipases A2 (P3) and natrin (P4).	205
89 Western blot analysis of soluble HuScFv prepared from HB2151 <i>E. coli</i> clones that were derived from bio-panning with phospholipases A2 (P5).	206
90 Western blot analysis of HuScFv prepared from <i>E. coli</i> HB2151 clones that were derived from bio-panning with neurotoxin (P8).	207

LIST OF FIGURES (Cont.)

Figure	Page
91 Optical densities of the proteins of <i>huscFv</i> -transformed <i>E. coli</i> lysate after passing through the affinity anti-E-Tag column.	209
92 Dot-ELISA for detecting HuScFv in whole cell lysates of HB2151 <i>E. coli</i> and protein fractions after purification.	210
93 SDS-PAGE (A) and Western blot analysis (B) of proteins in various fractions from the affinity anti-E-Tag column	211
94 Patterns of <i>MvaI</i> RFLP of <i>huscFv</i> sequences	213
95 Appearance of plaques on an LB agar plate containing IPTG and X-gal	214
96 Eighteen representatives of M13 phage genome (Lanes 1-18) extracted from the amplified phage particles in LB medium by using phenol/chloroform and ethanol precipitation. Each M13 phage genome was analyzed on 0.8% agarose gel.	215
97 The representative DNA sequencing result of M13 phage genome selected from the third round of bio-panning with HuScFv that was prepared from HB2151 <i>E. coli</i> clone P8/0/1 plaque number 1.	216
98 Pairwise alignment of the 12-mer mimotope “NPVEWFMSTVNT” and long neurotoxin of <i>N. naja siamensis</i> (α -neurotoxin) by using CLUSTALW 2.0.5 multiple sequence alignment.	217
99 Three dimensional structure of long neurotoxin of <i>N. naja siamensis</i> (gi 229777 pdb 1CTX).	218