

References

Books

- Pathummanon. (1998). *Medical Epidemiology*. Bangkok: A PHRCG Publishers.
- Perrin. (1993). *Isokinetic exercise and assessment*. Unpublished manuscript, Virginia.
- Waddell. (1998). *The back pain revolution*. Edinburgh: Churchill Livingstone.

Articles

- Adams, Mannion, & Dolan. (1999). Personal risk factors for first-time low back pain. *Spine*, 24(23), 2497-505.
- Alaranta, Luoto, Heliovaara, & Hurri. (1995). Static back endurance and the risk of low-back pain. *Clin Biomech (Bristol, Avon)*, 10(6), 323-24.
- Andersson. (1999). Epidemiological features of chronic low-back pain. *Lancet*, 354(9178), 581-5.
- Arab, Salavati, Ebrahimi, & Ebrahim Mousavi. (2007). Sensitivity, specificity and predictive value of the clinical trunk muscle endurance tests in low back pain. *Clin Rehabil*, 21(7), 640-7.
- Baecke, Burema, & Frijters. (1982). A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr*, 36(5), 936-42.
- Barnekow-Bergkvist, Hedberg, Janlert, & Jansson. (1998). Determinants of self-reported neck-shoulder and low back symptoms in a general population. *Spine*, 23(2), 235-43.
- Bejia, Abid, Ben Salem, Letaief, Younes, Touzi, et al. (2005). Low back pain in a cohort of 622 Tunisian schoolchildren and adolescents: an epidemiological study. *Eur Spine J*, 14(4), 331-6.
- Bombardier. (2000). Outcome assessments in the evaluation of treatment of spinal disorders: summary and general recommendations. *Spine*, 25(24), 3100-3.
- Burton, Balague, Cardon, Eriksen, Henrotin, Lahad, et al. (2005). How to prevent low back pain.[erratum appears in Best Pract Res Clin Rheumatol. 2005

- Dec;19(6):1095 Note: Balague, F [added]; Cardon, G [added]; Eriksen, HR [added]; Henrotin, Y [added]; Lahad, A [added]; Leclerc, A [added]; Muller, G [added]; van der Beek, AJ [added]]. *Best Pract Res Clin Rheumatol*, 19(4), 541-55.
- Burton, Clarke, McClune, & Tillotson. (1996). The natural history of low back pain in adolescents. *Spine*, 21(20), 2323-8.
- Cairns, Foster, & Wright. (2006). Randomized controlled trial of specific spinal stabilization exercises and conventional physiotherapy for recurrent low back pain. *Spine*, 31(19), E670-81.
- Cassidy, Carroll, & Cote. (1998). The Saskatchewan health and back pain survey. The prevalence of low back pain and related disability in Saskatchewan adults. *Spine*, 23(17), 1860-6; discussion 67.
- Cassidy, Cote, Carroll, & Kristman. (2005). Incidence and course of low back pain episodes in the general population. *Spine*, 30(24), 2817-23.
- Coorevits, Danneels, Cambier, Ramon, & Vanderstraeten. (2008). Assessment of the validity of the Biering-Sorensen test for measuring back muscle fatigue based on EMG median frequency characteristics of back and hip muscles. *J Electromyogr Kinesiol*.
- Croft, Macfarlane, Papageorgiou, Thomas, & Silman. (1998). Outcome of low back pain in general practice: a prospective study. *BMJ*, 316(7141), 1356-9.
- Daltroy, Cats-Baril, Katz, Fossel, & Liang. (1996). The North American spine society lumbar spine outcome assessment Instrument: reliability and validity tests. *Spine*, 21(6), 741-9.
- Danforth, Jensen, Kopelman, Lefebvre, & Reeder. (2001). Dose-response issues concerning physical activity and health: an evidence-based symposium. *Med Sci Sports Exerc*, 33(6), S351-S58.
- Davidson, & Keating. (2002). A comparison of five low back disability questionnaires: reliability and responsiveness. *Phys Ther*, 82(1), 8-24.
- de Vet, Heymans, Dunn, Pope, van der Beek, Macfarlane, et al. (2002). Episodes of low back pain: a proposal for uniform definitions to be used in research. *Spine*, 27(21), 2409-16.

- Denis, Shannon, Wessel, Stratford, & Weller. (2007). Association of low back pain, impairment, disability & work limitations in nurses. *J Occup Rehabil*, 17(2), 213-26.
- Deyo, Battie, Beurskens, Bombardier, Croft, Koes, et al. (1998). Outcome measures for low back pain research. A proposal for standardized use. *Spine*, 23(18), 2003-13.
- Dionne, Dunn, Croft, Nachemson, Buchbinder, Walker, et al. (2008). A consensus approach toward the standardization of back pain definitions for use in prevalence studies. *Spine*, 33(1), 95-103.
- Elders, & Burdorf. (2004). Prevalence, incidence, and recurrence of low back pain in scaffolders during a 3-year follow-up study. *Spine*, 29(6), E101-6.
- Eriksen, Natvig, & Bruusgaard. (1999). Smoking, heavy physical work and low back pain: a four-year prospective study. *Occup Med (Lond)*, 49(3), 155-60.
- Fairbank, Couper, Davies, & O'Brien. (1980). The Oswestry low back pain disability questionnaire. *Physiotherapy*, 66(8), 271-3.
- George. (2002). The six-month incidence of clinically significant low back pain in the Saskatchewan adult population. *Spine (Phila Pa 1976)*, 27(16), 1778-82.
- Ghaffari, Alipour, Farshad, Yensen, & Vingard. (2006). Incidence and recurrence of disabling low back pain and neck-shoulder pain. *Spine*, 31(21), 2500-6.
- Griffith, Hogg-Johnson, Cole, Krause, Hayden, Burdorf, et al. (2007). Low-back pain definitions in occupational studies were categorized for a meta-analysis using Delphi consensus methods. *J Clin Epidemiol*, 60(6), 625-33.
- Gundewall, Liljeqvist, & Hansson. (1993). Primary Prevention of Back Symptoms and Absence from Work: A Prospective Randomized Study Among Hospital Employees. *Spine*, 18(5), 587-94.
- Hamberg-van Reenen, Ariens, Blatter, van Mechelen, & Bongers. (2007). A systematic review of the relation between physical capacity and future low back and neck/shoulder pain. *Pain*, 130(1-2), 93-107.
- Hellsing, & Bryngelsson. (2000). Predictors of musculoskeletal pain in men: A twenty-year follow-up from examination at enlistment. *Spine (Phila Pa 1976)*, 25(23), 3080-6.

- Hestbaek, Korsholm, Leboeuf-Yde, & Kyvik. (2008). Does socioeconomic status in adolescence predict low back pain in adulthood? A repeated cross-sectional study of 4,771 Danish adolescents. *Eur Spine J*, 17(12), 1727-34.
- Hodges, & Richardson. (1996). Inefficient muscular stabilization of the lumbar spine associated with low back pain. A motor control evaluation of transversus abdominis. *Spine*, 21(22), 2640-50.
- Iwai, Nakazato, Irie, Fujimoto, & Nakajima. (2004). Trunk muscle strength and disability level of low back pain in collegiate wrestlers. *Med Sci Sports Exerc*, 36(8), 1296-300.
- Janwantanakul, Pensri, Jiamjarasrangsi, & Sinsongsook. (2009). Associations between prevalence of self-reported musculoskeletal symptoms of the spine and biopsychosocial factors among office workers. *J Occup Health*, 51(2), 114-22.
- Jarvik, & Deyo. (2002). Diagnostic evaluation of low back pain with emphasis on imaging. *Ann Intern Med*, 137(7), 586-97.
- Jorgensen. (1997). Human trunk extensor muscles: Physiology and ergonomics. *Acta Physiologica Scandinavica Supplementum*, 160(637), 1-58.
- Josephson, Hagberg, & Hjelm. (1996). Self-reported physical exertion in geriatric care. A risk indicator for low back symptoms? *Spine*, 21(23), 2781-5.
- Karahan, Kav, Abbasoglu, & Dogan. (2009). Low back pain: prevalence and associated risk factors among hospital staff. *Journal of Advanced Nursing*, 65(3), 516-24.
- Karatas, Gogus, & Meray. (2002). Reliability of isokinetic trunk muscle strength measurement. *Am J Phys Med Rehabil*, 81(2), 79-85.
- Keller, Johansen, Hellesnes, & Brox. (1999). Predictors of isokinetic back muscle strength in patients with low back pain. *Spine*, 24(3), 275-80.
- Kopec, & Sayre. (2004). Work-related psychosocial factors and chronic pain: a prospective cohort study in Canadian workers. *J Occup Environ Med*, 46(12), 1263-71.
- Krismer, & van Tulder. (2007). Strategies for prevention and management of musculoskeletal conditions. Low back pain (non-specific). *Best Pract Res Clin Rheumatol*, 21(1), 77-91.

- Kujala, Taimela, Viljanen, Jutila, Viitasalo, Videman, et al. (1996). Physical loading and performance as predictors of back pain in healthy adults. A 5-year prospective study. *Eur J Appl Physiol Occup Physiol*, 73(5), 452-8.
- Latimer, Maher, Refshauge, & Colaco. (1999). The reliability and validity of the Biering-Sorensen test in asymptomatic subjects and subjects reporting current or previous nonspecific low back pain. *Spine*, 24(20), 2085-9; discussion 90.
- Leboeuf-Yde. (2000). Alcohol and low-back pain: a systematic literature review. *J Manipulative Physiol Ther*, 23(5), 343-6.
- Leboeuf-Yde. (2000). Body Weight and Low Back Pain: A Systematic Literature Review of 56 Journal Articles Reporting on 65 Epidemiologic Studies. *Spine*, 25(2), 226.
- Leboeuf-Yde, & Kyvik. (1998). At what age does low back pain become a common problem? A study of 29,424 individuals aged 12-41 years. *Spine (Phila Pa 1976)*, 23(2), 228-34.
- Leboeuf-Yde, Kyvik, & Bruun. (1998). Low back pain and lifestyle. Part I: Smoking. Information from a population-based sample of 29,424 twins. *Spine*, 23(20), 2207-13; discussion 14.
- Leboeuf-Yde, Kyvik, & Bruun. (1999). Low back pain and lifestyle. Part II--Obesity. Information from a population-based sample of 29,424 twin subjects. *Spine*, 24(8), 779-83; discussion 83-4.
- Leclaire, Blier, Fortin, & Proulx. (1997). A cross-sectional study comparing the Oswestry and Roland-Morris Functional Disability scales in two populations of patients with low back pain of different levels of severity. *Spine*, 22(1), 68-71.
- Lee, Hoshino, Nakamura, Kariya, Saita, & Ito. (1999). Trunk muscle weakness as a risk factor for low back pain. A 5-year prospective study. *Spine*, 24(1), 54-7.
- Linton. (2000). A review of psychological risk factors in back and neck pain. *Spine*, 25(9), 1148-56.
- Mannion, Dolan, & Adams. (1996). Psychological Questionnaires: Do "Abnormal" Scores Precede or Follow First-time Low Back Pain? [Miscellaneous Article]. *Spine J*, 15(21(22)), 2603-11.

- Masset, Piette, & Malchaire. (1998). Relation between functional characteristics of the trunk and the occurrence of low back pain. Associated risk factors. *Spine*, 23(3), 359-65.
- Mazloun, Nozad, & Kumashiro. (2006). Occupational low back pain among workers in some small-sized factories in Ardabil, Iran. *Ind Health*, 44(1), 135-9.
- Mills, Taunton, & Mills. (2005). The effect of a 10-week training regimen on lumbo-pelvic stability and athletic performance in female athletes: A randomized-controlled trial. *Phys Ther Sport*, 6(2), 60-66.
- Miranda, Viikari-Juntura, Martikainen, Takala, & Riihimaki. (2002). Individual factors, occupational loading, and physical exercise as predictors of sciatic pain. *Spine*, 27(10), 1102-9.
- Mirtz, & Greene. (2005). Is obesity a risk factor for low back pain? An example of using the evidence to answer a clinical question. *Chiropr Osteopat*, 13(1), 2.
- Moreau, Green, Johnson, & Moreau. (2001). Isometric back extension endurance tests: a review of the literature. *J Manipulative Physiol Ther*, 24(2), 110-22.
- Mortimer, Wiktorin, Pernold, Svensson, & Vingard. (2001). Sports activities, body weight and smoking in relation to low-back pain: a population-based case-referent study. *Scand J Med Sci Sports*, 11(3), 178-84.
- Omokhodion, Umar, & Ogunnowo. (2000). Prevalence of low back pain among staff in a rural hospital in Nigeria. *Occupational Medicine (Oxford)*, 50(2), 107-10.
- Ozguler, Leclerc, Landre, Pietri-Taleb, & Niedhammer. (2000). Individual and occupational determinants of low back pain according to various definitions of low back pain. *J Epidemiol Community Health*, 54(3), 215-20.
- Panjabi. (2003). Clinical spinal instability and low back pain. *J Electromyogr Kinesiol*, 13(4), 371-9.
- Pensri, Foster, Srisuk, Baxter, & McDonough. (2005). Physiotherapy management of low back pain in Thailand: a study of practice. *Physiother Res Int*, 10(4), 201-12.
- Power, Frank, Hertzman, Schierhout, & Li. (2001). Predictors of low back pain onset in a prospective British study. *Am J Public Health*, 91(10), 1671-8.

- Proper, Koning, van der Beek, Hildebrandt, Bosscher, & van Mechelen. (2003). The effectiveness of worksite physical activity programs on physical activity, physical fitness, and health. *Clin J Sport Med, 13*(2), 106-17.
- Rissanen, Heliovaara, Alaranta, Taimela, Malkia, Knekt, et al. (2002). Does good trunk extensor performance protect against back-related work disability? *J Rehabil Med, 34*(2), 62-6.
- Rives, & Douglass. (2004). Evaluation and treatment of low back pain in family practice. *J Am Board Fam Pract, 17*, 823-31.
- Sakulsriprasert, Vachalathiti, Vongsirinavarat, & Kantasorn. (2006). Cross-cultural adaptation of modified Oswestry Low Back Pain Disability Questionnaire to Thai and its reliability. *J Med Assoc Thai, 89*(10), 1694-701.
- Sanjaroensuttikul. (2007). The Oswestry low back pain disability questionnaire (version 1.0) Thai version. *J Med Assoc Thai, 90*(7), 1417-22.
- Schneider, Randoll, & Buchner. (2006). Why Do Women Have Back Pain More Than Men?: A Representative Prevalence Study in the Federal Republic of Germany. [Article]. *The Clinical Journal of Pain, 22*(8), 738-47.
- Shiri, Karppinen, Leino-Arjas, Solovieva, & Viikari-Juntura. (2010). The association between obesity and low back pain: a meta-analysis. *Am J Epidemiol, 171*(2), 135-54.
- Shiri, Solovieva, Husgafvel-Pursiainen, Taimela, Saarikoski, Huupponen, et al. (2008). The association between obesity and the prevalence of low back pain in young adults: the Cardiovascular Risk in Young Finns Study. *Am J Epidemiol, 167*(9), 1110-9.
- Soukup, Glomsrod, Lonn, Bo, & Larsen. (1999). The Effect of a Mensendieck Exercise Program as Secondary Prophylaxis for Recurrent Low Back Pain: A Randomized, Controlled Trial With 12-Month Follow-up. *Spine J, 24*(15), 1585.
- Soukup, Lonn, Glomsrod, Bo, & Larsen. (2001). Exercises and education as secondary prevention for recurrent low back pain. *Physiother Res Int, 6*(1), 27-39.

- Spyropoulos, Papathanasiou, Georgoudis, Chronopoulos, Koutis, & Koumoutsou. (2007). Prevalence of low back pain in greek public office workers. *Pain Physician, 10*(5), 651-9.
- Staal, Hlobil, van Tulder, Koke, Smid, & van Mechelen. (2002). Return-to-work interventions for low back pain: a descriptive review of contents and concepts of working mechanisms. *Sports Medicine, 32*(4), 251-67.
- Staal, Rainville, Fritz, van Mechelen, & Pransky. (2005). Physical exercise interventions to improve disability and return to work in low back pain: current insights and opportunities for improvement. *J Occup Rehabil, 15*(4), 491-505.
- Stevenson, Weber, Smith, Dumas, & Albert. (2001). A longitudinal study of the development of low back pain in an industrial population. *Spine, 26*(12), 1370-7.
- Stewart, Latimer, & Jamieson. (2003). Back extensor muscle endurance test scores in coal miners in Australia. *J Occup Rehabil, 13*(2), 79-89.
- Storheim, Bo, Pederstad, & Jahnsen. (2002). Intra-tester reproducibility of pressure biofeedback in measurement of transversus abdominis function. *Physiother Res Int, 7*(4), 239-49.
- Stroyer, & Jensen. (2008). The role of physical fitness as risk indicator of increased low back pain intensity among people working with physically and mentally disabled persons: a 30-month prospective study. *Spine, 33*(5), 546-54.
- Takala, & Viikari-Juntura. (2000). Do functional tests predict low back pain? *Spine, 25*(16), 2126-32.
- Thomas, Silman, & Croft. (1999). Predicting who develops chronic low back pain in primary care: a prospective study. *BMJ, 318*, 1662-7.
- Thongjunjua. (2007). Effects of lumbar stabilization exercises on exercise level attained in healthy subjects. *Thai J Phys Ther, 1*, 98.
- Underwood. (2000). Exercise and the prevention of back pain disability. *Br J Sports Med, 34*(1), 5.
- van Poppel, Hooftman, & Koes. (2004). An update of a systematic review of controlled clinical trials on the primary prevention of back pain at the workplace. *Occupational Medicine, 54*(5), 345-52.

- van Tulder, Koes, & Bombardier. (2002). Low back pain. *Best Pract Res Clin Rheumatol*, 16(5), 761-75.
- van Tulder, Koes, & Bouter. (1995). A cost-of-illness study of back pain in The Netherlands. *Pain*, 62(2), 233-40.
- Vuori. (2001). Dose-response of physical activity and low back pain, osteoarthritis, and osteoporosis. *Med Sci Sports Exerc*, 33(6 Suppl), S551-86; discussion 609-10.
- Walker, Muller, & Grant. (2004). Low back pain in Australian adults: prevalence and associated disability. *Journal of Manipulative & Physiological Therapeutics*, 27(4), 238-44.
- Waxman, Tennant, & Helliwell. (2000). A prospective follow-up study of low back pain in the community. *Spine*, 25(16), 2085-90.
- Webster, & Snook. (1994). The cost of 1989 workers' compensation low back pain claims. *Spine*, 19(10), 1111-5.
- WHO. (2004). Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet*, 363, 157-63.
- Yip. (2001). A study of work stress, patient handling activities and the risk of low back pain among nurses in Hong Kong. *J Adv Nurs*, 36(6), 794-804.

Other Materials

- Annual Report Thammasat University 2007*. (2007). Bangkok: Thammasat University.
- Bernard. (1997, July 10997). Musculoskeletal disorders (MSDs) and workplace factors. *A critical review of epidemiologic evidence for work-related musculoskeletal disorders of the neck, upper extremity, and low back*. Retrieved 27/09/05, 2005, from <http://www.cdc.gov/niosh/ergoscui1.html>
- Intakochasarn. (2006). *Back pain of labourers in the context of labour process: A case study of ready-made clothes factory in Bangkok vicinity*. Mahidol University, Bangkok.

- Piriyaprasarth. (1999). *Comparison of concentric isokinetic of trunk muscles performance between female subjects with and without low back pain.* Mahidol, Bangkok.
- Sengkisiri. (1998). *A study on musculo-skeletal disorders in industrial: risk factors and its improvement.* Bangkok.
- Social Security Statistics 2006: Statistics of Workmen's Compensation Fund (Covered by WCF).* (2007). Bangkok: The War Veterans Organization of Thailand.