

บรรณานุกรม

บทความ

ปริญญาธิ์ สมฤทธิ. (มปป). “ข้าวลูกผสม: สถานภาพและแนวทางการวิจัยและพัฒนาของประเทศไทย”.
กรมวิชาการเกษตร.

Yuan L. (2004). “Hybrid Rice Technology for Food Security in the World”. China National
Hybrid Rice R & D Center. FAO RICE CONFERENCE 04/CRS.21.

เอกสารอื่นๆ

กระทรวงเกษตรแห่งสหรัฐอเมริกา.

<http://www.fao.org/es/ess/yearbook/vol_1_1/site_en.asp?page=production >

ปริญญาธิ์ สมฤทธิ. 2546. การประชุมเพื่อนำเสนอผลการศึกษา เรื่อง “แนวทางการวิจัยและพัฒนาข้าว
ลูกผสมของประเทศไทย”.วันที่ 20 กุมภาพันธ์ 2546. อาคารไบโอเทค อุทยานวิทยาศาสตร์
ประเทศไทย, ประเทศไทย.

Yuan L.P. (1987). “Strategy conception of hybrid rice breeding”. Hybrid Rice 1: 1–3.

Books

กิตติพัฒน์ อุโฆษกิจ. 2006. พันธุศาสตร์. กรุงเทพฯ: สำนักพิมพ์มหาวิทยาลัยธรรมศาสตร์.

ประพาส วีระแพทย์. 1983. ความรู้เรื่องข้าว. กรุงเทพฯ: ไทยวัฒนาพานิช.

Crawford, G.W. and C. Shen. 1998. The Origins of Rice Agriculture: Recent Progress in East
Asia. Antiquity 72:858–866.

Maruyama K., Araki H. and Kato H. (1991). "Thermosensitive genic male sterility induced by irradiation". Rice genetics II. International rice research institute, Manila, the Philippines, pp 227-235.

Virmani, S .S., Sun, Z.X., Mou, T.M., Jauhar Ali, A. and Mao, C.X. (2003). "Two-lines hybrid rice breeding manual". International Rice Research Institute, P.O. Box 933, Manila, Philippines.

Articles

Akagi H., Nakamura A., Yokozeki-Misono Y., Inagaki A., Takahashi H. (2004). "Positional cloning of the rice *Rf-1* gene, a restorer of BT-type cytoplasmic male sterility that encodes a mitochondria-targeting PPR protein". Theor Appl Genet 108: 1449-1457.

Allard R. W. (1956). "Formulas and tables to facilitate the calculation of recombination values in heredity". Hilgardia 24: 235-278.

Atsuma S., Takeo N. and Yasuo K. (1992). "Sterility in rice induced by chemical treatment". Bio sci. Biotech. Biochem. 56: 1619 -1622.

Borkakati, R.P. & Virmani S.S. (1996). "Genetics of thermosensitive genic male sterility in rice". Euphytica 88: 1-7.

Chen P.L., Chen B., Chen J.I. and Ho L.K.(1986). "Arsenic content and castration effect in reaction to male gametocide no.2 in rice plants". J. South China Agric. Univ. 7:22-27.

Dong NV, Subudhi PK, Luong PN, Quang VD, Quy TD. (2000). "Molecular mapping of a rice gene conditioning thermosensitive genic male sterility using AFLP, RFLP, and SSR techniques". Theor Appl Genet 100: 727-734.

- Hamzeiy, H., Vahdati-Mashhadian, N., Edwards, H.J. and Goldfarb, P.S. (2002). "Mutation analysis of the human *CYP3A4* gene 5' regulatory region: population screening using non-radioactive SSCP". Mutat. Res. 500: 103-110.
- Hanson, W. D. (1959). "Early generation analysis of lengths of heterozygous chromosome segments around a locus held heterozygous with backcrossing or selfing". Genetics 44, 833-837.
- Hoisington D, Khairallah M, Reeves J, Skovmand B. (1999). "Plant genetic resources: What can they contribute toward increased crop productivity?". Proc Natl Acad Sci USA 96: 5937-5943.
- Komori T, Ohta S, Murai N, Takakura Y, Kuraya T. (2004). "Map-based cloning of a fertility restorer gene, Rf-1 in rice (*Oryza sativa* L.)". Plant J. 37: 315-325.
- Latha R., Thiyagarajan K. and Senthilvel S. (2004). "Genetics, fertility behaviour and molecular marker analysis of a new TGMS line, TS6, in rice". Plant Breeding 123: 235-240.
- Lee D.S., Chen L.J., Suh H.S. (2005). "Genetic characterization and fine mapping of a novel thermo-sensitive genic male-sterile gene *tms6* in rice (*Oryza sativa* L.)". Theor Appl Genet 111:1271-1277.
- Liu N., Shan Y., Wang F.P., Xu C.G., Peng KM., Li X.H., Zhang Q. (2001). "Identification of an 85-kb DNA fragment containing *pms1*, a locus for photoperiod-sensitive genic male sterility in rice". Mol Genet Genomics 266: 271-275.
- Lopez M.T. and Virmani S.S. (2000). "Development of TGMS lines for developing two-line rice hybrids for the tropics". Euphytica 114: 211-215.

- Lukuwitz W, Grillmor CS, Scheible WR. (2000). "Positional cloning in Arabidopsis. Why it feels good to have a genomic initiative working for you". Plant Physiol 123: 795-806.
- Naveira, H. and Barbadilla, A. (1992). "The theoretical distribution of lengths of intact chromosome segments around a locus held heterozygous with backcrossing in a diploid species". Genetics 130, 205–209.
- Orti G., Hare M.P., Avise J.C. (1997). "Detection and isolation of nuclear haplotypes by PCR-SSCP". Molecular Ecology, 6, 575–580.
- Pitnjam K., Chakhonkaen S., Toojinda T., Muangprom A. (2008). "Identification of deletion in *tms2* and development of gene-based markers for selection". Planta. 228(5):813-22.
- Reddy O., Siddiq E., Ali N., Hussain A.J., Nimmakayala P. and Ramasamy P. (2000). "Genetic analysis of temperature-sensitive male sterility in rice". Theor Appl Genet 100: 794-801.
- Rongbai li, Pandey M.P., Sharma P. (2005). "Inheritance of thermosensitive genic male sterility in rice (*Oryza sativa* L.)". Curr science 88 (11):1809-1815.
- Small M.P., Gosling E.M. (2000). "Species relationships in population structure of *Littorina saxatilis* Olivi and *L. tenebrosa* Montagu in Ireland using single-strand conformational polymorphisms (SSCPs) of cytochrome *b* fragments". Molecular Ecology 9: 39–52.
- Srivastava H. (2004). "Mitochondria-nuclear interaction in male sterility an heterosis for productivity enhancement in crop plants". Journ Sust Agric 22(4):63-78.
- Stam, P. and Zeven, A. C. (1981). "The theoretical proportion of the donor genome in near-isogenic lines of self-fertilizers bred by backcrossing". Euphytica 30: 227–238.

- Subudhi P.K., Borkakati R.K., Virmani S.S., Huang N. (1997). "Molecular mapping of a *thermosensitive genetic male-sterility* gene in rice using bulked segregant analysis". Genome 40: 188–194.
- Sun C.Q., Jaing T.B., Fu Y.C. and Wang Z.K. (2002). "Indica-Japonica differentiation of paddy rice and its relationship with heterosis". Plant Breeding 121: 330-337.
- Sunnucks, P., Wilson, A.C.C., Beheregaray, L.B., Zenger, K., French, J., Taylor, A.C. (2000). "SSCP is not so difficult: The application and utility of single-stranded conformation polymorphism in evolutionary biology and molecular ecology". Molecular Ecology 9 (11): 1699-1710.
- Viraktamath, B.C. and Virmani, S.S. (2001). "Expression of thermosensitive genic male sterility in rice under varying temperature situations". Euphytica 122(1): 137-143.
- Wang B., Xu W.W., Wang J.Z., Wu W., Zheng H.G., Yang Z.Y., Ray J.D., Nguyen H.T. (1995). "Tagging and mapping the thermosensitive genic male-sterile gene in rice (*Oryza sativa* L.) with molecular markers". Theor Appl Genet 91: 1111–1114.
- Wang Y.G., Xing Q.H., Deng Q.Y., Liang F.S., Yuan L.P., Weng M.L., Wang B. (2003). "Fine mapping of the rice thermo-sensitive genic male-sterile gene *tms5*". Theor Appl Genet 107: 917–921.
- Yamaguchi Y., Ikeda R., Hirasawa H., Minami M., Ujihara A. (1997). "Linkage analysis of the thermo-sensitive genic male sterility gene *tms2* in rice (*Oryza sativa* L.)". Breed Sci 47: 371–373.
- Yang, Q., Liang, C., Zhuang, W., Li, J., Deng, H., Deng, Q., Wang, B. (2007). "Characterization and identification of the candidate gene of rice thermo-sensitive genic male sterile gene *tms5* by mapping". Planta 225 (2): 321-330.