

## Air Quality Assessment Using Interpolation Technique

Awkash Kumar<sup>a</sup>, Rashmi S. Patil<sup>a</sup>, Anil Kumar Dikshit<sup>a</sup> and Rakesh Kumar<sup>b</sup>

<sup>a</sup> Centre for Environmental Science and Engineering, Indian Institute of Technology Bombay,  
Mumbai-400076, India

<sup>b</sup> National Environmental Engineering Research Institute Mumbai Zone, Mumbai-400018, India

---

### Abstract

Air pollution is increasing rapidly in almost all cities around the world due to increase in population. Mumbai city in India is one of the mega cities where air quality is deteriorating at a very rapid rate. Air quality monitoring stations have been installed in the city to regulate air pollution control strategies to reduce the air pollution level. In this paper, air quality assessment has been carried out over the sample region using interpolation techniques. The technique Inverse Distance Weighting (IDW) of Geographical Information System (GIS) has been used to perform interpolation with the help of concentration data on air quality at three locations of Mumbai for the year 2008. The classification was done for the spatial and temporal variation in air quality levels for Mumbai region. The seasonal and annual variations of air quality levels for SO<sub>2</sub>, NO<sub>x</sub> and SPM (Suspended Particulate Matter) have been focused in this study. Results show that SPM concentration always exceeded the permissible limit of National Ambient Air Quality Standard. Also, seasonal trends of pollutant SPM was low in monsoon due rain fall. The finding of this study will help to formulate control strategies for rational management of air pollution and can be used for many other regions.

**Keywords:** air quality; interpolation technique; ArcGIS; IDW

---

### 1. Introduction

Urbanization is a major trend all over the world mainly due to rapid growth in population. More than half the population of the world now lives in urban areas. This may be attributed to the continuous increase in global population (Kim, 2005; Huff and Angeles, 2011). Urbanization is affecting air quality at global, regional and local scales (Contini *et al.*, 2012; Duh *et al.*, 2008; Nemitz *et al.*, 2008; Järvi *et al.*, 2009). Urban air quality is a complex subject with different socio-economic aspects in different parts of the world and even within a specific region. The population growth is directly correlated to many environmental quality parameters like water and air quality (Alcamo *et al.*, 2002; Bollen *et al.*, 2009; 2010; Kan *et al.*, 2012; Kanada *et al.*, 2013; Yennawar, 1978). Nowadays air pollution is becoming a major issue in urban areas especially megacities such as Mumbai, Delhi, and Chennai in India. It has many sources like vehicles, industries, bakeries, domestic sector, load from cremation and hotels. Mumbai is the capital of the state of Maharashtra and is the most populated city in India. Furthermore, it is the financial, commercial and entertainment capital of India with the total population of more than eighteen million people (Census, 2011).

Air quality studies are done with monitoring and modeling techniques (Gulia *et al.*, 2015; Kumar *et al.*, 2015; Li *et al.*, 2011). Air quality modeling requires emission data, meteorological data and surface characteristics for the region, collecting which can prove to be rigorous and tedious. Air quality monitoring is carried out by many authorities to communicate air quality status and this data can be easily collected. The monitoring data is a combination of emission sources and meteorological data which represents air quality status at a point. The objective of this study is to understand the air quality status through the use of interpolation techniques leading to delineating of spatial pattern over a region. The outcome of the study will significantly help in air pollution management, wherein a small dataset can be used for air quality mapping of a larger area. The study has been carried out in three locations within the Mumbai city to analyse air quality. ArcGIS was used to interpolate these values over the Mumbai region for the year 2008. Many communities and regulatory authorities have installed air quality monitoring stations to communicate air quality level or as a part of research and development. National Environmental Engineering Research Institute (NEERI) is a government organisation working continuously to ascertain the critical conditions of environment

and suggest control options for the environmental problems. NEERI measures ambient air quality concentration at three locations in the Mumbai region namely Kalbadevi, Worli and Parel. In the present study, concentration data-sets at three locations have been interpolated over the Mumbai region and interpolated spatial and seasonal patterns have been observed that can help in rational management of air quality. The study area “Mumbai city” is depicted Fig. 1 where the three stations Parel, Kalbadevi and Worli are presented. Parel is situated in central Mumbai and represents an influx of huge enterprises in the compounds of the long-gone cotton mills. Kalbadevi is a residential area and an old neighbourhood in Mumbai. Worli is a part of South Mumbai and is one of the original seven islands that constituted the Mumbai city. Now, it has become one of the busiest office areas in the city.

## 2. Methodology

### 2.1 Geographical Information System (GIS) and its applications

Geographical Information System (GIS) is an efficient tool to capture, store, manipulate, manage, and analyse various types of geographical data and represent graphical view for easy understanding. ArcGIS software has several inbuilt tools which

include many operations such as interpolation techniques, network analysis and spatial analysis. ArcGIS is widely used for modeling and simulation using data processing and various mathematical functions. This tool can help to estimate the levels of air pollution at various locations of the study region. This estimation is very useful to formulate control strategies for air quality management. ArcGIS has been applied in several studies for assessing air quality and exposure (Gulliver and Briggs, 2011; Cinderby and Forrester, 2005; Ketzal *et al.*, 2011; Jensen *et al.*, 2001; Kumar *et al.*, 2016; Maantay, 2007; Marquez and Smith, 1999; Pummakarnchana *et al.*, 2005; Sohrabinia and Khorshiddoust, 2007; Vienneau *et al.*, 2009; Wang *et al.*, 2009; Zhang *et al.*, 2008). There are many interpolation techniques available in ArcGIS such as Kriging, Spline and Inverse Distance Weighting (IDW). Interpolation techniques are considered good tools to study spatial and temporal patterns of air quality without requiring data on meteorology and emissions (Candiani *et al.*, 2013; Janssen *et al.*, 2008; van Loon, 1993). The urban Indian case studies show that Inverse Distance Weighting gives a fairly good performance for the interpolation of the concentration of air pollutants as compared to other interpolation techniques, because the regional and local aspects both are incorporated. The evaluation of IDW interpolation technique for the concentration of ambient air quality in Port Blair




Figure 1. Study area and three monitoring sites: Worli, Kalbadevi and Parel

(India) has been reported by Jha *et al.* (2011). Wong *et al.* (2004) have also demonstrated that IDW gives fairly good interpolation of air quality (Kumar *et al.*, 2016). IDW interpolates all values of the points within the sample range as averaging tool and gives better interpolation estimates when the minimum and maximum values of the surface are represented by sample data points. It estimates the value of a point as a basis of a function of two variables: the distance between sampling points and the point at which the value has to be estimated. The concentration of point will have heavier weight if it is proximal to the required point and vice versa. Here, weight is an inverse function of the distance, as demonstrated in the following equation:

$$Z_j = \frac{\sum_{i=1}^n W_i Z_i}{\sum_{i=1}^n W_i} \quad \text{and} \quad W_i = \frac{1}{d_{ji}^p} \quad \text{Eq 1.}$$

where  $Z_j$  is the value of concentration at the  $j^{\text{th}}$  point,  $W_i$  is the weight of observed  $i^{\text{th}}$  point,  $d_{ji}$  is the distance from the  $i^{\text{th}}$  point to the  $j^{\text{th}}$  point,  $p$  is the power and  $n$  is total number of points.

A base map was taken and geo-referencing was done to associate with physical earth space in ArcGIS. Shapefiles were created with the help of the geo-referenced map for air quality monitoring stations and for the Mumbai region. The air quality monitoring locations are denoted by point shapefile and area of south-west Mumbai is denoted by polygon shapefile in Mumbai region. Air quality data, monitored by NEERI (Mumbai), was fed in the attribute of point

shapefiles on the basis of monthly, seasonal and annual average. Colorimetric method for  $\text{SO}_2$ ,  $\text{NO}_x$  and gravimetric method for SPM were used to monitor air quality data with the help of High Volume Sampler. The monitoring station runs for twice a week and collect 104 data in a year as per (National Ambient Air Quality Standard) NAAQS. The Inverse Distance Weighting (IDW) was used to interpolate the data of three point shapefiles for the concentrations of  $\text{SO}_2$ ,  $\text{NO}_x$  and SPM in the study area.

### 3. Results and Discussion

#### 3.1 Results of monitoring data

Based on monitored data-set of ambient concentration at three locations, it can be concluded that the annual average concentration of  $\text{SO}_2$  and  $\text{NO}_x$  is low while concentration of SPM is consistently high all over Mumbai region for the year 2008. The annual permissible limits of NAAQS by Central Pollution Control Board (CPCB) are  $60 \mu\text{g}/\text{m}^3$  for  $\text{SO}_2$  and  $\text{NO}_x$  and  $140 \mu\text{g}/\text{m}^3$  for SPM (CPCB, 2008). Annual average values of the air quality parameters considered in this study are shown in Fig. 2. The maximum annual average concentrations of  $\text{SO}_2$  and  $\text{NO}_x$  are found to be 10 and  $44 \mu\text{g}/\text{m}^3$  in Kalbadevi amongst the three locations, which are well below the standards. Annual average concentrations of  $\text{SO}_2$  and  $\text{NO}_x$  are close to these values at Parel and Worli: 7 and  $36 \mu\text{g}/\text{m}^3$ , respectively. The annual average concentrations of SPM is 251, 253 and  $274 \mu\text{g}/\text{m}^3$  at Worli, Kalbadevi and Parel, respectively, which is above the permissible limit.




Figure 2. Annual average concentration ( $\mu\text{g}/\text{m}^3$ ) for  $\text{SO}_2$ ,  $\text{NO}_x$  and SPM as monitored at three locations

Table 1. Annual values for the three stations and permissible limit values for the three pollutants

Monitoring Station	Annual average concentration ( $\mu\text{g}/\text{m}^3$ )		
	SO <sub>2</sub>	NO <sub>x</sub>	SPM
Worli	8	38	251
Parel	7	36	274
Kalbadevi	10	44	253
Permissible limit	60	60	140

Table 1 enlists the annual values for the three stations and permissible limit values for the three pollutants, which have been provided by CPCB. The SPM concentration at all three locations is higher than the CPCB standard. Table 2 shows seasonal concentrations of three pollutants at the three stations. The SO<sub>2</sub> concentration is maximum at Kalbadevi from the post-monsoon to winter season which shows industrial emission effect. Worli and Parel have higher concentration for SPM in the monsoon which may be because of resuspension of dust. Kalbadevi and Parel have maximum concentration in the winter. Table 3 shows monthly concentrations of three pollutants at the three stations in which daily standard of air quality of pollutants have been included. Ambient air quality standards are not included in Table 2 because standards are given only on a daily and annual basis. It is evident that SPM exceeds the permissible limit of air quality standard at all three locations while the other two parameters (SO<sub>2</sub> and NO<sub>x</sub>) are within limits. The monthly concentration of SPM at all three locations are within limits in the monsoon season because of washout due to rainfall.

Figs. 3 (a), (b) and (c) show bar plots of seasonal average concentration for air pollutants SO<sub>2</sub>, NO<sub>x</sub> and SPM at the three monitoring sites respectively. Seasons are considered as follows: December and January represent the winter season, February, March, April and May are the pre-monsoon season, June, July, August and September are the monsoon months and the post-monsoon season comprises of October

and November. Seasonal average concentrations of SO<sub>2</sub>, NO<sub>x</sub> and SPM were highest in the winter season, and maximum seasonal concentrations were observed at Kalbadevi for SO<sub>2</sub>, NO<sub>x</sub> and SPM (24, 76 and 414  $\mu\text{g}/\text{m}^3$ , respectively). SPM concentrations were high in each season except during monsoon at Worli and Parel. It may be because these are dense and congested regions where resuspension of particulate matter occurs more. Figs. 4 (a), (b) and (c) represent monthly variations of SO<sub>2</sub>, NO<sub>x</sub> and SPM, respectively, for the year 2008. The concentration of pollutants was found to start decreasing from winter season to pre-monsoon season and concentrations of all pollutants was least in the monsoon season.

### 3.2 Results of IDW interpolation

The collected monitored dataset was used in ArcGIS 9.3 to interpolate the concentration data-set of air quality using IDW. Figs. 5 to 9 show the results of use of IDW interpolation technique. Figs. 5 (a), (b) and (c) show contour plots of annual average for air pollutants SO<sub>2</sub>, NO<sub>x</sub> and SPM respectively. Annual average concentrations of SO<sub>2</sub> and NO<sub>x</sub> are maximum at Kalbadevi, but SPM is higher in Parel at 274  $\mu\text{g}/\text{m}^3$ . In these figures, spatial patterns indicate that NO<sub>x</sub> and SO<sub>2</sub> may be transported from Kalbadevi to Worli and from Worli to Parel, However, in the case of SPM, pollutant may be transported from Parel to Worli and Kalbadevi over the year.

Table 2. Seasonal concentrations ( $\mu\text{g}/\text{m}^3$ ) of three pollutants at the three stations

Locations/ Season	Worli			Kalbadevi			Parel		
	SO <sub>2</sub>	NO <sub>x</sub>	SPM	SO <sub>2</sub>	NO <sub>x</sub>	SPM	SO <sub>2</sub>	NO <sub>x</sub>	SPM
Winter	16	70	323	24	74	414	14	62	410
Premonsoon	9	36	231	8	36	259	8	32	296
Monsoon	3	15	226	4	19	142	3	16	151
Postmonsoon	10	58	269	14	71	300	7	51	343

Table 3. Monthly concentrations ( $\mu\text{g}/\text{m}^3$ ) of three pollutants at the three locations

Locations/ Pollutants	Worli			Kalbadevi			Parel			Standard (Daily)		
	SO <sub>2</sub>	NO <sub>x</sub>	SPM	SO <sub>2</sub>	NO <sub>x</sub>	SPM	SO <sub>2</sub>	NO <sub>x</sub>	SPM	SO <sub>2</sub>	NO <sub>x</sub>	SPM
JAN	18	68	322	23	64	397	15	64	436	80	80	200
FEB	14	56	345	13	60	398	13	39	417	80	80	200
MAR	9	40	209	10	39	261	8	37	273	80	80	200
APR	9	30	228	8	28	249	7	33	296	80	80	200
MAY	4	17	142	2	17	127	4	20	196	80	80	200
JUN	2	13	207	3	16	147	2	15	176	80	80	200
JUL	2	9	211	3	16	147	3	12	138	80	80	200
AUG	3	13	285	3	17	127	3	15	125	80	80	200
SEP	3	25	202	7	27	147	4	22	163	80	80	200
OCT	9	55	275	10	58	276	7	43	330	80	80	200
NOV	11	61	262	18	83	323	7	58	356	80	80	200
DEC	13	71	324	25	83	431	12	60	383	80	80	200




Figure 3. Seasonal variation of concentration ( $\mu\text{g}/\text{m}^3$ ) of (a) SO<sub>2</sub>, (b) NO<sub>x</sub> and (c) SPM in 2008




Figure 4. Monthly variation of (a) SO<sub>2</sub>, (b) NO<sub>x</sub> and (c) SPM concentration ( $\mu\text{g}/\text{m}^3$ ) in 2008




Figure 5. Annual average concentration ( $\mu\text{g}/\text{m}^3$ ) of (a)  $\text{SO}_2$ , (b)  $\text{NO}_x$  and (c) SPM

Figs. 6 (a), (b) and (c) exhibit contour plots of average of monsoon period of the year. This is less for all pollutants, though SPM in Worli and Parel is still higher than the permissible limit of ambient air quality standards. SPM concentration is extremely high at  $226 \mu\text{g}/\text{m}^3$  in Worli for the monsoon period. Here,  $\text{SO}_2$  may have been transported from Parel to Kalbadevi to Worli.  $\text{NO}_x$  may be transported from Kalbadevi to Parel to Worli. However, for SPM, spatial pattern is found to be from Worli to Parel and Kalbadevi.

Figs. 7 (a), (b) and (c) are contour plots of average for  $\text{SO}_2$ ,  $\text{NO}_x$  and SPM during post-monsoon months of the year.  $\text{SO}_2$  and  $\text{NO}_x$  concentrations are as low as 7 and  $51 \mu\text{g}/\text{m}^3$  in Parel. The highest concentrations of  $\text{SO}_2$  and  $\text{NO}_x$  are in Kalbadevi at 14 and  $71 \mu\text{g}/\text{m}^3$  respectively. SPM concentration is high in Parel at

$343 \mu\text{g}/\text{m}^3$  and low in Worli at  $269 \mu\text{g}/\text{m}^3$ , though both are higher than the standard limit. Here too, it may be observed that pollutants  $\text{SO}_2$  and  $\text{NO}_x$  may be transported from Kalbadevi to Worli to Parel, but SPM may be transported in the opposite direction.

Contour plots for winter season are shown in Figs. 8 (a), (b) and (c).  $\text{SO}_2$  and  $\text{NO}_x$  are low in Parel at 14 and  $62 \mu\text{g}/\text{m}^3$ , respectively. Kalbadevi has a concentration as high as 24, 74 and  $414 \mu\text{g}/\text{m}^3$  for  $\text{SO}_2$ ,  $\text{NO}_x$  and SPM respectively. Remarkably, here SPM apparently followed the opposite flow direction: from Kalbadevi and Parel to Worli. The concentration of SPM in Worli was  $323 \mu\text{g}/\text{m}^3$ , which is low. Here, too, pollutants  $\text{SO}_2$  and  $\text{NO}_x$  had the same spatial pattern of being transported from Kalbadevi to Worli to Parel but SPM may go from Kalbadevi and Parel to Worli.




Figure 6. Monsoon (June, July, August and September) concentrations ( $\mu\text{g}/\text{m}^3$ ) of (a)  $\text{SO}_2$ , (b)  $\text{NO}_x$  and (c) SPM




Figure 7. Post-monsoon (October and November) concentration ( $\mu\text{g}/\text{m}^3$ ) of (a)  $\text{SO}_2$ , (b)  $\text{NO}_x$  and (c) SPM

Figs. 9 (a), (b) and (c) show contour plots for the pre-monsoon season. Here, the spatial pattern of  $\text{SO}_2$ ,  $\text{NO}_x$  and SPM concentrations do not match. For  $\text{SO}_2$ , Parel and Kalbadevi display fairly low concentrations ( $8 \mu\text{g}/\text{m}^3$ ) which are comparable to the values observed in Worli ( $9 \mu\text{g}/\text{m}^3$ ). For  $\text{NO}_x$ , Worli and Kalbadevi have the same concentration at  $36 \mu\text{g}/\text{m}^3$ , which is maximum. For SPM, Parel showed higher concentration ( $296 \mu\text{g}/\text{m}^3$ ) than Worli ( $231 \mu\text{g}/\text{m}^3$ ). For the entire region, SPM concentration was higher than the permissible limit, except during the monsoon season.

The average wind speed and maximum temperature of every month has been represented in Fig. 10. The maximum wind speed is in the monsoon season, leading to low concentration level of pollutants. The temperature was high and wind speed was low in

October, November and December. Fig. 11 shows SPM concentration in terms of monthly average. SPM concentration is low from May to September, which corresponds to the monsoon season in the study area. National Ambient Air Quality Standard (NAAQS; CPCB, 2008) prescribes annual and daily air quality standard 140 and  $200 \mu\text{g}/\text{m}^3$ , respectively for SPM but the measured values were higher than that except during the monsoon season at all locations. The concentration is low during the monsoon season and high during the remaining seasons due to precipitation. Rainfall is high from June to September peaking in July, as shown in Fig. 11. The high rainfall washout the pollutants and show lower pollution levels in that period. Temperature, wind speed and rainfall data was collected from Indian Meteorological Department, Mumbai.




Figure 8. Winter (December and January) concentration ( $\mu\text{g}/\text{m}^3$ ) of (a)  $\text{SO}_2$ , (b)  $\text{NO}_x$  and (c) SPM




Figure 9. Pre-monsoon (February, March, April and May) concentration ( $\mu\text{g}/\text{m}^3$ ) of (a)  $\text{SO}_2$ , (b)  $\text{NO}_x$  and (c) SPM

### 5. Conclusions

Interpolation technique of ArcGIS was used for air pollutants such as  $\text{SO}_2$ ,  $\text{NO}_x$  and SPM for Mumbai region. Inverse Distance Weighting (IDW) tool of ArcGIS was used for air quality assessment with a view to understand the variability of air pollutant concentrations across the city. The analyses have been performed for four distinct seasons: winter, pre-monsoon, monsoon and post-monsoon. The city experiences distinct drop in air pollution during rainy season (monsoon), which spans over four months (June-September). The spatial pattern of annual average, winter and post-monsoon seasons displayed

similar pattern for  $\text{SO}_2$  and  $\text{NO}_x$ . The pre-monsoon and post-monsoon seasons showed a similar pattern for SPM. Except these, all showed different patterns for pollutant over the region. All three pollutants showed different patterns in pre-monsoon and monsoon. IDW technique appears to be an appropriate navigation tool for analysis of spatial variation of air quality. It does not require meteorological and emission data, with monitored concentration of pollutants being the only dataset needed. Monitoring data is easy and economical to obtain while collection of emission and meteorological data needs more resources in terms of time and costs.




Figure 10. Monthly maximum temperature and average wind speed




Figure 11. SPM concentrations (average of each month) for three locations of Mumbai compared with annual and daily air quality standard (140 and 200  $\mu\text{g}/\text{m}^3$  respectively)

However, this study has some limitations. For instance, air quality data was available only at three monitoring sites over the entire region and measurement on meteorological parameters was not available at sites. Enhanced air quality monitoring network and inclusion of meteorological parameters will help understand the atmospheric transportation of pollutants, leading to rigorous spatial interpolation and prediction.

## References

- Alcamo J, Mayerhofer P, Guardans R, van Harmelen T, van Minnen J, Onigkeit J, Posch M, de Vries B. An integrated assessment of regional air pollution and climate change in Europe : findings of the AIR-CLIM Project. *Environmental Science and Policy* 2002; 5(4): 257-72.
- Bollen J, Hers S, van der Zwaan B. An integrated assessment of climate change, air pollution, and energy security policy. *Energy Policy* 2010; 38(8): 4021-30.
- Bollen J, van der Zwaan B, Brink C, Eerens H. Local air pollution and global climate change: A combined cost-benefit analysis. *Resource and Energy Economics* 2009; 31(3): 161-81.
- Candiani G, Carnevale C, Finzi G, Pisoni E, Volta M. A comparison of reanalysis techniques: applying optimal interpolation and Ensemble Kalman Filtering to improve air quality monitoring at mesoscale. *Science of the Total Environment* 2013; 458-460: 7-14.
- Census. Census of India Organization [homepage on the Internet]. Ministry of Home Affairs, Government of India. 2011. Available from: [http://www.censusindia.gov.in/Central Pollution Control Board \(CPCB\). National Ambient Air Quality Monitoring 2008. 2008](http://www.censusindia.gov.in/Central Pollution Control Board (CPCB). National Ambient Air Quality Monitoring 2008. 2008).
- Cinderby S, Forrester J. 2005. Facilitating the local governance of air pollution using GIS for participation. *Applied Geography* 2005; 25(2): 143-58.
- Contini D, Donato A, Elefante C, Grasso FM. Analysis of particles and carbon dioxide concentrations and fluxes in an urban area : Correlation with traffic rate and local micrometeorology. *Atmospheric Environment* 2012; 46, 25-35.
- Duh JD, Shandas V, Chang H, George LA. Rates of urbanisation and the resiliency of air and water quality. *Science of the Total Environment* 2008; 400(1-3): 238-56.
- Gulia S, Shrivastava A, Nema AK, Khare M. Assessment of urban air quality around a heritage site using AERMOD: a case study of Amritsar city, India. *Environmental Modeling and Assessment* 2015; 20(6): 599-608. doi: 10.1007/s10666-015-9446-6
- Gulliver J, Briggs D. STEMS-Air: A simple GIS-based air pollution dispersion model for city-wide exposure assessment. *Science of the Total Environment* 2011; 409(12): 2419-29.
- Huff G, Angeles L. Globalization, industrialization and urbanization in Pre-World War II Southeast Asia. *Explorations in Economic History* 2011; 48(1): 20-36.
- Janssen S, Dumont G, Fierens F, Mensink C. Spatial interpolation of air pollution measurements using CORINE land cover data. *Atmospheric Environment* 2008; 42(20): 4884-903.
- Järvi L, Rannik Ü, Mammarella I, Sogachev, Aalto PP, Keronen P, Siivola E, Kulmala M, Vesala T. Annual particle flux observations over a heterogeneous urban area. *Atmospheric Chemistry and Physics* 2009; 9: 7847-56. doi: 10.5194/acp-9-7847-2009

- Jensen SS, Berkowicz R, Hansen HS, Hertel O. A danish decision-support GIS tool for management of urban air quality and human exposures. *Transportation Research Part D: Transport and Environment* 2001; 6(4): 229-41.
- Jha DK, Sabesan M, Das A, Vinithkumar NV, Kirubakaran R. Evaluation of interpolation technique for air quality parameters in Port Blair, India. *Universal Journal of Environmental Research and Technology* 2011; 1(3): 301-10.
- Kan H, Chen R, Tong S. Ambient air pollution, climate change, and population health in China. *Environmental International* 2012; 42: 10-19.
- Kanada M, Fujita T, Fujii M, Ohnishi S. The long-term impacts of air pollution control policy : historical links between municipal actions and industrial energy efficiency in Kawasaki City, Japan. *Journal of Cleaner Production* 2013; 58: 92-101.
- Ketzel M, Berkowicz R, Hvidberg M, Jensen SS, Raaschou-Nielsen O. Evaluation of AirGIS : a GIS-based air pollution and human exposure modelling system. *International Journal of Environment and Pollution* 2011; 47: 226-38. doi: 10.1504/IJEP.2011.047337
- Kim S. Industrialization and urbanization: did the steam engine contribute to the growth of cities in the United States?. *Explorations in Economic History* 2005; 42(4): 586-98.
- Kumar A, Dikshit AK, Fatima S, Patil RS. Application of WRF model for vehicular pollution modelling using AERMOD. *Atmospheric and Climate Sciences* 2015; 5(2): 57-62. doi: 10.4236/acs.2015.52004
- Kumar A, Gupta I, Brandt J, Kumar R, Dikshit AK, Patil RS. Air quality mapping using GIS and economic evaluation of health impact for Mumbai city, India. *Journal of the Air and Waste Management Association* 2016; 66(5): 470-81.
- Li C, Hsu NC, Tsay SC. A study on the potential applications of satellite data in air quality monitoring and forecasting. *Atmospheric Environment* 2011; 45(22): 3663-75.
- Maantay J. Asthma and air pollution in the Bronx: methodological and data considerations in using GIS for environmental justice and health research. *Health and Place* 2007; 13(1): 32-56.
- Marquez LO, Smith NC. A framework for linking urban form and air quality. *Environmental Modelling and Software* 1999; 14(6): 541-48.
- Nemitz E, Jimenez JL, Huffman JA, Ulbrich IM, Canagaratna MR, Worsnop DR, Guenther AB. An eddy-covariance system for the measurement of surface/atmosphere exchange fluxes of submicron aerosol chemical species-first application above an urban area. *Aerosol Science and Technology* 2008; 42(8): 636-57. doi: 10.1080/02786820802227352
- Pummakarnchana O, Tripathi N, Dutta J. Air pollution monitoring and GIS modeling: a new use of nanotechnology based solid state gas sensors. *Science and Technology of Advanced Materials* 2005; 6(3-4): 251-55.
- Sohrabinia M, Khorshiddoust AM. Application of satellite data and GIS in studying air pollutants in Tehran. *Habitat International* 2007; 31(2): 268-75.
- van Loon M. Testing interpolation and filtering techniques in connection with a semi-Lagrangian method. *Atmospheric Environment. Part A. General Topics* 1993; 27(15): 2351-64.
- Vienneau D, de Hoogh K, Briggs D. A GIS-based method for modelling air pollution exposures across Europe. *Science of the Total Environment* 2009; 408(2): 255-66.
- Wang YQ, Zhang XY, Draxler RR. TrajStat: GIS-based software that uses various trajectory statistical analysis methods to identify potential sources from long-term air pollution measurement data. *Environmental Modelling and Software* 2009; 24(8): 938-39.
- Wong DW, Yuan L, Perlin SA. Comparison of spatial interpolation methods for the estimation of air quality data. *Journal of Exposure Analysis and Environmental Epidemiology* 2004; 14(5): 404-15. doi: 10.1038/sj.jea.7500338
- Yennawar PK. Air pollution measurement system in India. *Studies in Environmental Science* 1978; 2: 91-93.
- Zhang Q, Wei Y, Tian W, Yang K. GIS-based emission inventories of urban scale: a case study of Hangzhou, China. *Atmospheric Environment* 2008; 42(20): 5150-65.

---

*Received 10 May 2016*

*Accepted 6 June 2016*

**Correspondence to**

Dr. Awkash Kumar

Centre for Environmental Science and Engineering,

Indian Institute of Technology Bombay,

Mumbai -400 076,

India

E-mail: [awkash@iitb.ac.in](mailto:awkash@iitb.ac.in); [awkash.narayan@gmail.com](mailto:awkash.narayan@gmail.com)