

REFERENCE

- Bartlett, R. (1999). Characterizing soil redox behavior. In D. Sparks, *Soil Physical Chemistry* (pp. 371-397). Boca Raton, Florida: CRC Press.
- Bongen, A. S. (2009). *Using agricultural land for carbon sequestration*. Retrieved March 8, 2010, from Purdue agriculture department, Purdue university: www.agry.purdue.edu/soils/Csequest.PDF
- Bossio, D., Horwath, W., Mutters, R., and van Kessel, C. (1999). Methane pool and flux dynamics in a rice field following straw incorporation. *Soil Biology and Biochemistry*, 31, 1313-1322.
- Brady, N. C., and Weil, R. R. (2002). *The nature and properties of soils*. (13, Ed.) New Jersey: Prentice-Hall, Inc.
- Bronson, K., Neue, H., Singh, U., and Abao, J. (1997). Automatic chamber measurements of methane and nitrous oxide flux in a flooded rice soil: I. Residue, nitrogen and water management. *Soil Science Society of America Journal*, 61, 981-987.
- Brown, P., and Allison, F. (1916). The influence of some common humus forming materials of narrow and wide nitrogen-carbon ratios on bacterial activities. *Soil Science*, 1, 49-75.
- Buendia, L. V., Neue, H.-U., Wassmann, R., Lantin, R. S., and Javellana, A. M. (1997). Understanding the nature of methane emission from rice ecosystems as basis of mitigation strategies. *Applied Energy*, 56, 433-444.
- Cai, Z., Xing, G., Yan, X., Xu, H., Tsuruta, H., Yagi, K., et al. (1999). Measurements of CH₄ and N₂O emissions from rice paddies in Fengqiu, China. *Soil Science and Plant Nutrition*, 45 (1), 1-13.
- Chalk, P., and Smith, C. (1983). Gaseous loss of nitrogen from plant-soil systems. *Developments in Plant and Soil Science*, 9, 65-89.
- Chidthaisong, A., and Watanabe, I. (1997). Methane formation and emission from flooded rice soil incorporated with ¹³C-labeled rice straw. *Soil Biology and Biochemistry*, 29, 1173-1181.
- Chidthaisong, A., Obata, H., and Watanabe, I. (1999). Methane formation and substrate utilisation in anaerobic rice soils as affected by fertilisation. *Soil Biology and Biochemistry*, 31, 135-143.
- Cicerone, R., and Shetter, J. (1981). Sources of atmospheric methane: measurements in rice paddies and a discussion. *Journal of Geophysical Research*, 86C, 7203-7209.

- Conrad, R., Klose, M., Noll, M., Kemnitz, D., and Bodelier, P. L. (2008). Soil type links microbial colonization of rice roots to methane emission. *Global change biology* , 14, 657-669.
- DEDE. (2003). *Rice in Thailand*. Bangkok: Department of Alternative Energy Development and Efficiency (DEDE).
- Devevre, O., and Horwath, W. (2000). Decomposition of rice straw and microbial carbon use efficiency under different soil temperatures and moistures. *Soil Biology and Biochemistry* , 32 (11-12), 1773-1785.
- Dobermann, A., and Fairhurst, T. (2002). Rice straw management. *Better Crops International* , 16 (Special Supplement), 7-11.
- Doryland, C. (1916). The influence of energy material upon the relation of soil microorganisms to soluble plant food. *N.Dak.Agr.Exp.Sta.Bul.* , 116.
- Duiker, S., and Lal, R. (2000). Carbon budget study using CO₂ flux measurements from a no till system in central Ohio. *Soil and Tillage Research* , 54, 21-30.
- Fitzgerald, G. (1998). *Fate of Carbon in California Rice Systems*. California: University of California.
- Gadde, B., Bonnet, S., Menke, C. M., and Garivait, S. (2009). Air pollutant emissions from rice straw open field burning in India, Thailand and the Philippines. *Environmental Pollution* , 157, 1554-1558.
- Ghosh, S., Majumdar, D., and Jain, M. (2003). Methane and nitrous oxide emissions from an irrigated rice of north India. *Chemosphere* , 51, 181-195.
- Guo, L., and Lin, E. (2001). Carbon sink in cropland soils and the emission of greenhouse gases from paddy soils: a review of work in China. *Chemosphere-global change science* , 413-418.
- Holzappel-Pschorn, A., and Seiler, W. (1986). Methane emission during a cultivation period from an Italian rice paddy. *Journal of Geophysical Research* , 91, 11803-11814.
- Hua, X., Guangxi, X., Cai, Z.-C., and Tsuruta, H. (1997). Nitrous oxide emissions from three rice paddy fields in China. *Nutrient Cycling in Agroecosystems* , 49, 23-28.
- Ikenaga, M., Asakawa, S., Muraoka, Y., and Kimura, M. (2004). Methanogenic archaeal communities in rice roots grown in flooded soil pots: estimation by PCR-DGGE and sequence analyses. *Soil Science and Plant Nutrition* , 50, 701-711.
- IPCC. (2001). *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*. (J. Houghton, Y. Ding, D. Griggs, M. Noguer, P. van der Linden, X. Dai, et al., Eds.) Cambridge: Cambridge University Press.

- IPCC. (2007). *Climate change 2007 synthesis report, Summary for policymakers*. Gland, Switzerland: IPCC Secretariat.
- IPCC. (1996). *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories; Reference Manual (Volume 3)*. IPCC Guidelines for National Greenhouse Gas Inventories.
- Iqbal, J., Hu, R., Lin, S., Hatano, R., Feng, M., and Lu, L. (2009). CO₂ emission in a subtropical red paddy soil (Ultisol) as affected by straw and N-fertilizer applications: A case study in Southern China. *Agriculture, Ecosystems and Environment*, 131, 292-302.
- IRRI, I. R. (2002). *Growth stages of the rice plant*. Retrieved March 1, 2010, from International Rice Research Institute Web site: <http://www.knowledgebank.irri.org/growthStages/growthStages.htm>
- Jacinthe, P. A., Lal, R., and Kimble, J. M. (2002). Carbon budget and seasonal carbon dioxide emission from a central Ohio Luvisol as influenced by wheat residue amendment. *Soil and Tillage Research*, 67, 147-157.
- Jakobsen, P., Patrick Jr, W., and Williams, B. (1981). Sulfide and methane formation in soils and sediments. *Soil Science*, 132, 279-287.
- Jiang, J., Huang, Y., and Zong, L. (2003). Influence of water controlling and straw application on CH₄ and N₂O emissions from rice field. *China Environmental Science*, 23, 552-556.
- Kanokkanjana, K. (2010). *An emission assessment of carbonaceous aerosols from agricultural open burning in Thailand: integrating experimental data and remote sensing*. Bangkok: The joint graduate school of energy and environment.
- Kimura, M. (2000). Anaerobic Microbiology in Waterlogged Rice Fields. In J.-M. Bollag, and G. Stotzky, *Soil biochemistry* (Vol. 10, pp. 35-138). New York: Marcel dekker, Inc.
- Kimura, M., Murasa, J., and Lu, Y. (2004). Carbon cycling in rice field ecosystems in the context of input, decomposition and translocation of organic materials and the fates of their end products (CO₂ and CH₄). *Soil biology and Biochemistry*, 1399-1416.
- Kimura, M., Watanabe, I., Patcharapreecha, P., Panichsakpatana, S., Wada, H., and Takai, Y. (1990). Quantitative estimation of decomposition process of plant debris in paddy field. II. Amounts and C/N ratios of plant debris in tropical paddy soils. *Soil science and plant nutrition*, 36, 33-42.
- Le Mer, J., and Roger, P. (2001). Production, oxidation, emission and consumption of methane by soils: A review. *Eur. J. Soil Biol.*, 37, 25-50.
- Li, C., Kou, Z., Yang, J., Cai, M., Wang, J., and Cao, C. (2010). Soil CO₂ fluxes from direct seeding rice fields under two tillage practices in central China. *Atmospheric Environment*, 44, 2696-2704.

- Liou, R., Huang, S., and Lin, C. (2003). Methane emission from fields with differences in nitrogen fertilizers and rice varieties in Taiwan paddy soils. *Chemosphere* , 50 (2), 237-246.
- Liu, S., Qin, Y., Zou, J., and Liu, Q. (2010). Effects of water regime during rice-growing season on annual direct N₂O emission in a paddy rice-winter wheat rotation system in southeast China. *Science of the Total Environment* , 408, 906-913.
- Luo, Y., and Zhou, X. (2006). *Soil respiration and the environment*. Burlington: Elsevier, Inc.
- Ma, J., Ma, E., Xu, H., Yagi, K., and Cai, Z. (2009). Wheat straw management affects CH₄ and N₂O emissions from rice fields. *Soil Biology and Biochemistry* , 41, 1022-1028.
- Mayer, H., and Conrad, R. (1990). Factors influencing the population of methanogenic bacteria and the initiation of methane production upon flooding of paddy soil. *FEMS Microbiology Ecology* , 73, 103-112.
- Minamikawa, K., and Sakai, N. (2006). The practical use of water management based on soil redox potential for decreasing methane emission from a paddy field in Japan. *Agriculture, Ecosystems and Environment* , 116, 181-188.
- Minamikawa, K., Sakai, N., and Hayashi, H. (2005). The effects of ammonium sulfate application on methane emission and soil carbon content of a paddy field in Japan. *Agriculture, Ecosystems and Environment* , 107 (4), 371-379.
- Naser, H. M., Nagata, O., Tamura, S., and Hatano, R. (2007). Methane emissions from five paddy fields with different amounts of rice straw application in central Hokkaido, Japan. *Soil Science and Plant Nutrition* , 53 (1), 95-101.
- Neue, H., Wassmann, R., Lantin, S., Alberto, M. C., Aduna, J., and Javellana, A. (1996). Factors affecting methane emission from rice fields. *Atmospheric Environment* , 30, 1751-1754.
- Nishimura, S., Yonemura, S., Sawamoto, T., Shirato, Y., Akiyama, H., Sudo, S., et al. (2008). Effect of land use change from paddy rice cultivation to upland crop cultivation on soil carbon budget of a cropland in Japan. *Agriculture, Ecosystems and Environment* , 125, 9-20.
- OAE. (2007). *Agricultural Statistics*. Retrieved March 9, 2010, from Office of Agricultural Economics: http://www.oae.go.th/oae_report/stat_agri/report_result_content.php
- OAE. (2007). *The survey's report of major rice crop year 2005/06*. Bangkok: Office of Agricultural Economics (OAE).

OEPP. (2000). *Thailand's Initial National Communication under the United Nations Framework Convention on Climate Change*. Bangkok: Ministry of Science, Technology and Environment.

Penman, J., Kruger, D., Galbally, I., Hiraishi, T., Nyenzi, B., Emmanul, S., et al. (2000). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. Japan: Institute for Global Environmental Strategies.

Penning, H., and Conrad, R. (2006). Effect of inhibition of Acetoclastic Methanogenesis on growth of Archaeal populations in an anoxic model environment. *Environmental microbiology*, 72 (1), 178-184.

Pidwirny, M. (2006). *The carbon cycle Fundamentals of Physical Geography, 2nd Edition*. Retrieved February 1, 2010, from Physical Geography.net: <http://www.physicalgeography.net/fundamentals/9r.html>

Rath, A. K., Ramakrishnan, B., and Sethunathan, N. (2002). Effect of application of ammonium thiosulphate on production and emission of methane in a tropical rice soil. *Agriculture, Ecosystems and Environment*, 90, 319-325.

Salter, F. J. (1930). The carbon-nitrogen ratio in relation to the accumulation of organic matter in soils. *Soil Science*, 31 (6), 413-430.

Sass, R. (2002). CH₄ emissions from rice agriculture. In IPCC, *Background Papers – IPCC Expert Meetings on Good Practice Guidance and Uncertainty* (pp. 399-417). Institute for Global Environmental Strategies (IGES).

Sass, R., Fisher, F., Harcombe, P., and Turner, F. (1991). Mitigation of methane emissions from rice field: possible adverse effects of incorporated rice straw. *Global Biogeochemical Cycles*, 5, 275-287.

Seiler, W., Holzappel-Pschorn, A., Conrad, R., and Scharffe, D. (1984). Methane emission from rice paddies. *Journal of Atmospheric Chemistry*, 1, 241-268.

Shang, S., and Hsiu, L. (1998). Effect of environmental conditions on methane production and emission from paddy soil. *Agriculture, Ecosystems and Environment*, 69, 69-80.

Singh, S., Singh, J., and Kashyap, A. K. (1999). Methane flux from irrigated rice fields in relation to crop growth and N-fertilization. *Soil Biology and Biochemistry*, 31, 1219-1228.

Singh, Y., and Singh, B. (2001). Efficient Management of Primary Nutrition in the Rice-Wheat System. In P. Kataki, *The Rice-Wheat Cropping Systems of South Asia: Efficient Production Management* (pp. 23-85). New York, USA: Food Products Press.

Singh, Y., Singh, B., Ladha, J., Khind, C., Gupta, R., Meelu, O., et al. (2004). Long-term effect of organic inputs on yield and soil fertility in the rice-wheat rotation. *Soil Science Society of America Journal*, 68, 845-853.

- Smakgahn, K. (2003). *Empirical model and process base model application on methane and nitrous oxide emission from Thai rice fields*. Bangkok: The joint graduate school of energy and environment.
- Smith, P. D. (2007). *Agriculture. In Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the*. NY, USA.: Cambridge University Press.
- Solomon, S. D. (2007). *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. New York: Canbridge University Press.
- Suramaythangkoor, T., and Gheewala, S. H. (2008). Potential of practical implementation of rice straw-based power generation in Thailand. *Energy policy*, 36, 3193-3197.
- Tanji, K., Gao, S., Scardaci, S., and Chow, A. (2003). Characterizing redox status of paddy soils with incorporated rice straw. *Geoderma*, 114, 333-353.
- Towprayoon, S., Smakgahn, K., and Poonkaew, S. (2005). Mitigation of methane and nitrous oxid emissions from drained irrigated rice fields. *Chemosphere*, 59, 1547-1556.
- Wang, B., Neue, H., and Samonte, H. (1999). Factors controlling diel patterns of methane emission via rice. *Nutrient Cycling in Agroecosystems*, 53, 229-235.
- Watanabe, A., Katoh, K., and Kimura, M. (1993). Effect of rice straw application on CH₄ emission from paddy fields. I. Effect of weathering of rice straw in the field during off-crop season. *Soil Science and Plant Nutrition*, 39, 701-706.
- Watanabe, A., Yamada, H., and Kimura, M. (2005). Analysis of temperature effects on seasonal and inter-annual variation in CH₄ emission from rice-planted pots. *Agriculture, Ecosystems and environment*, 105, 439-443.
- Watanabe, I., and Furusaka, C. (1980). Microbial ecology of flooded rice soils. In M. Alexander, *Advances in microbial ecology* (Vol. 4, pp. 125-168). Plenum publishing corporation.
- Watanabe, T., Chairoj, P., Tsuruta, H., Masarngsan, W., Wongwiwatchai, C., Wonprasaid, S., et al. (2000). Nitrous oxide emissions from fertilized upland fields in Thailand. *Nutrient Cycling in Agroecosystems*, 58, 55-65.
- Whitman, W. B., Bowen, T., and Boone, D. (2006). The methanogenic bacteria. In M. Dworkin, S. Falkow, E. Rosenberg, K. Schleifer, and E. Stackebrandt, *The Prokaryotes* (3 ed., Vol. 3, pp. 165-207). New York: Springer.
- Witt, C., Cassman, K., Olk, D., Biker, U., Liboon, S., Samson, S., et al. (2000). Crop rotation and residue management effects on carbon sequestration, nitrogen cycling and productivity of irrigated rice systems. *Plant and Soil*, 263-278.

- Wrage, N., Velthof, G., van Beusichem, M., and Oenema, O. (2001). Role of nitrifier denitrification in the production of nitrous oxide. *Soil Biology and Biochemistry*, 33, 1723-1732.
- Xing, G., Zhao, X., Xiong, Z., Yan, X., Xu, H., Xie, Y., et al. (2009). Nitrous oxide emission from paddy fields in China. *Acta Ecologica Sinica*, 29, 45-50.
- Yagi, K. (1997). Methane emission from paddy fields. *Bull. Natl. Inst. Agro-Environ. Sci.* (14), 96-210.
- Yagi, K., Tsuruta, H., Kanda, K., and Minami, K. (1996). Effect of water management on methane emission from a Japanese rice paddy field: Automated methane monitoring. *Global Biogeochemical Cycles*, 10, 255-267.
- Yamane, I., and Sato, K. (1967). Effect of temperature on the decomposition of organic substances in flooded soil. *Soil Science and Plant Nutrition*, 13, 94-100.
- Yan, X., Du, L., Shi, S., and Xing, G. (2000). Nitrous oxide emission from wetland rice soil as affected by the application of controlled availability fertilizers and mid-season aeration. *Biology and Fertility of Soils*, 32, 60-66.
- Yan, X., Shi, S., Du, L., and Xing, G. (2000). Pathways of N₂O emission from rice paddy soil. *short communication, Soil Biology and Biochemistry*, 32, 437-440.
- Zheng, J., Zhang, X., Li, L., Zhang, P., and Pan, G. (2007). Effect of long-term fertilization on C mineralization and production of CH₄ and CO₂ under anaerobic incubation from bulk samples and particle size fractions of a typical paddy soil. *Agriculture, Ecosystems and Environment*, 120 (2-4), 129-138.
- Zou, J., Huang, Y., Jiang, J., Zheng, X., and Sass, R. (2005). A 3-year field measurement of methane and nitrous oxide emissions from rice paddies in China: Effect of water regime, crop residue, and fertilizer application. *Global Biogeochemical Cycles*, 19, GB2021, doi:10.1029/2004GB002401.
- Zou, J., Huang, Y., Zong, L., Jiang, J., Zheng, X., and Wang, Y. (2003). Effects of water regime and straw application in paddy rice season on N₂O emission from following wheat cropping season. *Scientia Agricultura Sinica*, 36, 409-414.
- ZP, W., Masscheleyn, R. D., and Jr., W. P. (1993). Soil redox and pH effects on methane production in a flooded rice soil. *Soil Science Society of America Journal*, 57, 382-385.



