

บรรณานุกรม

- [1] ปราโมทย์ ประสาทกุล. เป็นคนแก่อีก 20 ปีข้างหน้า จะเหงา หรือไม่เหงา 2549.
- [2] กุศล สุนทรธาดา. โรคเรื้อรังของประชากรในพื้นที่เฝ้าระวังทางประชากรกาญจนบุรี. ประชากรและการพัฒนา 2550.
- [3] Cornoni-Huntley JC, Foley DJ, Guralnik JM. Co-morbidity analysis: a strategy for understanding mortality, disability and use of health care facilities of older people. *Int J Epidemiol* 1991;20 Suppl 1:S8-17.
- [4] Wadhwa A, Sabharwal M, Sharma S. Nutritional status of the elderly. *Indian J Med Res* 1997;106:340-8.
- [5] สุวัฒน์ มหัตนรินทร์กุล. โครงการจัดทำโปรแกรมสำเร็จรูปในการสำรวจสุขภาพจิตในพื้นที่โรงพยาบาลสวนปรุง กรมสุขภาพจิต กระทรวงสาธารณสุข 2545.
- [6] Vergara N, Montoya JE, Luna HG, Amparo JR, Cristal-Luna G. Quality of life and nutritional status among cancer patients on chemotherapy. *Oman Med J* 2013;28:270-4.
- [7] Moreira AC, Carolino E, Domingos F, Gaspar A, Ponce P, Camilo ME. Nutritional status influences generic and disease-specific quality of life measures in haemodialysis patients. *Nutr Hosp* 2013;28:951-7.
- [8] Rasheed S, Woods RT. An investigation into the association between nutritional status and quality of life in older people admitted to hospital. *J Hum Nutr Diet* 2013;n/a-n/a.
- [9] Auld G, Baker S, Bauer L, Koszewski W, Procter SB, Steger MF. EFNEP's Impact on the Quality of Life of Its Participants and Educators. *J Nutr Educ Behav* 2013.
- [10] Langius JAE, Zandbergen MC, Eerenstein SEJ, van Tulder MW, Leemans CR, Kramer MHH, et al. Effect of nutritional interventions on nutritional status, quality of life and mortality in patients with head and neck cancer receiving (chemo)radiotherapy: a systematic review. *Clin Nutr Edinb Scottl* 2013;32:671-8.
- [11] Nadine S. Food insufficiency and the nutritional status if the elderly population: USDA center for nutrition policy and promotion. May 2000.
- [12] Swain JF, McCarron PB, Hamilton EF, Sacks FM, Appel LJ. Characteristics of the diet patterns tested in the optimal macronutrient intake trial to prevent heart disease (OmniHeart): options for a heart-healthy diet. *J Am Diet Assoc* 2008;108:257-65.

- [13] Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403.
- [14] Schusdziarra V, Hausmann M, Wiedemann C, Hess J, Barth C, Wagenpfeil S, et al. Successful weight loss and maintenance in everyday clinical practice with an individually tailored change of eating habits on the basis of food energy density. *Eur J Nutr* 2011;50:351–61.
- [15] Handbook of obesity: clinical applications. 2nd ed. New York: M. Dekker; 2004.
- [16] Heymsfield SS, Wang Z. Evaluation of Total and Regional Adiposity. *Handb. Obes.*, New York: Marcel Dekker; 2004, p. 33.
- [17] Dey DK, Bosaeus I, Lissner L, Steen B. Changes in body composition and its relation to muscle strength in 75-year-old men and women: a 5-year prospective follow-up study of the NORA cohort in Göteborg, Sweden. *Nutr* Burbank Los Angeles Cty Calif 2009;25:613–9.
- [18] Heshka S, Ruggiero A, Bray GA, Foreyt J, Kahn SE, Lewis CE, et al. Altered body composition in type 2 diabetes mellitus. *Int J Obes* 2005 2008;32:780–7.
- [19] Scherzer R, Shen W, Bacchetti P, Kotler D, Lewis CE, Shlipak MG, et al. Comparison of dual-energy X-ray absorptiometry and magnetic resonance imaging-measured adipose tissue depots in HIV-infected and control subjects. *Am J Clin Nutr* 2008;88:1088–96.
- [20] Wang ZM, Pierson RN Jr, Heymsfield SB. The five-level model: a new approach to organizing body-composition research. *Am J Clin Nutr* 1992;56:19–28.
- [21] Gallagher D, Kuznia P, Heshka S, Albu J, Heymsfield SB, Goodpaster B, et al. Adipose tissue in muscle: a novel depot similar in size to visceral adipose tissue. *Am J Clin Nutr* 2005;81:903–10.
- [22] LaForgia J, Dollman J, Dale MJ, Withers RT, Hill AM. Validation of DXA body composition estimates in obese men and women. *Obes Silver Spring Md* 2009;17:821–6.
- [23] Withers RT, LaForgia J, Pillans RK, Shipp NJ, Chatterton BE, Schultz CG, et al. Comparisons of two-, three-, and four-compartment models of body composition analysis in men and women. *J Appl Physiol Bethesda Md* 1985 1998;85:238–45.
- [24] De Lorenzo A, Andreoli A, Matthie J, Withers P. Predicting body cell mass with bioimpedance by using theoretical methods: a technological review. *J Appl Physiol Bethesda Md* 1985 1997;82:1542–58.

- [25] Sheng HP, Huggins RA. A review of body composition studies with emphasis on total body water and fat. *Am J Clin Nutr* 1979;32:630–47.
- [26] Bray GA, DeLany JP, Volaufova J, Harsha DW, Champagne C. Prediction of body fat in 12-y-old African American and white children: evaluation of methods. *Am J Clin Nutr* 2002;76:980–90.
- [27] Ginde SR, Geliebter A, Rubiano F, Silva AM, Wang J, Heshka S, et al. Air displacement plethysmography: validation in overweight and obese subjects. *Obes Res* 2005;13:1232–7.
- [28] Völgyi E, Tylavsky FA, Lyytikäinen A, Suominen H, Alén M, Cheng S. Assessing body composition with DXA and bioimpedance: effects of obesity, physical activity, and age. *Obesity (Silver Spring)* 2008;16:700–5.
- [29] Fakhrawi DH, Beeson L, Libanati C, Feleke D, Kim H, Quansah A, et al. Comparison of body composition by bioelectrical impedance and dual-energy x-ray absorptiometry in overweight/obese postmenopausal women. *J Clin Densitom* 2009;12:238–44.
- [30] Lukaski HC, Johnson PE, Bolonchuk WW, Lykken GI. Assessment of fat-free mass using bioelectrical impedance measurements of the human body. *Am J Clin Nutr* 1985;41:810–7.
- [31] Kyle UG, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Gómez JM, et al. Bioelectrical impedance analysis—part I: review of principles and methods. *Clin Nutr Edinb Scotl* 2004;23:1226–43.
- [32] กมลรัชฎ์จงชนากร. การใช้เครื่องวิเคราะห์ส่วนประกอบของร่างกายในเวชปฏิบัติ. วารสารโรงพยาบาลเจริญกรุงประชารักษ์ 2552;ปีที่ 5.
- [33] Slinde F, Grönberg A, Engström C-P, Rossander-Hulthén L, Larsson S. Body composition by bioelectrical impedance predicts mortality in chronic obstructive pulmonary disease patients. *Respir Med* 2005;99:1004–9.
- [34] Fürstenberg A, Davenport A. Assessment of body composition in peritoneal dialysis patients using bioelectrical impedance and dual-energy x-ray absorptiometry. *Am J Nephrol* 2011;33:150–6.
- [35] Wagner D, Adunka C, Kniepeiss D, Jakoby E, Schaffellner S, Kandlbauer M, et al. Serum albumin, subjective global assessment, body mass index and the bioimpedance analysis in the assessment of malnutrition in patients up to 15 years after liver transplantation. *Clin Transplant* 2011;25:E396–400.

- [36] Buffa R, Mereu RM, Putzu PF, Floris G, Marini E. Bioelectrical impedance vector analysis detects low body cell mass and dehydration in patients with Alzheimer's disease. *J Nutr Health Aging* 2010;14:823–7.
- [37] Schwenk A, Beisenherz A, Römer K, Kremer G, Salzberger B, Elia M. Phase angle from bioelectrical impedance analysis remains an independent predictive marker in HIV-infected patients in the era of highly active antiretroviral treatment. *Am J Clin Nutr* 2000;72:496–501.
- [38] Sánchez-Lara K, Turcott JG, Juárez E, Guevara P, Núñez-Valencia C, Oñate-Ocaña LF, et al. Association of nutrition parameters including bioelectrical impedance and systemic inflammatory response with quality of life and prognosis in patients with advanced non-small-cell lung cancer: a prospective study. *Nutr Cancer* 2012;64:526–34.
- [39] Hill GL, Jonathan E. Rhoads Lecture. Body composition research: implications for the practice of clinical nutrition. *JPEN J Parenter Enteral Nutr* 1992;16:197–218.
- [40] Walter-Kroker A, Kroker A, Mattiucci-Guehlke M, Glaab T. A practical guide to bioelectrical impedance analysis using the example of chronic obstructive pulmonary disease. *Nutr J* 2011;10:35.
- [41] Van der Meij BS, Langius JAE, Smit EF, Spreeuwenberg MD, von Blomberg BME, Heijboer AC, et al. Oral nutritional supplements containing (n-3) polyunsaturated fatty acids affect the nutritional status of patients with stage III non-small cell lung cancer during multimodality treatment. *J Nutr* 2010;140:1774–80.
- [42] Ha L, Hauge T, Iversen PO. Body composition in older acute stroke patients after treatment with individualized, nutritional supplementation while in hospital. *Bmc Geriatr* 2010;10:75.
- [43] Guigoz Y, Vellas B. The Mini Nutritional Assessment (MNA) for grading the nutritional state of elderly patients: presentation of the MNA, history and validation. *Nestlé Nutr Work Ser Clin Perform Program* 1999;1:3–11; discussion 11–12.
- [44] Rubenstein LZ, Harker JO, Salvà A, Guigoz Y, Vellas B. Screening for undernutrition in geriatric practice: developing the short-form mini-nutritional assessment (MNA-SF). *J Gerontol A Biol Sci Med Sci* 2001;56:M366–372.
- [45] Kaiser MJ, Bauer JM, Ramsch C, Uter W, Guigoz Y, Cederholm T, et al. Validation of the Mini Nutritional Assessment short-form (MNA-SF): a practical tool for identification of nutritional status. *J Nutr Health Aging* 2009;13:782–8.
- [46] Kahneman D, Deaton A. High income improves evaluation of life but not emotional well-being. *Proc Natl Acad Sci* 2010;107:16489–93.

- [47] Ravasco P, Monteiro-Grillo I, Camilo M. Individualized nutrition intervention is of major benefit to colorectal cancer patients: long-term follow-up of a randomized controlled trial of nutritional therapy. *Am J Clin Nutr* 2012;96:1346–53.
- [48] Scott MK, Shah NA, Vilay AM, Thomas J 3rd, Kraus MA, Mueller BA. Effects of peridialytic oral supplements on nutritional status and quality of life in chronic hemodialysis patients. *J Ren Nutr Off J Counc Ren Nutr Natl Kidney Found* 2009;19:145–52.
- [49] Ha L, Hauge T, Spenning AB, Iversen PO. Individual, nutritional support prevents undernutrition, increases muscle strength and improves QoL among elderly at nutritional risk hospitalized for acute stroke: a randomized, controlled trial. *Clin Nutr Edinb Scottl* 2010;29:567–73.
- [50] Sugawara K, Takahashi H, Kasai C, Kiyokawa N, Watanabe T, Fujii S, et al. Effects of nutritional supplementation combined with low-intensity exercise in malnourished patients with COPD. *Respir Med* 2010;104:1883–9.
- [51] Stange I, Bartram M, Liao Y, Poeschl K, Kolpatzik S, Uter W, et al. Effects of a low-volume, nutrient- and energy-dense oral nutritional supplement on nutritional and functional status: a randomized, controlled trial in nursing home residents. *J Am Med Dir Assoc* 2013;14:628.e1–8.
- [52] Burke L, Lee AH, Jancey J, Xiang L, Kerr DA, Howat PA, et al. Physical activity and nutrition behavioural outcomes of a home-based intervention program for seniors: a randomized controlled trial. *Int J Behav Nutr Phys Act* 2013;10:14.
- [53] Leenders M, Verdijk LB, Van der Hoeven L, Van Kranenburg J, Nilwik R, Wodzig WKWH, et al. Protein supplementation during resistance-type exercise training in the elderly. *Med Sci Sports Exerc* 2013;45:542–52.