

LIST OF CONTENTS

Chapter	Page
I INTRODUCTION	1
Rational for the study.....	1
Objectives of the study.....	2
Scope of the study.....	2
Hypotheses of the study.....	2
Anticipated outcomes.....	3
II LITERATURE REVIEW	4
Rice (<i>Oryza sativa</i> L.).....	4
Components of rice seed.....	4
Starch in rice seed.....	5
Rice damage in cultivation process.....	6
Postharvest loss and damage of rice product.....	10
Stored product insect pests.....	11
Red flour beetles: <i>Tribolium castaneum</i> (Coleoptera: Tenebrionidae).....	11
Identification.....	12
Biology and life cycle of red flour beetle: <i>Tribolium castaneum</i>	12
Amylase.....	14
Alpha-amylase.....	15
Alpha-amylase in animals.....	15
Human α -amylase.....	16
Alpha-amylases in microorganisms.....	18
Cereal α -amylases.....	19
Insect α -Amylase.....	20
Beta-amylase.....	22
Gamma-amylase (Glucoamylase).....	22
Alpha-amylase inhibitor.....	22

LIST OF CONTENTS (CONT.)

Chapter	Page
III RESEARCH METHODOLOGY	24
Materials	24
Methods.....	28
IV RESULTS AND DISCUSSION	36
Amplification and cloning of <i>T. castaneum</i> α -amylase gene (TcAm).....	36
The 1,650 bp of TcAm gene sequence analysis.....	36
Expression of 1,650 pb recombinant <i>Tribolium castaneum</i> α -amylase.....	37
Growth, survival and α -amylase gene expression of <i>Tribolium castaneum</i> on different starch diets.....	48
Cloning and sequence analysis of 1,470 bp TcAm gene	52
Expression & Purification of recombinant <i>Tribolium castaneum</i> α -amylase	53
Enzymatic activity and biochemical properties of recombinant <i>Tribolium</i> <i>castaneum</i> α -amylase	63
V CONCLUSION	69
REFERENCE	71
APPENDIX	85
BIOGRAPHY	93

LIST OF TABLES

Table	Page
1	Oligonucleotide primers used for DNA amplification of <i>Tribolium castaneum</i> α -amylase gene and insect β -actin gene actin30
2	Activity of recombinant <i>Tribolium castaneum</i> α -amylase was determined with the 2-Chloro-4-nitrophenyl- α -D-maltotrioside substrate at 50 °C 30 min.66

LIST OF FIGURES

Figure		Page
1	The different of indica, javanica and japonica rice.	4
2	The components of rice seed	5
3	Structures of starch consist of amylopectin and amylose.....	6
4	Rice diseases.....	7
5	Rice animal pests.....	8
6	Rice weeds.....	9
7	Yellow stem borer and Bagnall	10
8	Red flour beetles: <i>Tribolium castaneum</i> (Coleoptera: Tenebrionidae)	12
9	Life cycle of red flour beetle	13
10	Life cycle of red flour beetle 2	14
11	Stereo drawing of a schematic representation of the polypeptide chain fold of human pancreatic α -amylase. Also indicated is the relative positioning of the three structural domains present in this protein.	17
12	Ribbon diagram of HSAm α showing the active site and secondary saccharide binding sites.....	17
13	Stereo view of the ribbon model of the <i>Bacillus</i> sp. α -amylase structure.....	18
14	Ribbon diagram of barley α -amylase	20
15	Three acidic side chains in (A) PPA: Asp197, Glu233 and Asp300 corresponding to (B) TMA: Asp185, Glu222 and Asp287	21
16	Agarose gel electrophoresis of PCR product.....	38
17	Agarose gel electrophoresis of colony PCR screening of the 1,650 bp TcAm recombinant plasmid with specific primers.....	38
18	Agarose gel electrophoresis of <i>Bam</i> HI and <i>Xho</i> I restriction enzyme analysis of 1,650 bp TcAm.....	39
19	Diagrams depicting comparison between (A) <i>Tribolium castaneum</i> α -amylase mRNA (NCBI Reference Sequence: NM_001114376.1) and (B) TcAm cDNA cloned in this study.....	39

LIST OF FIGURES (CONT.)

Figure		Page
20	The comparison of 1,650 bp TcAm cDNA cloned nucleotide sequences with <i>Tribolium castaneum</i> α -amylase mRNA (NCBI Reference Sequence: NM_001114376.1).....	40
21	The cDNA sequence and its deduced protein sequence of <i>Tribolium castaneum</i> α -amylase gene. Abnormal stop codon (grey box) is shown in the first extra region.	47
22	10% SDS-PAGE gel of 1650 bp α -amylase - pET32a (+) recombinant cells was induced with 1mM IPTG at 37 °C.	48
23	Larvae of <i>Tribolium castaneum</i> were fed with difference diet after 3 week.	50
24	Effect of different diets on <i>Tribolium castaneum</i> mortality.....	51
25	Agarose gel electrophoresis of PCR product amplified from <i>Tribolium castaneum</i> DNA, isolated RNA in 1, 2, 3, 4, 6 and 8 week.	51
26	Agarose gel electrophoresis of colony PCR screening of the 1,470 bp TcAm recombinant plasmid with specific primers	54
27	Agarose gel electrophoresis of BamHI and XhoI restriction enzyme analysis of 1,470 bp TcAM.....	55
28	Three clones of 1,470 bp <i>Tribolium castaneum</i> α -amylase genes nucleotide sequence alignment compared with the NCBI nucleotide database (NCBI Reference Sequence: NM_001114376.1) tool.....	56
29	Comparison of deduced amino acid sequences of three cDNAs (Gene Bank Accession Number KF247314, KF247315, KF247316) cloned from <i>Tribolium castaneum</i> , encoding three putative α -amylases, and the NCBI database (NCBI Reference Sequence: NM_001114376.1)..	59

LIST OF FIGURES (CONT.)

Figure		Page
30	10% SDS-PAGE gel of pET32a (+) – 1,470 bp α -amylase recombinant cells was induced with 1mM IPTG at 37 °C for 1hr.	60
31	10% SDS-PAGE of purified TcAm using Ni ²⁺ affinity chromatography....	60
32	Mascot search of LC-MS/MS result showing that the tryptic digests of 53.3 kDa protein have exactly matched to α -amylase of <i>Tribolium castaneum</i>	61
33	Identification of peptide mass fingerprints of α -amylase protein that was induced with 1 mM IPTG at 37 °C for 2 hr by LC-MS. The protein has exactly matched to α -amylase of <i>Tribolium castaneum</i>	62
34	Starch zymogram analysis of recombinant TcAm under non-denature condition. Native PAGE stained with Coomassie brilliant blue R-250	65
35	Kinetic of recombinant TcAm measured with 2-Chloro-4-nitrophenyl- α -D-maltotrioxide substrate at 50 °C, 30 min runtime and 1 min interval.....	66
36	Relative enzyme activity (% of maximum) of purified TcAm measured at different temperatures ranging from 20-80 °C for 30 min.....	67
37	Determination of the thermal stability at different temperatures for different time points.	67
38	Effect of pH on enzyme activity was assayed at different pH values range from 4.0-9.0 using 2-Chloro-4-nitrophenyl- α -D-maltotrioxide as a substrate.....	68
39	Effects of some metal ions and reagent on the TcAm activity was determined using 2-Chloro-4-nitrophenyl- α -D-maltotrioxide as a substrate at 50 °C, pH 7 for 30 min.	68