

ACKNOWLEDGEMENTS

The success of this thesis can be succeeded by the attentive support from many people. First of all, I would like to express my heartfelt gratitude and deep appreciation to Asst. Prof. Anadi Nitithamyong, my major advisor for her mercy, invaluable dedication, wisdom and valuable guidance and encouragement throughout my study. This work would not succeed without financially supported by funding from the Office of The National Reserch Council of Thailand.

I also would like to express my sincere to my co-advisor, Dr. Nattapol Tungsuphoom for his kindness, suggestion, valuable advice and guidance in this study.

Very special thanks contribute to Prof. Vanna Tulyathan for her valuable time to be the chair of my examination. My special thanks to Asst. Prof. Sitima Jittinandana, Miss Rungrat Chamchan and Mr. Somsak Mekkluen for their supports and assistance in laboratory equipments and facilities. My special thanks to Mrs. Patcharin Poochaiwatananon, Department of Chemistry, Faculty of Science, Mahidol University for her technical assistance in NMR spectroscopy and thank to Mr. Nawapol Uduay of The Salaya Central Scientific Instrument Facility, Mahidol University for his help and advice in FTIR spectroscopy. I also would like to thank for kindness and help of all staffs in Food Science Units of The Institute of Nutrition, Mahidol University.

I deep appreciation and grateful to my beloved friends, Miss Paninart Kanjanatiwat, Miss Rungrapa Tubrisod, Miss Numphung Rungruen, Miss Pattarmon Mackamarn and Miss Pattamas Karaket and Mr. Worayut Kongkeaw and all friends for their love, understanding, help, encouragement and every things they give to me. Moreover, I express my sincere to those not signed out here.

Finally, I am greatly indebted to my family for invaluable benevolences. The usefulness of this thesis, I dedicate to my family and all educators of my life

Gunsanee Suparpkul