

## REFERENCES

- Abbadì A, Domergue F, Bauer J, Napier J, Welte R, Zähringer U, Cirpus P, Heintz E. Biosynthesis of very-long-chain polyunsaturated fatty acids in transgenic oilseeds: Constraints on their accumulation. *Plant Cell*. 2004; 16: 2734-2748.
- Abdel-Aal EM, Young JC, Rabalski I. Anthocyanin composition in black, blue, pink, purple, and red cereal grains. *J Agric Food Chem*. 2006; 54(13): 4696-4704.
- Abdel-Aal E-SM, Hucl P. A rapid method for quantifying total anthocyanins in blue aleurone and purple pericarp wheats. *Cereal Chem*. 1999; 76(3): 350-354.
- Abraham SK, Graf U. Protection by coffee against somatic genotoxicity in *Drosophila*: role of bioactivation capacity. *Food Chem Toxicol*. 1996; 34: 1-14.
- Abraham SK, Singh SP, Kesavan PC. In vivo antigenotoxic effects of dietary agents and beverages co-administered with urethane: assessment of the role of glutathione S-transferase activity. *Mutat Res*. 1998; 413(2): 103-110.
- Abraham SK. Antigenotoxicity of coffee in the *Drosophila* assay for somatic mutation and recombination. *Mutagenesis*. 1994; 9: 383-386.
- Adams SP. Mechanisms of nutrition bar hardening: effect of hydrolyzed whey protein and carbohydrate source. Master thesis. Utah State University, USA. 2008.
- Afaq F, Syed DN, Malik A, Hadi N, Sarfaraz S, Kweon MH, Khan N, Zaid MA, Mukhtar H. Delphinidin, an anthocyanidin in pigmented fruits and vegetables, protects human HaCaT keratinocytes and mouse skin against UVB mediated oxidative stress and apoptosis. *J Invest Dermatol*. 2007; 127: 222-232.
- Aguilar-Garcia C, Gavino G, Baragaño-Mosqueda M, Hevia P, Gavino VC. Correlation of tocopherol, tocotrienol,  $\gamma$ -oryzanol and total polyphenol

- content in rice bran with different antioxidant capacity assays. *Food Chem.* 2007; 102(4): 1228-1232.
- Aidoo KE, Nout MJR, Sarkar PK. Occurrence and function of yeasts in Asian indigenous fermented foods. *FEMS Yeast Res.* 2006; 6: 30-39.
- Akim A, Ling LC, Rahmat A, Zakaria ZA. Antioxidant and anti-proliferative activities of Roselle juice on Caov-3, MCF-7, MDA-MB-231 and HeLa cancer cell lines. *Afr J Pharm Pharmacol.* 2011; 5(7): 957-965.
- Amarowicz R, Pegg RB, Rahimi-Moghaddam P, Barl B, Weil JA. Free-radical scavenging capacity and antioxidant activity of selected plant species from the Canadian prairies. *Food Chem.* 2004; 84: 551-562.
- AOAC. Official methods of analysis (17th ed.). Gaithersburg, MD: Association of Official Analytical Chemists. 2000.
- Azzi A, Davies KJA, Kelly F. Free radical biology-terminology and critical thinking. *FEBS Lett.* 2004; 558: 3-6.
- Baars AJ. Biotransformation of xenobiotics in *Drosophila melanogaster* and its relevance for mutagenicity testing. *Drug Metab Rev.* 1980; 11:191-221.
- Bagchi D, Sen CK, Bagchi M, Atalay M. Anti-angiogenic, antioxidant, and anti-carcinogenic properties of a novel anthocyanin-rich berry extract formula. *Biochemistry Mosc.* 2004; 69: 75-80.
- Baier SK, Guthrie BD, Elmore DL, Smith SA, Lendon CA, Muroski AR, Aimutis W. Influence of extrusion on protein conformation and shelf-life extension of nutritional bars. Paper presented at the international symposium on the properties of water X, 2-7 September, Bangkok, Thailand. 2007
- Beuchat LR. Microbial stability as affected by water activity. *Cereal Food World.* 1981; 26: 345-349.
- Buglass AJ. Alcoholic fermentation. In: Buglass AJ (ed.), *Handbook of alcoholic beverages: technical, analytical and nutritional aspects*. United States: John Wiley and Sons. 2011; pp 73-74.
- Canas BJ, Havery DC, Robinson LR, Sullivan MP, Joe FL Jr, Diachenko GW. Ethyl carbamate levels in selected fermented foods and beverages. *J. AOAC Int.* 1989; 72: 873-6.

- Castro FDD. Cereal bars with soy protein and wheat germ, physicochemical characteristics and texture during the storage. *Arch Latinoam Nutr.* 2005; 55(3): 299
- Champagne ET, Wood DF, Juliano BO, Bechtel DB. The rice grain and its gross composition. In: Champagne ET (ed.), *Rice*. American association of cereal chemistry. St. Paul, USA. 2004; pp 77-107
- Chaudhary RC. Speciality rices of the world: Effect of WTO and IPR on its production trend and marketing. *J Food Agr Environ.* 2003; 1(2): 34-41.
- Chen CC, Shih YC, Chiou PWS, Yu B. Evaluating nutritional quality of single stage- and two stage-fermented soybean meal. *Asian-Aust J Anim Sci.* 2010; 23(5): 598 - 606.
- Chen PN, Kuo WH, Chiang CL, Chiou HL, Hsieh YS, Chu SC. Black rice anthocyanins inhibit cancer cells invasion via repressions of MMPs and u-PA expression. *Chem Biol Interact.* 2006; 163: 218-229.
- Cheng Q, Bai X, Wang Y. Isolation and fermentation condition of milk clotting enzyme producing strain from glutinous rice wine. *Chin J Biotechnol.* 2008; 24(6): 999-1003.
- Cheriot S, Billaud C, Pöchtrager S, Wagner KH, Nicolas J. A comparison study between antioxidant and mutagenic properties of cysteine glucose-derived Maillard reaction products and neoformed products from heated cysteine and hydroxymethylfurfural. *Food Chem.* 2009; 114: 132-138.
- Choi JH, Choi CY, Lee KJ, Hwang YP, Chung YC, Jeong HG. Hepatoprotective effects of an anthocyanins fraction from purple-fleshed sweet potato against acetaminophen-induced liver damage in mice. *J Med Food.* 2009; 12: 320-326.
- Crout JR. Drug product containing urethane. Opportunity to heating on proposal to withdraw approval of new drug application for profenil injection. U.S. Federal Registry Document No. 76-5599; 1976.
- Daglia M, Papetti A, Aceti C, Sordelli B, Gregotti C, Gazzani G. Isolation of high molecular weight components and contribution to the protective activity of coffee against lipid peroxidation in a rat liver microsome system. *J Agric Food Chem.* 2008; 56(24): 11653-11660.

- Dahl GA, Miller JA, Miller EC. Vinyl carbamate as a promutagen and a more carcinogenic analog of ethyl carbamate. *Cancer Res.* 1978; 38: 3793-3804.
- Daker M, Abdullah N, Vikineswary S, Goh PC, Kuppusamy UR. Antioxidant from maize and maize fermented by *Marasmiellus* sp. as stabiliser of lipid-rich foods. *Food Chem.* 2008; 107: 1092-1098.
- De Flora S, Astengo M, Serra D, Bennicelli C. Inhibition of urethane-induced lung tumors in mice by dietary N-acetylcysteine. *Cancer Lett.* 1986; 32: 235-241.
- de Llanos R, Querol A, Peman J, Gobernado M, Fernandez-Espinar MT. Food and probiotic strains from the *Saccharomyces cerevisiae* species as a possible origin of human systemic infections. *Int J Food Microbiol.* 2006; 110(3): 286-290.
- Dordević TM, Šiler-Marinković SS, Dimitrijević-Branković SI. Effect of fermentation on antioxidant properties of some cereals and pseudo cereals. *Food Chem.* 2010; 119: 957-963.
- Erukainure OL, Oke OV, Adenekan SO, Ajiboye JA. Antioxidant activities, total phenolic and flavonoid levels of watermelon rinds subjected to *saccharomyces cerevisiae* solid media fermentation. *World Journal of Ferm. Tech and Bioeng.* 2011; 2: 11-16.
- Faist V, Lindenmeier M, Geisler C, Erbersdobler HF, Hofmann T. Influence of molecular weight fractions isolated from roasted malt on the enzyme activities of NADPH-cytochrome c-reductase and glutathione-S-transferase in Caco-2 cells. *J Agric Food Chem.* 2002; 50(3): 602-606.
- FAO. Rice and human nutrition. International year of rice 2004. Available from: <http://www.fao.org/rice2004/en/f-sheet/factsheet3.pdf>.
- Feng R, Ni HM, Wang SY, Tourkova IL, Shurin MR, Harada H, Yin XM. Cyanidin-3-rutinoside, a natural polyphenol antioxidant, selectively kills leukemic cells by induction of oxidative stress. *J Biol Chem.* 2007; 282: 13468-13476.
- Floyd RA. Neuroinflammatory processes are important in neurodegenerative diseases: an hypothesis to explain the increased formation of reactive

- oxygen and nitrogen species as major factors involved in neurodegenerative disease development. *Free Radical Biol Med.* 1999; 26(9-10): 1346-1355.
- Frei H, Würgler FE. Statistical methods to decide whether mutagenicity test data from *Drosophila* assay indicate a positive, negative or inconclusive result. *Mutat Res.* 1988; 203: 297-308.
- Frölich A, Würgler FE. *Drosophila* wing-spot test: improved detectability of genotoxicity of polycyclic aromatic hydrocarbons. *Mutat Res* 1990a; 234: 71-80.
- Frolich A, Würgler FE. Genotoxicity of ethyl carbamate in the *Drosophila* wing-spot test: dependence on genotype-controlled metabolic capacity. *Mutat Res.* 1990; 244: 201-208.
- Frölich A, Würgler FE. Genotoxicity of ethyl carbamate in the *Drosophila* wing spot test: dependence on genotype-controlled metabolic capacity. *Mutat Res* 1990b; 244: 201-8.
- Fukumoto LR, Mazza G. Assessing antioxidant and pro-oxidant activity of phenolic compounds. *J Agric Food Chem.* 2000; 48(8): 3597-3604.
- Gliszczynska-Swiglo A, Tyrakowska, B. Quality of commercial apple juices evaluated on the basis of the polyphenol content and the TEAC antioxidant activity. *J Food Sci.* 2003; 68: 1844-1849.
- Gonzalez-Castejon M, Rodriguez-Casado A. Pharmacological Dietary phytochemicals and their potential effects on obesity: a review. *Pharmacol Res.* 2011; 64: 438-455.
- Graf U, Würgler FE, Katz AI, Frei H, Juon H, Hall CB. Somatic mutation and recombination test in *Drosophila melanogaster*. *Environ Mutagen.* 1984; 6: 153-88.
- Graf D, van Schaik N. Improved high bioactivation cross for the wing somatic mutation and recombination test in *Drosophila melanogaster*. *Mutat Res.* 1992; 271: 59-67.
- Graf U, Abraham SK, Rincon JG, Würgler FE. Antigenotoxicity studies in *Drosophila melanogaster*. *Mutat Res.* 1998; 402: 203-209.
- Graf U, Singer D. Somatic mutation and recombination test in *Drosophila*

- melanogaster* (wing spot test): effects of extracts of airborne particulate matters from fire-exposed and non fire-exposed building ventilation filters. *Chemosphere*. 1989; 19: 1094-1097.
- Graf U, van Schaik N. Improved high bioactivation cross for the wing somatic mutation and recombination test in *Drosophila melanogaster*. *Mutat Res*. 1992; 271: 59-67.
- Graf U, Würgler FE, Katz AI, Frei H, Juon H, Hall CB. Somatic mutation and recombination test in *Drosophila melanogaster*. *Environ Mutagen*. 1984; 6: 153-188.
- Griffin SP, Bhagooli R. Measuring antioxidant potential in corals using the FRAP assay. *J Exp Marine Bio Ecol*. 2004; 302: 201-211.
- Guengerich FP, Kim DH, Iwasaki M. Role of human cytochrome P-450 IIE1 in the oxidation of many low molecular weight cancer suspects. *Chem Res Toxicol*. 1991; 4: 168-179.
- Guo C, Yang J, Wei J, Li Y, Xu J, Jiang Y. Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay. *Nutr Res*. 2003; 23: 1719-1726.
- Gutteridge JMC, Mitchell J. Redox imbalance in the critically ill. *Br Med Bull* 1999; 55(1): 49-75.
- Hallstrom I, Blanck A, Atuma S. Genetic variation in cytochrome P-450 and xenobiotic metabolism in *Drosophila melanogaster*. *Biocheml Pharmacol*. 1984; 33: 13-20.
- Hamss RE, Idaomara M, Moraga AA, Serrano AM. Antimutagenic properties of bell and black peppers. *Food ChemToxicol*. 2003; 41: 41-47.
- Handoyo T, Morita N. Structural and functional properties of fermented soybean (tempeh) by using *Rhizopus oligosporus*. *Int J Food Prop*. 2006; 9(2): 347-355.
- Harborne JB, Grayer RJ. The anthocyanins. In: Harborne JB (ed.), *The Flavonoids: advances in research since 1980*. London: Chapman and Hall. 1988; pp 1-20.
- Hayase F, Usui T, Watanabe H. Chemistry and some biological effects of model melanoidins and pigments as Maillard intermediates. *Mol Nutr Food Res*.

- 2006; 50: 1171-1179.
- HHS/OS/OPHS. Solicitation of written comments on proposed definition of bioactive food components. U.S. Dept. of Health and Human Services/Office of the Secretary/Office of Public Health and Safety, Washington, DC. Fed. Reg. 2004; 69: 55821-55822.
- Hoffler U, Dixon D, Peddada S, Ghanayem BI. Inhibition of urethane-induced genotoxicity and cell proliferation in CYP2E1-null mice. *Mutat Res.* 2005; 572: 58-72.
- Hoffler U, El-masri HA, Ghanayem BI. Cytochrome P450 2E1 (cyp2E1) is the principal enzyme responsible for urethane metabolism: comparative studies using cyp 2E1-null and wild type mice. *J Pharmacol Exp Ther.* 2003; 305: 557-564.
- Hou DX, Kai K, Li JJ, Lin S, Terahara N, Wakamatsu M, Fujii M, Young MR, Colburn N. Anthocyanidins inhibit activator protein 1 activity and cell transformation: structure-activity relationship and molecular mechanisms. *Carcinogenesis.* 2004; 25: 29-36.
- Hou DX. Potential mechanisms of cancer chemoprevention by anthocyanins. *Curr Mol Med.* 2003; 3: 149-159.
- Hsieh ML, Chou CC. Mutagenicity and antimutagenic effect of soymilk fermented with lactic acid bacteria and bifidobacteria. *Int J Food Microbiol.* 2006; 111: 43-47.
- Hu C, Zawistowski J, Ling W, Kitts DD. Black rice (*Oryza sativa* L. *indica*) pigmented fraction suppresses both reactive oxygen species and nitric oxide in chemical and biological model systems. *J Agric Food Chem.* 2003; 51(18): 5271-5277.
- Hung YH, Huang HY, Chou CC. Mutagenic and antimutagenic effects of methonal extracts of unfermented and fermented black soybeans. *Int J Food Microbiol.* 2007; 118: 62-68.
- Hwang IG, Kim HY, Woob KS, Lee J, Jeong HS. Biological activities of Maillard reaction products (MRPs) in a sugar-amino acid model system. *Food Chem.* 2011; 126: 221-227.
- Hwang YP, Choi JH, Choi JM, Chung YC, Jeong HG. Protective mechanisms of

- anthocyanins from purple sweet potato against tert-butyl hydroperoxide-induced hepatotoxicity. *Food Chem Toxicol.* 2011a; 49: 2081-2089.
- Hwang YP, Choi JH, Yun HJ, Han EH, Kim HG, Kim JY, Park B.H, Khanal T, Choi JM, Chung YC, Jeong HG. Anthocyanins from purple sweet potato attenuate dimethylnitrosamine-induced liver injury in rats by inducing Nrf2-mediated antioxidant enzymes and reducing COX-2 and iNOS expression. *Food Chem Toxicol.* 2011b; 49: 93-99.
- Hyogo A, Kobayashi T, Garcia del Saz E, Seguchi H. Antioxidant effects of protocatechuic acid, ferulic acid, and caffeic acid in human neutrophils using a fluorescent substance. *Int J Morphol.* 2010; 28(3): 911-920.
- Hyun JW, Chung HS. Cyanidin and malvidin from *Oryza sativa* cv. Heugjinjubyeo mediate cytotoxicity against human monocytic leukemia cells by arrest of G(2)/M phase and induction of apoptosis. *J Agric Food Chem.* 2004; 52(8): 2213-2217.
- IARC. Urethane. In: IARC monographs on the evaluation of carcinogenic risk of chemicals to humans, some anti-thyroid and related substances, nitrofurans and industrial chemicals. Lyon, France: International Agency for Research on Cancer. 1974; 7: 111-140.
- Ichikawa H, Ichiyanagi T, Xu B, Yoshii Y, Nakajima M, Konishi T. Antioxidant activity of anthocyanin extract from purple black rice. *J Med Food.* 2001; 4(4): 211-218.
- IFIC. Functional foods attitudinal research. International Food Information Council, Washington, DC. 2002. Available from: <http://ific.org/proactive/newsroom/release.vtml?id=20761>.
- Ingelman-Sundberg M, Johansson I, Yin H, Terelius Y, Eliasson E, Clot P, Albano E. Ethanol-inducible cytochrome P4502E1: genetic polymorphism, regulation, and possible role in the etiology of alcohol-induced liver disease. *Alcohol.* 1993; 10: 447-452.
- Jaeger H, Janositz A, Knorr D. The Maillard reaction and its control during food processing. The potential of emerging technologies. *Pathol Biol.* 2010; 58: 207-213.
- James MJ, Ursin VM, Cleland LG. Metabolism of stearidonic acid in human subjects:



- comparison with the metabolism of other n-3 fatty acids. *Am J Clin Nutr.* 2003; 77: 1140-1145.
- Ji LL. Free radicals and exercise: implication in health and fitness. *J Exerc Sci Fit.* 2003; 1(1): 15-22.
- Jing H, Kitts DD. Chemical and biochemical properties of casein-sugar Maillard reaction products. *Food Chem Toxicol.* 2000; 40: 1007-1015.
- Ju HK, Cho EJ, Jang ME, Lee YY, Hong SS, Park JH, Kwon SW. Characterization of increased phenolic compounds from fermented Bokbunja (*Rubus coreanus* Miq.) and related antioxidant activity. *J Pharmaceut Biomed Anal.* 2009; 49: 820-827.
- Karthikeyan R, Manivasagam T, Anantharaman P, Balasubramanian T, Somasundaram ST. Chemopreventive effect of *Padina boergesenii* extracts on ferric nitrilotriacetate (Fe-NTA)-induced oxidative damage in Wistar rats. *J Appl Phycol.* 2011; 23(2): 57-263.
- Kemper RA, Myers SR, Hurst HE. Detoxification of vinyl carbamate epoxide by glutathione: evidence for participation of glutathione-S-transferase in metabolism of ethyl carbamate. *Toxicol Appl Pharmacol.* 1995; 135(1): 110-118.
- Koide T, Kamei H, Hashimoto Y, Kojima T, Hasegawa M. Antitumor effect of hydrolyzed anthocyanin from grape rinds and red rice. *Cancer Biother. Radiopharm.* 1996; 11(4): 273-277.
- Kotanidou A, Chol AM, Winchurch RA, Otterbein L, Fessler HE. Urethan anesthesia protects rats against lethal endotoxemia and reduces TNF-alpha release. *J Appl Physiol.* 1996; 81(5): 2305-2311.
- Kraiket S. Improvement of fermented Thai rice noodle (khanomjeen) with natural colors as a source of antimutagen and antioxidant. Master thesis, Mahidol University, Bangkok, Thailand. 2007.
- Kuhn MA. Oxygen Free Radicals and Antioxidants: an overview of how antioxidants protect the body from disease. *Am J Nurs.* 2003; 103(4): 58-62.
- Kurata N, Hurst HE, Benz FW, Kemper RA, Waddell WJ. Studies on inhibition and induction of metabolism of ethyl carbamate by acetone and related

- compounds. Evidence for metabolism by cytochrome P450. *Drug Metab Dispos.* 1991a; 19: 388-393.
- Kurata N, Hurst HE, Kemper RA, Waddell WJ. Studies on induction of metabolism of ethyl carbamate in mice by ethanol. *Drug Metab Dispos.* 1991b; 19: 239-240.
- Kwak CS, Lee MS, Chul S. Higher antioxidant properties of Chungkookjang, a fermented soybean paste, may be due to increased aglycone and malonylglycoside isoflavone during fermentation. *Nutr Res.* 2007; 27: 719-727.
- Lee IH, Chou CC. Distribution profiles of isoflavone isomers in black bean koji prepared with various filamentous fungi. *J Agric Food Chem.* 2006; 54(4): 1309-1314.
- Lee IH, Hung YH, Chou CC. Solid-state fermentation with fungi to enhance the antioxidative activity, total phenolic and anthocyanin contents of black bean. *Int J Food Microbiol.* 2008; 121: 150-156.
- Lee IH, Hung YH, Chou CC. Total phenolic and anthocyanin contents, as well as antioxidant activity, of black bean koji fermented by *Aspergillus awamori* under different culture conditions. *Food Chem.* 2007; 104: 936-942.
- Leithauser MT, Liem A, Stewart BC, Miller EC, Miller JA. 1,N6- ethenoadenosine formation, mutagenicity and murine tumor induction as indicators of the generation of an electrophilic epoxide metabolite of the closely related carcinogens ethyl carbamate (urethane) and vinyl carbamate. *Carcinogenesis.* 1990; 11(3): 463-473.
- Liang CH, Syu JL, Mau JL. Antioxidant properties of solid-state fermented adlay and rice by *Phellinus linteus*. *Food Chem.* 2009; 116: 841-845.
- Lieber CS. Biochemical and molecular basis of alcohol-induced injury to liver and other tissues. *N Engl J Med.* 1988; 319(25): 1639-1650.
- Lieber CS. Interaction of alcohol with other drugs and nutrients. Implication for the therapy of alcoholic liver disease. *Drugs.* 1990; 40: 23-44.
- Lin CH, Chou CC. Suppression on the mutagenicity of 4-nitroquinoline-N-oxide by the methanol extracts of soybean koji prepared with various filamentous fungi. *Int J Food Microbiol.* 2006; 110: 43-47.

- Lindenmeier M, Faist V, Hofmann T. Structural and functional characterization of pronyl-lysine, a novel protein modification in bread crust melanoidins showing in vitro antioxidative and phase I/II enzyme modulating activity. *J Agric Food Chem.* 2002; 50(24): 6997-7006.
- Ling WH, Cheng QX, Ma J, Wang T. Red and black rice decrease atherosclerotic plaque formation and increase antioxidant status in rabbits. *J Nutr.* 2001; 131(5): 1421-1426.
- Liu RH. Potential synergy of phytochemicals in cancer prevention: mechanism of action. *J Nutr.* 2004; 134: 3479S-3485S.
- Liu RH. Whole grain phytochemicals and health. *J Cereal Sci.* 2007; 46(3): 207-219.
- Lobato LP, Iakmiu Camargo Pereira AE, Lazaretti MM, Barbosa DS, Carreira CM, Mandarino JM, Grossmann MV. Snack bars with high soy protein and isoflavone content for use in diets to control dyslipidaemia. *Int J Food Sci Nutr.* 2011; 63(1): 49-58.
- Lotong, N. Seed inoculum and their production technology (in Thai) (2<sup>nd</sup> ed.). Bangkok: Funny Publishing. 1992; pp 18.
- Loveday SM, Hindmarsh JP, Creamer LK, Singh H. Physicochemical changes in a model protein bar during storage. *Food Res Int.* 2009; 42: 798-806.
- Manosroi A, Ruksiriwanich W, Kietthanakorn B, Manosroi W, Manosroi J. Relationship between biological activities and bioactive compounds in the fermented rice sap. *Food Res Int.* 2011; 44(9): 2757-2765.
- Martin MA, Ramos S, Mateos R, Rufian-Henares JA, Morales FJ, Bravo L, Luis G. Biscuit melanoidins of different molecular masses protect human HEPG2 cells against oxidative stress. *J Agric Food Chem.* 2009; 57(16): 7250-7258.
- Mazza G, Miniati E. Anthocyanins in fruits, vegetables and grains. CRC Press, Boca Raton. 1993.
- McGhie TK, Walton MC. The bioavailability and absorption of anthocyanins: towards a better understanding. *Mol Nutr. Food Res.* 2007; 51: 702-713.
- Mendonc CX Jr, Almeida CRM, Mori AV, Watanabe C. Effect of dietary vitamin A on egg yolk retinol and tocopherol levels. *J Appl Poult Res.* 2002; 11: 373-378.

- Mergener M, Martins MR, Antunes MV, Calice da Silva C, Lazzaretti C, Fontanive TO, Suyenaga ES, Ardenghi PV, Maluf SW, Gamaro GD. Oxidative stress and DNA damage in older adults that do exercises regularly. *Clin Biochem.* 2009; 42: 1648-1653.
- Meyers KJ, Watkins CB, Pritts MP, Liu RH. Antioxidant and antiproliferative activities of strawberries. *J Agric Food Chem.* 2003; 51(23): 6887-6892.
- Michalska A, Amigo-Benavent M, Zielinski H, del Castillo MD. Effect of bread making on formation of Maillard reaction products contributing to the overall antioxidant activity of rye bread. *J Cereal Sci.* 2008; 48: 123–132.
- Miller JA, Miller EC. The metabolic activation and nucleic acid adducts of naturally occurring carcinogens: recent results with ethyl carbamate and the spice flavors safrole and estragole. *Br J Cancer.* 1983; 48(1): 1-15.
- Mirvish SS. The carcinogenic action and metabolism of urethane and N hydroxyl urethan. *Adv Cancer Res.* 1968; 11: 1-42.
- Mishra DK, Dolan KD, Yang L. Confidence intervals for modeling anthocyanin retention in grape pomace during nonisothermal heating. *J Food Sci.* 2008; 73: E9-E15.
- Miwa M, Watanabe T, Kawasumi T, Hayase F. Protective effects of melanoidins derived from soy sauce and soy paste on NO induced DNA damage. *Food Sci Technol Res.* 2002; 8 (3): 231-234.
- Moktan B, Saha J, Sarkar PK. Antioxidant activities of soybean as affected by *Bacillus*-fermentation to kinema. *Food Res Int.* 2008; 41: 586-593.
- Mori I, Yasuhara K, Hayashi SM, Nonoyama T, Nomura T, Mitsumori K. Carcinogen dose-dependent variation in the transgene mutation spectrum in urethane-induced lung tumors in transgenic mice carrying the human prototype c-Ha-ras gene. *Cancer Lett.* 2000; 153: 199-209.
- Mukherjee PK, Marcheselli VL, Serhan CN, Bazan NG. Neuroprotecin D1: a docosahexanoic acid-derived docosatriene protects human retinal pigment epithelial cells from oxidative stress. *Proc Natl Acad Sci USA.* 2004; 101(22): 8491-8496.
- Norlen P, Kitano M, Lindstrom E, Hakanson R. Anaesthetic agents inhibit gastrin-stimulated but not basal histamine release from rat stomach ECL cells. *Br*

- J Pharmacol. 2000; 130(4): 725-730.
- O'Donnell CD. 10 trends in nutraceutical ingredients. Prepared Foods. 2003; 172(3): 89-92.
- Olsson ME, Gustavsson KE, Andersson S, Nilsson A, Duan RD. Inhibition of cancer cell proliferation in vitro by fruit and berry extracts and correlations with antioxidant levels. J Agric Food Chem. 2004; 52(24): 7264-7271.
- Omata Y, Ogawa Y, Saito Y, Yoshida Y, Nik E. Assessment of the antioxidant capacity of a fermented grain food product, Antioxidant Biofactor (AOB), by using pyranine and pyrogallol red as a combined probe. Food Chem. 2009; 114: 429-433.
- Ough CS. Ethyl carbamate in fermented beverages and foods. I. naturally occurring ethyl carbamate. J Agric Food Chem. 1976; 24(2): 323-328.
- Pacher P, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. Physiol Rev. 2007; 87(1): 315-424.
- Papetti A, Daglia M, Aceti C, Quaglia M, Gregotti C, Gazzani G. Isolation of an in vitro and ex vivo antiradical melanoidin from roasted barley. J Agric Food Chem. 2006; 54(4): 1209-1216.
- Park KK, Liem A, Stewart BC, Miller JA. Vinyl carbamate epoxide, a major strong electrophilic, mutagenic and carcinogenic metabolite of vinyl carbamate and ethyl carbamate (urethane). Carcinogenesis. 1993; 14: 441-450.
- Parry J, Su L, Moore J, Cheng Z, Luther M, Rao JN, Wang JY, Yu LL. Chemical compositions, antioxidant capacities, and antiproliferative activities of selected fruit seed flours. J Agric Food Chem. 2006; 54(11): 3773-3778.
- Pitiporn S. Miracle of Thai rice. Matichon Weekly. Bangkok: Matichon Publisher. 2008; 461: pp 92.
- Prior RL, Wu X. Anthocyanins: structural characteristics that result in unique metabolic patterns and biological activities. Free Radic Res. 2006; 40(10): 1014-1028.
- Ratnam DV, Ankola DD, VBhardwaj V, Sahana DK, Ravi Kumar MNV. Role of antioxidants in prophylaxis and therapy: a pharmaceutical perspective. J Controlled Release. 2006; 113(3): 189-207.
- Reen RK, Nines R, Stoner GD. Modulation of N-nitrosomethylbenzylamine

- metabolism by black raspberries in the esophagus and liver of Fischer 344 rats. *Nutr Cancer*. 2006; 54: 47-57.
- Renis M, Calandra L, Scifo C, Tomasello B, Cardile V, Vanella L, Bei R, Fauci LL, Galvano F. Response of cell cycle/stress-related protein expression and DNA damage upon treatment of CaCo2 cells with anthocyanins. *Br J Nutr*. 2008; 100(1): 27-35.
- Rhodes CJ. Toxicology of the human environment: the critical role of free radicals. London; Taylor and Francis. 2000.
- Roberts DB. Basic *Drosophila* care and techniques. In: Roberts DB (ed), *Drosophila: a practical approach*. Oxford: IRL Press. 1986; pp 1-38.
- Rodriguez AR, Tellez GO. Structure-activity relationships of several anisidine and dibenzanthracene isomers in the w/w+ somatic assay of *Drosophila melanogaster*. *Mutat Res*. 2002; 514: 193-200.
- Rufián-Henares JA, Morales FJ. Functional properties of melanoidins: in vitro antioxidant, antimicrobial and antihypertensive activities. *Food Res Int*. 2007; 40: 995-1002.
- Ryland D, Vaisey-Genser M, Arntfield SD, Malcolmson LJ. Development of a nutritious acceptable snack bar using micronized flaked lentils. *Food Res Int*. 2010; 43: 642-649.
- Sadabpodi K, Kangsadalampai K, Tongyong L. Antioxidant activity and antimutagenicity of Hom Nil rice and black glutinous rice. *J Health Res*. 2010; 24: 49-54.
- Sangkitikomol W, Tencomnao T, Rocejanasaroj A. Antioxidant effects of anthocyanins-rich extract from black sticky rice on human erythrocytes and mononuclear leukocytes. *Afr J Biotechnol*. 2010; 9(48): 8222-8229.
- Sarikaya R, Cakir S. Genotoxicity testing of four food preservatives and their combinations in the *Drosophila* wing spot test. *Environ Toxicol Pharmacol*. 2005; 20(3): 424-230.
- Scheibmeir HD, Christensen K, Whitaker SH, Jegaethesan J, Clancy R, Pierce JD. A review of free radicals and antioxidants for critical care nurses. *Intensive Crit Care Nurs*. 2005; 21(1): 24-28.
- Schlatter DB, Lutz WK. The carcinogenic potential of ethyl carbamate (urethane):

- risk assessment at human dietary exposure levels. *Food Chem Toxicol.* 1990; 28: 205-211.
- Schlatter J, Lutz WK. The carcinogenic potential of ethyl carbamate (urethane): risk assessment at human dietary exposure levels. *Food Chem Toxicol.* 1990; 28: 205-211.
- Ścibisz I, Mitek M. Effect of processing and storage conditions on phenolic compounds and antioxidant capacity of high bush blueberry jams. *Pol J Food Nutr Sci.* 2009; 59 (1): 45-52.
- Sharma P, Velu V, Indrani D, Singh RP. Effect of dried guduchi (*Tinospora cordifolia*) leaf powder on rheological, organoleptic and nutritional characteristics of cookies. *Food Res Int.* 2012. In Press, Corrected Proof.
- Shih PH, Yeh CT, Yen GC. Anthocyanins induce the activation of phase II enzymes through the antioxidant response element pathway against oxidative stress induced apoptosis. *J Agric Food Chem.* 2007; 55(23): 9427-9435.
- Singh VP, Sanyal MK, Dubey PC, Sachan N, Kumar V. Chicken snacks as affected by storage conditions under aerobic and vacuum packaging at 30±2°C. *Afr J Food Sci.* 2011; 5(11): 620-625.
- Singletary KW, Jung KJ, Giusti M. Anthocyanin-rich grape extract blocks breast cell DNA damage. *J Med Food.* 2007; 10: 244-251.
- Somoza V, Wenzel E, Lindenmeier M, Grothe D, Erbersdobler HF, Hofmann T. Influence of feeding malt, bread crust, and a pronylated protein on the activity of chemopreventive enzymes and antioxidative defense parameters in vivo. *J Agric Food Chem.* 2005; 53(21): 8176-8182.
- Sompong R, Siebenhandl-Ehn S, Linsberger-Martin G, Berghofer E. Physicochemical and antioxidative properties of red and black rice varieties from Thailand, China and Sri Lanka. *Food Chem.* 2011; 124: 132-140.
- Stoewsand GS, Anderson JL, Munson L. Inhibition by wine of tumorigenesis induced by ethyl carbamate (urethane) in mice. *Food Chem Toxicol.* 1991; 29: 291-295.
- Summa C, McCourt J, Cämmerer B, Fiala A, Probst M, Kun S, Anklam E, Wagner KH. Radical scavenging activity, anti-bacterial and mutagenic effects of

- cocoa bean Maillard reaction products with degree of roasting. *Mol Nutr Food Res.* 2008; 52(3): 342-351.
- Suzuki M, Kimura T, Yamagishi K, Shinmoto H, Yamaki K. Comparison of mineral contents in 8 cultivars of pigmented brown rice. *J Jpn Soc Food Sci Technol* 2004; 51(8): 424-7.
- Syamaladevi RM, Andrews PK, Davies NM, Walters T, Sablani SS. Storage effects on anthocyanins, phenolics and antioxidant activity of thermally processed conventional and organic blueberries. *J Sci Food Agric.* 2012; 92(4): 916-924.
- Trošt K, Golc-Wondra A, Prošek M, Milivojević L. Anthocyanin degradation of blueberry-aronia nectar in glass compared with carton during storage. *J Food Sci.* 2008; 73(8): S405-S411.
- Uggla A, Busk L. Ethyl carbamate (Urethane) in alcoholic beverages and foodstuffs-*a Nordic View*, Nordic Council of Ministers, Copenhagen. 1992.
- Ursin VM. Modification of plant lipids for human health: development of functional land-based omega-3 fatty acids. *J Nutr.* 2003; 133: 4271-4274.
- Valls-Bellés V, Torres M, Muñiz P, Boix L, González-Sanjosé ML, Codoñer-Franch P. The protective effects of melanoidins in adriamycin-induced oxidative stress in isolated rat hepatocytes. *J Sci Food Agric.* 2004; 84(13): 1701-1707.
- Vallverdú-Queralt A, Arranz S, Medina-Remón A, Casals-Ribes I, Lamuela-Ravent RM. Changes in phenolic content of tomato products during storage. *J Agric Food Chem.* 2011; 59(17): 9358-9365.
- Vergara-Valencia N, Granados-Pérez E, Agama-Acevedo E, Tovar J, Ruales J, Bello-Pérez LA. Fibre concentrate from mango fruit: characterization, associated antioxidant capacity and application as a bakery product ingredient. *LWT-Food Sci Technol.* 2007; 40: 722-729.
- Verzelloni E, Tagliazucchi D, Conte A. From balsamic to healthy: traditional balsamic vinegar melanoidins inhibit lipid peroxidation during simulated gastric digestion of meat. *Food Chem Toxicol.* 2010; 48: 2097-2102.
- Vichapong J, Sookserm M, Srijesdaruk V, Swatsitang P, Srijaranai S. High performance liquid chromatographic analysis of phenolic compounds and



- their antioxidant activities in rice varieties. *LWT-Food Sci Technol*. 2010; 43: 1325-1330.
- Vijay P, Vijayvergia R. An antioxidant potential of Hingot and Hadjod. *J B Clin Pharm*. 2010; 1: 33-36.
- Viña J, Gomez-Cabrera MC, Lloret A, Marquez R, Miñana JB, Pallardó FV, Sastre J. Free radicals in exhaustive physical exercise: mechanism of production, and protection by antioxidants. *IUBMB Life*. 2000; 50: 271-277.
- Vipassanatham N, Kangsadalampai K, Tongyonk L. Antimutagenicity of hom nil rice and black glutinous rice using somatic mutation and recombination test. *J Health Res*. 2012; 26: 45-50.
- Vogel EW, Nivard MJM. Performance of 181 chemicals in a *Drosophila* assay predominantly monitoring interchromosomal mitotic recombination. *Mutagenesis*. 1993; 8(1): 57-81.
- Vogel EW, Zijlstra JA. Mechanistic and methodological aspect of chemically induced somatic mutation and recombination in *Drosophila melanogaster*. *Mutat Res*. 1987; 182: 243-264.
- Vogel EW. Evaluation of potential mammalian genotoxins using *Drosophila*: the need for a change in test strategy. *Mutagenesis*. 1987; 2(3): 161-171.
- Waddell WJ, Marlowe C, Pierce WM. Inhibition of the localization of urethane in mouse tissues by ethanol. *Food Chem Toxicol*. 1987; 25(7): 527-531.
- Wang HY, Qian H, Yao WR. Melanoidins produced by the Maillard reaction: structure and biological activity. *Food Chem*. 2011; 128: 573-584.
- Wang HY, Yu XY, Qian H, Yao WR. Separation of antioxidant components from meat flavor and their antioxidant capacities. *Food Science*. 2010; 31(19): 59-62.
- Wang LS, Stoner GD. Anthocyanins and their role in cancer prevention. *Cancer Lett*. 2008; 269: 281-290.
- Wang Q, Han P, Zhang M, Xia M, Zhu H, Ma J, Hou M, Tang Z, Ling W. Supplementation of black rice pigment fraction improves antioxidant and anti-inflammatory status in patients with coronary heart disease. *Asia Pac J Clin Nutr* 2007; 16 (Suppl 1): 295-301.
- Wang SY, Jiao H. Scavenging capacity of berry crops on superoxide radicals,

- hydrogen peroxide, hydroxyl radicals, and singlet oxygen. *J Agric Food Chem.* 2000; 48(11): 5677-5684.
- Wang Y, Zhang M, Mujumdar AS, Influence of green banana flour substitution for cassava starch on the nutrition, color, texture and sensory quality in two types of snacks. *LWT-Food Sci. Technol.* 2012; 47: 175-182.
- Wang YJ, Sheen LY, Chou CC. Storage effects on the content of anthocyanin, mutagenicity and antimutagenicity of black soybean koji. *LWT-Food Sci. Technol.* 2010; 43; 702-707.
- Wen X, Enokizo A, Hattori H, Kobayashi S, Murata M, Homma S. Effect of roasting on properties of the zinc-chelating substance in coffee brews. *J Agric Food Chem.* 2005; 53(7): 2684-2689.
- Wilkinson JT, Clapper ML. Detoxication enzymes and chemoprevention. *Proc Soc Exp Biol Med.* 1997; 216: 192-200.
- Wongpiyachon S. Utilization of rice for wine and whisky. Master thesis. Kasetsart University, Bangkok, Thailand. 1995.
- Wu X, Beecher GR, Holden JM, Haytowitz DB, Gebhardt SE, Prior RL. Concentration of anthocyanins in common foods in the United States and estimation of normal consumption. *J Agric Food Chem.* 2006; 54(11): 4069-4075.
- Würgler FE, Vogel EW, *In vivo* mutagenicity testing using somatic cells of *Drosophila melanogaster*. In: De Serres FJ (ed), Chemical mutagens, principles and methods for their detection. New York: Plenum Press. 1986: pp 1-72.
- Xia M, Ling, WH, Ma J, Kitts DD, Zawistowski J. Supplementation of diets with the black rice pigment fraction attenuates atherosclerotic plaque formation in apolipoprotein E deficient mice. *J Nutr.* 2003; 133: 744-751.
- Yamamoto T, Pierce WM, Hurst HE, Chen D, Waddell WJ. Inhibition of the metabolism of urethane by ethanol. *Drug Metab Dispos.* 1988; 16(3): 355-358.
- Yang J, Paulino R, Janke-Stedronsky S, Abawi F. Free-radical-scavenging activity and total phenols of noni (*Morinda citrifolia* L.) juice and powder in processing and storage. *Food Chem.* 2007; 102: 302-308.

- Yawadio R, Tanimori S, Morita N. Identification of phenolic compounds isolated from pigmented rices and their aldose reductase inhibitory activities. *Food Chem.* 2007; 101: 1616-1625.
- Zawistowski J, Kopec A, Kitts DD. Effects of a black rice extract (*Oryza sativa* L. *indica*) on cholesterol levels and plasma lipid parameters in Wistar Kyoto rats. *J Funct Food.* 2009; 1: 50-56.
- Zimmerli B, Schlatter J. Ethyl carbamate: analytical methodology, occurrence, formation, biology activity and risk assessment. *Mutat Res.* 1991; 259: 325-530.