

REFERENCES

- Alexander NB, Schultz AB, Warwick DN. Rising from a chair: effects of age and functional ability on performance biomechanics. **J Gerontol** 1991; 46: 91-8.
- Amatachaya S, Thaweewannakij T, Adirek-udomrat J, Siritaratiwat W. Factors related to obstacle crossing in independent ambulatory patients with spinal cord injury. **J Spinal Cord Med** 2010; 33: 144-9.
- American Spinal Injury Association. International standards for neurological classifications of spinal cord injury. Revised ed. Chicago, Ill: **American Spinal Injury Association** 2000:1-23.
- Bahrami F, Riener R, Jabedar-Maralani P, Schmidt G. Biomechanical analysis of sit-to-stand transfer in healthy and paraplegic subjects. **Clin Biomech (Bristol, Avon)** 2000; 15: 123-33.
- Barbeau H, Elashoff R, Deforge D, Ditunno J, Saulino M, Dobkin BH. Comparison of speeds used for the 15.2-meter and 6-minute walks over the year after an incomplete spinal cord injury: the SCILT Trial. **Neurorehabil Neural Repair** 2007; 21: 302-6.
- Batani H, Maki BE. Assistive devices for balance and mobility: benefits, demands, and adverse consequences. **Arch Phys Med Rehabil** 2005; 86: 134-45.
- Behrman AL, Harkema SJ. Locomotor training after human spinal cord injury: a series of case studies. **Phys Ther** 2000; 80: 688-700.
- Belgen B, Beninato M, Sullivan PE, Narielwalla K. The association of balance capacity and falls self-efficacy with history of falling in community-dwelling people with chronic stroke. **Arch Phys Med Rehabil** 2006; 87: 554-61.
- Bennett L, Murray MP, Murphy EF, Sowell TT. Locomotion assistance through cane impulse. **Int J Rehabil Res** 1979; 2: 519.
- Bohannon RW. Alternatives for measuring knee extension strength of the elderly at home. **Clin Rehabil** 1998; 12: 434-40.
- _____, Smith J, Hull D, Palmeri D, Barnhard R. Deficits in lower extremity muscle and gait performance among renal transplant candidates. **Arch Phys Med Rehabil** 1995; 76: 547-51.

- Brooks D, Davis AM, Naglie G. Validity of 3 physical performance measures in inpatient geriatric rehabilitation. **Arch Phys Med Rehabil** 2006; 87: 105-10.
- Brotherton SS, Krause JS, Nietert PJ. Falls in individuals with incomplete spinal cord injury. **Spinal Cord** 2007; 45: 37-40.
- Brunt D, Greenberg B, Wankadia S, Trimble MA, Shechtman O. The effect of foot placement on sit to stand in healthy young subjects and patients with hemiplegia. **Arch Phys Med Rehabil** 2002; 83: 924-9.
- Burns SP, Golding DG, Rolle WA, Jr., Graziani V, Ditunno JF, Jr. Recovery of ambulation in motor-incomplete tetraplegia. **Arch Phys Med Rehabil** 1997; 78: 1169-72.
- Cesari M, Kritchevsky SB, Penninx BW, Nicklas BJ, Simonsick EM, Newman AB, et al. Prognostic value of usual gait speed in well-functioning older people--results from the Health, Aging and Body Composition Study. **J Am Geriatr Soc** 2005; 53: 1675-80.
- Chandler JM, Duncan PW, Kochersberger G, Studenski S. Is lower extremity strength gain associated with improvement in physical performance and disability in frail, community-dwelling elders? **Arch Phys Med Rehabil** 1998; 79: 24-30.
- Cheng PT, Liaw MY, Wong MK, Tang FT, Lee MY, Lin PS. The sit-to-stand movement in stroke patients and its correlation with falling. **Arch Phys Med Rehabil** 1998; 79: 1043-6.
- Chou CY, Chien CW, Hsueh IP, Sheu CF, Wang CH, Hsieh CL. Developing a short form of the Berg Balance Scale for people with stroke. **Phys Ther** 2006; 86: 195-204.
- Collen FM, Wade DT, Bradshaw CM. Mobility after stroke: reliability of measures of impairment and disability. **Int Disabil Stud** 1990; 12: 6-9.
- Crozier KS, Cheng LL, Graziani V, Zorn G, Herbison G, Ditunno JF, Jr. Spinal cord injury: prognosis for ambulation based on quadriceps recovery. **Paraplegia** 1992; 30: 762-7.
- Csuka M, McCarty DJ. Simple method for measurement of lower extremity muscle strength. **Am J Med** 1985; 78: 77-81.
- Cunningham DA, Paterson DH, Himann JE. Determinants of independence in the elderly. **Can J Appl Physiol** 1993; 18: 243-54

- Curt A , Dietz V. Ambulatory capacity in spinal cord injury: significance of somatosensory evoked potentials and ASIA protocol in predicting outcome. **Arch Phys Med Rehabil** 1997; 78: 39-43.
- Dawson J, Shamley D, Jamous MA. A structured review of outcome measures used for the assessment of rehabilitation interventions for spinal cord injury. **Spinal Cord** 2008; 46: 768-80.
- De Groot IB, Bussmann HJ, Stam HJ, Verhaar JA. Small increase of actual physical activity 6 months after total hip or knee arthroplasty. **Clin Orthop Relat Res** 2008; 466: 2201-8.
- Dean CM, Shepherd RB. Task-related training improves performance of seated reaching tasks after stroke. A randomized controlled trial. **Stroke** 1997; 28: 722-8.
- Ditunno JF, Jr. American spinal injury standards for neurological and functional classification of spinal cord injury: past, present and future. 1992 Heiner Sell Lecture of the American Spinal Injury Association. **J Am Paraplegia Soc** 1994; 17: 7-11.
- _____, Jr., Ditunno PL, Graziani V, Scivoletto G, Bernardi M, Castellano V, et al. Walking index for spinal cord injury (WISCI): an international multicenter validity and reliability study. **Spinal Cord** 2000; 38: 234-43.
- Dobkin B, Apple D, Barbeau H, Basso M, Behrman A, Deforge D, et al. Weight-supported treadmill vs over-ground training for walking after acute incomplete SCI. **Neurology** 2006; 66: 484-93.
- Drabsch T, Lovenfosse J, Fowler V, Adams R, Drabsch P. Effects of task-specific training on walking and sit-to-stand after total hip replacement. **J Physiother** 1998; 44: 193-98.
- Eriksrud O, Bohannon RW. Relationship of knee extension force to independence in sit-to-stand performance in patients receiving acute rehabilitation. **Phys Ther** 2003; 83: 544-51.
- Finch E, Brooks D, Stratford P, Mayo N. **Physical Rehabilitation Outcome Measures: A guide to enhanced clinical decision making**. 2nd ed. Hamilton: BC Decker; 2002.

- Graham JE, Ostir GV, Fisher SR, Ottenbacher KJ. Assessing walking speed in clinical research: a systematic review. **J Eval Clin Pract** 2008; 14: 552-62.
- Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. **J Gerontol** 1994; 49: 85-94.
- Haisma JA, Post MW, van der Woude LH, Stam HJ, Bergen MP, Sluis TA, et al. Functional independence and health-related functional status following spinal cord injury: a prospective study of the association with physical capacity. **J Rehabil Med** 2008; 40: 812-8.
- Hardy SE, Perera S, Roumani YF, Chandler JM, Studenski SA. Improvement in usual gait speed predicts better survival in older adults. **J Am Geriatr Soc** 2007; 55: 1727-34.
- Headley S, Germain M, Mailloux P, Mulhern J, Ashworth B, Burris J, et al. Resistance training improves strength and functional measures in patients with end-stage renal disease. **Am J Kidney Dis** 2002; 40: 355-64.
- Holden MK, Gill KM, Magliozzi MR. Gait assessment for neurologically impaired patients. Standards for outcome assessment. **Phys Ther** 1986; 66: 1530-9.
- _____, Gill KM, Magliozzi MR, Nathan J, Piehl-Baker L. Clinical gait assessment in the neurologically impaired. Reliability and meaningfulness. **Phys Ther** 1984; 64: 35-40.
- Hornby TG, Zemon DH, Campbell D. Robotic-assisted, body-weight-supported treadmill training in individuals following motor incomplete spinal cord injury. **Phys Ther** 2005; 85: 52-66.
- Howe T, Oldham J. Functional tests in elderly osteoarthritic subjects: variability of performance. **Nurs Stand** 1995; 9: 35-8.
- Hughes C, Osman C, Woods AK. Relationship among performance on stair ambulation, functional reach, and Timed Up and Go tests in older adults. **Issues Aging** 1998; 21: 18-22.
- Hughes MA, Schenkman ML. Chair rise strategy in the functionally impaired elderly. **J Rehabil Res Dev** 1996; 33: 409-12.

- Hughes MA, Weiner DK, Schenkman ML. Chair rise strategies in the elderly. **Clin Biomech** 1994; 9: 187-192.
- Jackson AB, Carnel CT, Ditunno JF, Read MS, Boninger ML, Schmeler MR, et al. Outcome measures for gait and ambulation in the spinal cord injury population. **J Spinal Cord Med** 2008; 31: 487-99.
- Janssen WG, Bussmann HB, Stam HJ. Determinants of the sit-to-stand movement: a review. **Phys Ther** 2002; 82: 866-79.
- Kim CM, Eng JJ, Whittaker MW. Level walking and ambulatory capacity in persons with incomplete spinal cord injury: relationship with muscle strength. **Spinal Cord** 2004; 42: 156-62.
- Koh ES, Williams AJ, Povlsen B. Upper-limb pain in long-term poliomyelitis. **QJM** 2002; 95: 389-95.
- Krawetz P, Nance P. Gait analysis of spinal cord injured subjects: effects of injury level and spasticity. **Arch Phys Med Rehabil** 1996; 77: 635-8.
- Lapointe R, Lajoie Y, Serresse O, Barbeau H. Functional community ambulation requirements in incomplete spinal cord injured subjects. **Spinal Cord** 2001; 39: 327-35.
- LeBrasseur NK, Sayers SP, Ouellette MM, Fielding RA. Muscle impairments and behavioral factors mediate functional limitations and disability following stroke. **Phys Ther** 2006; 86: 1342-50.
- Leroux A, Fung J, Barbeau H. Postural adaptation to walking on inclined surfaces: II. Strategies following spinal cord injury. **Clin Neurophysiol** 2006; 117: 1273-82.
- Lin YC, Davey RC, Cochrane T. Tests for physical function of the elderly with knee and hip osteoarthritis. **Scand J Med Sci Sports** 2001; 11: 280-6.
- Lipsitz LA, Jonsson PV, Kelley MM, Koestner JS. Causes and correlates of recurrent falls in ambulatory frail elderly. **J Gerontol** 1991; 46: M114-22.
- Lord SR, Murray SM, Chapman K, Munro B, Tiedemann A. Sit-to-stand performance depends on sensation, speed, balance, and psychological status in addition to strength in older people. **J Gerontol A Biol Sci Med Sci** 2002; 57: M539-43.
- Lusardi MM, Pellecchia GL, Schulman M. Functional performance in community living older adults. **J Geriatr Phys Ther** 2003; 26:14-22.

- Mancini M, Horak FB. The relevance of clinical balance assessment tools to differentiate balance deficits. **Eur J Phys Rehabil Med** 2010; 46: 239-48.
- Maynard FM, Jr., Bracken MB, Creasey G, Ditunno JF, Jr., Donovan WH, Ducker TB, et al. International standards for neurological and functional classification of spinal cord injury. American Spinal Injury Association. **Spinal Cord** 1997; 35: 266-74.
- Melis EH, Torres-Moreno R, Barbeau H, Lemaire ED. Analysis of assisted-gait characteristics in persons with incomplete spinal cord injury. **Spinal Cord** 1999; 37: 430-9.
- Mong Y, Teo TW, Ng SS. 5-repetition sit-to-stand test in subjects with chronic stroke: reliability and validity. **Arch Phys Med Rehabil** 2010; 91: 407-13.
- National SCI Statistical Center. Annual statistical report for the model SCI care system. Birmingham: University of Alabama at Birmingham, 2006.
- Newcomer KL, Krug HE, Mahowald ML. Validity and reliability of the timed-stands test for patients with rheumatoid arthritis and other chronic diseases. **J Rheumatol** 1993; 20: 21-7.
- New PW, Rawicki HB, Bailey MJ. Nontraumatic spinal cord injury: demographic characteristics and complications. **Arch Phys Med Rehabil** 2002; 83: 996-1001.
- Palombaro KM, Craik RL, Mangione KK, Tomlinson JD. Determining meaningful changes in gait speed after hip fracture. **Phys Ther** 2006; 86: 809-16.
- Penrod LE, Hegde SK, Ditunno JF, Jr. Age effect on prognosis for functional recovery in acute, traumatic central cord syndrome. **Arch Phys Med Rehabil** 1990; 71: 963-8.
- Perera S, Mody SH, Woodman RC, Studenski SA. Meaningful change and responsiveness in common physical performance measures in older adults. **J Am Geriatr Soc** 2006; 54: 743-9.
- Perry J, Garrett M, Gronley JK, Mulroy SJ. Classification of walking handicap in the stroke population. **Stroke** 1995; 26: 982-9.
- Persad CC, Cook S, Giordani B. Assessing falls in the elderly: should we use simple screening tests or a comprehensive fall risk evaluation? **Eur J Phys Rehabil Med** 2010; 46: 249-59.

- Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. **J Am Geriatr Soc** 1991; 39: 142-8.
- Read MS, Sisto SA, Ditunno JF. Standardized ambulation assessments following spinal cord injury. **Top Spinal Cord Inj Rehabil** 2008; 14: 39–60
- Richards CL, Malouin F, Wood-Dauphinee S, Williams JI, Bouchard JP, Brunet D. Task-specific physical therapy for optimization of gait recovery in acute stroke patients. **Arch Phys Med Rehabil** 1993; 74: 612-20.
- Rossier P , Wade DT. Validity and reliability comparison of 4 mobility measures in patients presenting with neurologic impairment. **Arch Phys Med Rehabil** 2001; 82: 9-13.
- Runge M, Rehfeld G, Resnicek E. Balance training and exercise in geriatric patients. **J Musculoskelet Neuronal Interact** 2000; 1: 61-5.
- Salbach NM, Mayo NE, Higgins J, Ahmed S, Finch LE, Richards CL. Responsiveness and predictability of gait speed and other disability measures in acute stroke. **Arch Phys Med Rehabil** 2001; 82: 1204-12.
- Saraf P, Rafferty MR, Moore JL, Kahn JH, Hendron K, Leech K, et al. Daily stepping in individuals with motor incomplete spinal cord injury. **Phys Ther** 2010; 90: 224-35.
- Schenkman M, Hughes MA, Samsa G, Studenski S. The relative importance of strength and balance in chair rise by functionally impaired older individuals. **J Am Geriatr Soc** 1996; 44: 1441-6.
- Scivoletto G, Romanelli A, Mariotti A, Marinucci D, Tamburella F, Mammone A, et al. Clinical factors that affect walking level and performance in chronic spinal cord lesion patients. **Spine** 2008; 33: 259-64.
- Shumway-Cook A, Brauer S, Woollacott M. Predicting the probability for falls in community-dwelling older adults using the Timed Up & Go Test. **Phys Ther** 2000; 80: 896-903.
- Steffen TM, Hacker TA, Mollinger L. Age- and gender-related test performance in community-dwelling elderly people: Six-Minute Walk Test, Berg Balance Scale, Timed Up & Go Test, and gait speeds. **Phys Ther** 2002; 82: 128-37.
- Subbarao JV. Walking after spinal cord injury. Goal or wish? **West J Med** 1991; 154: 612-4.

- Tang SF, Tuel SM, McKay WB, Dimitrijevic MR. Correlation of motor control in the supine position and assistive device used for ambulation in chronic incomplete spinal cord-injured persons. **Am J Phys Med Rehabil** 1994; 73: 268-74.
- Ouellette MM, LeBrasseur NK, Bean JF, Phillips E, Stein J, Frontera WR, et al. High-intensity resistance training improves muscle strength, self-reported function, and disability in long-term stroke survivors. **Stroke** 2004; 35: 1404-9.
- van Asbeck FW, Post MW, Pangalila RF. An epidemiological description of spinal cord injuries in The Netherlands in 1994. **Spinal Cord** 2000; 38: 420-4.
- van Hedel HJ. Gait speed in relation to categories of functional ambulation after spinal cord injury. **Neurorehabil Neural Repair** 2009; 23: 343-50.
- _____, Dietz V, Curt A. Assessment of walking speed and distance in subjects with an incomplete spinal cord injury. **Neurorehabil Neural Repair** 2007; 21: 295-301.
- _____, Wirz M, Curt A. Improving walking assessment in subjects with an incomplete spinal cord injury: responsiveness. **Spinal Cord** 2006; 44: 352-6.
- _____, Wirz M, Dietz V. Assessing walking ability in subjects with spinal cord injury: validity and reliability of 3 walking tests. **Arch Phys Med Rehabil** 2005; 86: 190-6.
- _____, Wirz M, Dietz V. Standardized assessment of walking capacity after spinal cord injury: the European network approach. **Neurol Res** 2008; 30: 61-73.
- van Iersel MB, Munneke M, Esselink RA, Benraad CE, Olde Rikkert MG. Gait velocity and the Timed-Up-and-Go test were sensitive to changes in mobility in frail elderly patients. **J Clin Epidemiol** 2008; 61: 186-91.
- van Wieringen P. Ecological and dynamical approaches to rehabilitation: an epilogue. **Hum Mov Sci** 1996; 15: 315-23.
- Waters RL, Adkins RH, Yakura JS. Definition of complete spinal cord injury. **Paraplegia** 1991; 29: 573-81.
- _____, Adkins R, Yakura J, Sie I. Motor and sensory recovery following incomplete tetraplegia. **Arch Phys Med Rehabil** 1994;75:306-11.
- _____, Yakura JS, Adkins R, Barnes G. Determinants of gait performance following spinal cord injury. **Arch Phys Med Rehabil** 1989; 70: 811-8.

- Weiss A, Suzuki T, Bean J, Fielding RA. High intensity strength training improves strength and functional performance after stroke. **Am J Phys Med Rehabil** 2000; 79: 369-76; quiz 91-4.
- Wernig A , Muller S. Laufband locomotion with body weight support improved walking in persons with severe spinal cord injuries. **Paraplegia** 1992; 30: 229-38.
- Whitney SL, Wrisley DM, Marchetti GF, Gee MA, Redfern MS, Furman JM. Clinical measurement of sit-to-stand performance in people with balance disorders: validity of data for the Five-Times-Sit-to-Stand Test. **Phys Ther** 2005; 85: 1034-45.
- Wirz M, van Hedel HJ, Rupp R, Curt A, Dietz V. Muscle force and gait performance: relationships after spinal cord injury. **Arch Phys Med Rehabil** 2006; 87: 1218-22.
- Wyndaele M , Wyndaele JJ. Incidence, prevalence and epidemiology of spinal cord injury: what learns a worldwide literature survey? **Spinal Cord** 2006; 44: 523-9.
- Zörner B, Dietz V, Curt A. Clinical algorithm for predicting walking capacity based on the ASIA motor score in acute SCI. Paper presented at: Combined ASIA/ISCOS Meeting; 2006; Boston, MA.