

Topic: Investigation of Microbial Dynamics and Metabolic Profiles in Biohydrogen Production from Food Waste

Name of student: Ms. Suwimon Kanchanasuta **Student ID:** 52920109

Name of Supervisor: Asst. Prof. Dr. Nipon Pisutpaisal

ABSTRACT

Dark H₂ fermentation of a synthetic non-sterile food waste was carried out in 0.5 L batch reactors to study the effects of organic loading by the activity of the organisms in the reactors and to compare the activity of *Clostridium butyricum* TISTR1032 with anaerobic sludge under the conditions of initial pH 6 and 37°C. Better performance of H₂ fermentation regarding the H₂ yield and specific rate by *C. butyricum* over the anaerobic sludge was observed at 2.5% VS food waste. Up-scaling for 5 L semi-batch reactors by *C. butyricum* with 2.5% VS food waste was studied under controlled and uncontrolled pH condition to monitor the dominant organisms and metabolic profiles in long term fermentation. An accomplishment of H₂ production in 5 L semi-batch reactors based on the H₂ yield, specific H₂ production and maximum H₂ production under uncontrolled pH condition performed higher than those under controlled pH condition. Metabolite profiles and DGGE results indicated that solventogenesis was induced and it is the limitation factor under uncontrolled pH condition. On the other hand, acetogenesis and lactic acid production was presumed to result in the competition of *C. butyricum* and finally affected the stability of the overall hydrogen fermentation process under controlled pH condition. To better understand the metabolic flux reaction in dark fermentation by anaerobic sludge from complex substrates, rice starch fermentation was constructed in 0.5 L batch reactor and metabolic flux analysis was used to evaluate intracellular flux. The results indicated that acetogenesis and butyric acid oxidation pathways were the important actions resulting in the net H₂ production.

Keywords: hydrogen fermentation, food waste, *C.butyricum*, microbial dynamics, Metabolic Flux Analysis (MFA)

ACKNOWLEDGEMENTS

I would like to express my gratitude to the Joint Graduation School of Energy and Environment (JGSEE), King Mongkut's University of Technology Thonburi, National Research Council of Thailand (NRCT), and Electricity Generation Authority of Thailand (Grant No. 52-2115-031-JOB NO. 518-KMUTNB) for financial support. I also acknowledge Dr. Dong-Hoon Kim (KIER, Korea) for the MetaFluxNet software version 1.8 for the metabolic flux analysis. I greatly appreciate Dr. Somkiet Techkarnjanaruk and Ms. Nimaradee Boonapatcharone for their valuable assistance in the microbial population analysis. I would also like to express my sincere thanks to my thesis advisor, Dr. Nipon Pisutpaisal, for his invaluable help and constant encouragement throughout the course of this research. Finally, I most gratefully acknowledge my family for their support and encouragement.

CONTENTS

CHAPTER	TITLE	PAGE
	ABSTRACT	i
	ACKNOWLEDGEMENT	ii
	CONTENTS	iii
	LIST OF TABLES	v
	LIST OF FIGURES	vi
1	INTRODUCTION	1
	1.1. Rational/Problem Statement	1
	1.2. Literature Review	3
	1.3. Research Objectives	18
2	THEORIES	20
	2.1 Biohydrogen production method	20
	2.2 Denaturing Gradient Gel Electrophoresis	25
3	METHODOLOGY	28
	3.1 Material	28
	3.2 Method	29
	3.3 Analytical techniques	30
	3.4 Kinetic analysis	34
4	Comparative Performance between Heat-Shocked Anaerobic Sludge and <i>Clostridium butyricum</i> TISTR 1032 Inocula in Biohydrogen Production from Food Waste	35
	4.1 Hydrogen fermentation by <i>C.butyricum</i> in 0.5 L reactor	35
	4.2 Hydrogen fermentation by mixed culture in 0.5 L reactor	43
	4.3 Comparative performance of anaerobic sludge and <i>C.butyricum</i> in biohydrogen fermentation from food waste	44
	4.4 Conclusion	45
5	Investigation of Metabolic Profiles and Microbial Community on Biohydrogen Production by <i>Clostridium butyricum</i> TISTR1032 from Food Waste	51

CONTENTS (Cont')

CHAPTER	TITLE	PAGE
	5.1 Hydrogen production and cumulative hydrogen production in 5 L semi-batch reactor	51
	5.2 Metabolites accumulated during fermentative process	56
	5.3 Substrate degradation and utilization during fermentative Process	61
	5.4 Carbon distribution under uncontrolled pH condition	64
	5.5 Microbial dynamics analysis	66
	5.6 Kinetic parameter of hydrogen production	72
	5.7 Conclusion	72
6	Metabolic Flux Analysis of Hydrogen Production from Rice Starch by Anaerobic Sludge under Varying Organic Loading	75
	6.1 Carbon balance and recovery for starch fermentation	75
	6.2 Metabolic flux analysis	76
	6.3 Maximizing H ₂ production via acetic acid production	77
	6.4 Conclusion	79
7	CONCLUSION	87
	REFERENCES	89
	APPENDIXES	104

LIST OF TABLES

TABLES	TITLE	PAGE
1.1	The pure bacterial cultures for fermentative hydrogen production.	11
1.2	Summary of MFA applying for H ₂ dark fermentation.	18
2.1	Advantages and disadvantages of different hydrogen production process.	27
4.1	Yield of biohydrogen production by pure culture from various substrates.	42
4.2	Summarized yield and rate of biohydrogen production by <i>C. Butyricum</i> .	42
4.3	Yield of bio-hydrogen production by mixed culture from various substrates.	49
4.4	Summarized yield of biohydrogen production by mixed culture.	50
5.1	Fermentative periods and cumulative hydrogen production under uncontrolled pH condition.	53
5.2	Function of co-existed organism in fermentative reactor involved with hydrogen production.	68
5.3	Summarized kinetic parameters in 5 L semi batch reactor.	73
5.4	Comparison of kinetic parameters in biohydrogen production.	74
6.1	Carbon balance for starch fermentation after 20 hrs incubation by mixed culture.	80

LIST OF FIGURES

FIGURE	TITLE	PAGE
1.1	Hydrogen role in the anaerobic degradation of organic matter by microbial consortia.	8
1.2	Metabolic pathway in <i>Clostridium spp.</i> fermenting glucose.	15
1.3	Biochemical pathway utilized by <i>Clostridium spp.</i> for the conversion of carbohydrates to hydrogen, carbon dioxide, organic acids and solvents.	16
2.1	An example of DGGE gel showing band compositions of various population samples.	26
4.1	Hydrogen content, hydrogen volume and cumulative hydrogen produced by <i>C. butyricum</i> from varying food waste concentrations.	37
4.2	VFA accumulation in the reactor content after 96 hr fermentation.	38
4.3	COD removal after 96 hr fermentation.	39
4.4	Cumulative hydrogen (H_{max}) and maximum hydrogen production rate (R_{max}).	40
4.5	Hydrogen production yield based on mole of hexose, and gram of food waste addition.	41
4.6	Hydrogen content, hydrogen volume and cumulative hydrogen produced by anaerobic sludge in various F/M ratio.	46
4.7	VFA concentration after 70 hr fermentation.	47
4.8	Maximum hydrogen cumulative value in mL, mmol and maximum hydrogen production rate in unit mL/ hr, mmol/ hr.	48
5.1	Hydrogen production and cumulative hydrogen by <i>C. butyricum</i> under uncontrolled pH condition.	54
5.2	Hydrogen production and cumulative hydrogen by <i>C. butyricum</i> under controlled pH condition.	55
5.3	Variation of pH, metabolites and total concentration of undissociated acid by <i>C. butyricum</i> during hydrogen fermentative under uncontrolled pH condition.	59

LIST OF FIGURES (Cont')

FIGURE	TITLE	PAGE
5.4	Variation of pH, metabolites and total concentration of undissociated acid by <i>C.butyricum</i> during the hydrogen fermentative under uncontrolled pH condition.	60
5.5	Variation of TS and VS during the hydrogen fermentation under uncontrolled pH condition and controlled pH condition.	62
5.6	Variation of soluble carbohydrate during the hydrogen fermentation under uncontrolled pH condition and controlled pH condition.	63
5.7	Carbon distribution during the hydrogen fermentation under uncontrolled pH condition.	65
5.8	Carbon distribution in liquid phase during the hydrogen fermentation under uncontrolled pH condition.	65
5.9	Amplification of partial 16S rRNA genes sequence with GC-clamp for DGGE analysis.	69
5.10	Dynamics of microbial community structure in long term under uncontrolled pH condition.	70
5.11	Structure of microbial dynamics in first feed compared between that in controlled pH condition (Re1) and that in uncontrolled pH condition (Re2).	71
6.1	Cumulative hydrogen production by anaerobic sludge at different starch concentrations.	81
6.2	Hydrolysis, Utilization and H ₂ production, Carbon balance analysis at 20 hr starch fermentation at different concentrations.	82
6.3	Molar flux distributions of dark fermentation using starch as substrate at different concentrations.	83
6.4	Metabolite fluxes on starch fermentation at different concentrations.	84

LIST OF FIGURES (Cont')

FIGURE	TITLE	PAGE
6.5	(A) Acetic acid production measured from experiment and calculated from flux distribution and Acetic acid production by ACCOA intermediate pathway (R15), acetogenesis (R 17) and (B) butyric acid oxidation {(R26)*2} on starch fermentation at different concentrations.	85
6.6	H ₂ production (R15) from ferredoxin pathway, butyric acid degradation (R26*2) and hydrogen consumption (R 17*4) fluxes and H ₂ productivity on starch fermentation at different concentrations.	86