## **CONTENTS**

|                  |   |  | PAGE |  |
|------------------|---|--|------|--|
| EN               | IGLI:   | SH ABSTRACT  | ii   |  |
| TH               | iv  |  |      |  |
| ACKNOWLEDGEMENTS |   |  |      |  |
| CONTENTS         |   |  |      |  |
| LIST OF TABLES   |   |  |      |  |
| LIST OF FIGURES  |   |  |      |  |
| LIST OF SYMBOLS  |   |  |      |  |
| LI               | LIST OF ABBREVIATIONS                                     |  |      |  |
| CI               | HAP   | ΓER  |      |  |
| 1.               | INT   | RODUCTION  | 1    |  |
| 2.               | LIT   | 3  |      |  |
|                  | 2.1   | Introduction   | 3    |  |
|                  | 2.2   | Preparation of CeO <sub>2</sub> nanoparticles                                | 3    |  |
|                  | 2.3   | Preparation of Ni/CeO <sub>2</sub> and Ni/CeO <sub>2</sub> -ZrO <sub>2</sub> | 4    |  |
|                  | 2.4   | Methane steam reforming over CeO <sub>2</sub> , Ni/CeO <sub>2</sub> and      | 5    |  |
|                  |   | Ni/CeO <sub>2</sub> -ZrO <sub>2</sub> based catalyst                         |      |  |
| 3.               | THEORY  |  | 7    |  |
|                  | 3.1   | Introduction   | 7    |  |
|                  | 3.2   | Cerium oxide   | 7    |  |
|                  | 3.3   | Colloidal emulsion aphrons (CEAs)  | 10   |  |
|                  | 3.4   | Influence of various factors on the nanoparticle size                        | 14   |  |
|                  | 3.5   | Gas sorption measurement of adsorbents                                       | 17   |  |
|                  | 3.6   | Methane steam reforming  | 21   |  |
| 4.               | . PREPARATION OF NANO-SIZED CeO <sub>2</sub> BY DIFFERENT |  |      |  |
|                  | MICROEMULSION METHODS                                     |  |      |  |
|                  | 4.1   | Introduction   | 25   |  |
|                  | 4.2   | Objectives   | 26   |  |

|                  | 4.3   | Working scopes   | 26  |  |
|------------------|---|--|-----|--|
|                  | 4.4   | Experimental   | 26  |  |
|                  | 4.5   | Results and discussion                                     | 35  |  |
|                  | 4.6   | Summary  | 50  |  |
| 5.               | PRE   | EPARATION OF Ni LOADING ON NANO-SIZED                      | 51  |  |
|                  | CeO <sub>2</sub> AND CeO <sub>2</sub> -ZrO <sub>2</sub> BY COLLOIDAL EMULSION |  |     |  |
|                  | APHRONS METHOD  |  |     |  |
|                  | 5.1   | Introduction   | 51  |  |
|                  | 5.2   | Objectives   | 52  |  |
|                  | 5.3   | Working scopes   | 52  |  |
|                  | 5.4   | Experimental   | 52  |  |
|                  | 5.5   | Results and discussion                                     | 55  |  |
|                  | 5.6   | Summary  | 72  |  |
| 6.               | COI   | NCLUSIONS AND RECOMMENDATIONS                              | 73  |  |
|                  | 6.1   | Conclusions  | 73  |  |
|                  | 6.2   | Recommendations  | 75  |  |
| RI               | EFER  | RENCES   | 76  |  |
| Αŀ               | PPEN  | DICES  | 84  |  |
|                  | A   | The size distribution evaluated from transmission electron | 84  |  |
|                  |   | microscope (TEM) micrograph                                |     |  |
|                  | В   | Data of nitrogen adsorption-desorption isotherm and pore   | 94  |  |
|                  |   | size distribution  |     |  |
| CURRICULUM VITAE |   |  | 103 |  |