

## REFERENCES

- Adejumo, T.O., Hettwer, U. and Karlovsky, P. (2007). Survey of maize from southwestern Nigeria for zearalenone, α- and β-zearalenols, fumonisin B1 and enniatins produced by Fusarium species. *Food Additive*, 24, 993-1000.
- Agaliya, P.J. and Jeevaratnam, K. (2012). Screening of *Lactobacillus plantarum* isolated from fermented idli batter for probiotic properties. *African Journal of biotechnology*, 11, 12856-12864.
- Amiri, Z.R., Khandelwal, P. and Aruna, B.R. (2010). Development of acidophilus milk via selected probiotics & prebiotics using artificial neural network. *Advances in Bioscience and Biotechnology*,1, 224-231. Retived from http://www.SciRP.org/journal/abb/.
- Anukul, N., Vangnai, K. and Mahakarnchanakul, M. (2013). Significance of regulation limits in mycotoxin contamination in Asia and risk management programs at the national level. *Journal of Food and Drug analysis*, 21, 227-241.
- Avantaggiato, G., Havenaar, R. and Visconti, A. (2003). Assessing the zearalenone-binding activity of adsorbent materials during passage through a dynamic in vitro gastrointestinal model. *Food and Chemical Toxicology*, 41, 1283-1290.
- Barnard, J.A. and Warwick, G. (1993). Butyrate rapidly induces growth inhibition and differentiation in HT-29 cells. *Cell Growth Differ*, 4, 495-501.
- Barros, G.G., Oviedo, M.S., Ramirez, M.L. and Chulze, S.N. (2011). Safety aspect in soybean food and feed chains: fungal and mycotoxins contamination. In Tzi-Bun Ng. (Ed). Soybean- Biochemistry, Chemistry and Physiology. (Vol., pp 7-20). InTech, Retived from: http://www.intechopen.com/books/soybean-biochemistry-chemistry-and-physiology/safety-aspects-in-soybeanfood-and-feed-chains-fungal-and-mycotoxins-contamination. Published online 26, April, 2011.
- Begley, M., Hill, C. and Gahan, C. G. M. (2006). Bile salt hydrolase activity in probiotics. *Applied and Environmental Microbiology*, 72, 1729-1738.
- Bennett, J.W. and Klich, M. (2003). Mycotoxins. *Clinical Microbiology Reviews*, 16, 497-516.

- Berthiller, F., Schuhmacher, R., Adam, G. and Krska, R. (2009). Formation, determination and significance of masked and other conjugated mycotoxins. *Anal Bioanal Chemistry*, 39(5), 1243-1252.
- Bogdanov, S., Juredic, T. and Sieber, R. (2008). Honey for nutrition and health: a Review. *American Journal of the College of Nutrition*, 27, 677-689.
- Bordignon, J.R., Kazuhiko, N., Tadashi, Y. and Sayuki, N. (2004). Hydrolysis of isoflavone and consumption oligosaccharides during lactic acid fermentation of soybean. *International Research Center for Agricuture Sciences (JIRCAS)*, Tsukuba, Ibaraki 305-8686, Japan (http://www. Jircas.affrc.go.jp)
- Božanić, R., Brletić, S. and Lovković, S. (2008). Influence of temperature and sugar addition on soymilk fermentation by probiotic bacteria. *Mljekarstvo*, 58, 61-68,
- Broggi, L.E., González, H.H.L., Resnik, S.L. and Pacin, A. (2007). Alternaria alternata prevalence in cereal grains and soybean seeds from Entre Ríos, *Argentina*. *Revista Iberoamericana Micologia*, 24, 47-51.
- Bueno, D.J., L. di Marco, Oliver. G. and Bardón. A. (2006). In vitro binding of zearalenone to different adsorbents. *Journal Food Protection*, 68,613-615.
- Carrão-Panizzi, M.C., Beleia, A. P., Prudencio-Ferreira, S.H., Oliveira, M.C.N., Kitamura, K. (1999). Effect of isoflavones on beany flavor and astringency of soymilk and cooked whole soybean grain. Pesq. agropec. bras., *Brasilia*, 34,1045-1052.
- Chick, H., Shin, H.S. and Ustunol, Z. (2001). Growth and acid production by lactic acid bacteria and bifidobacteria grown in skim milk containing honey. *Journal of Food Science*, 66, 478-481.
- Chow J. (2002). Probiotics and prebiotics: A brief overview, *Journal of Renal Nutrition*, 12, 76–86.
- Chumchuere, S. and Robinson, R.K., (1999). Selection of starter cultures for the fermentation of soya milk. *Food Microbiology*, 16, 129–137.
- Collado, M.C. (2009). Role of probiotics in health and diseases. In Lee, Y.K. and Salminen, S. (Ed). *Hand book of probiotics and prebiotics*. (2<sup>nd</sup> ed, pp. 3-176). New jersey: John Wiley & Sons, Inc publishing.

- Crittendall, R. (2006). Emerging prebiotic carbohydrates. In Gibson, G.R. and Rastall, R.A. (Ed). *Prebiotics: development & application*. (pp.111-134). West Sussex, UK: Willey-Blackwell, John Wiley & Sons, Inc publishing.
- Crittenden, R.G., Morris, L.F., Harvey, M.L., Tran, M.L., Mitchell, H.L. and Playne, M.J. (2001). Selection of a *Bifidobacterium* strain to complement resistant starch in a symbiotic yoghurt. *Journal of Applied Microbiol*, 90, 268–278.
- Cummings, J.H. (1997). Short-chain fatty acid enemas in the treatment of distal ulcerative colitis. *European Journal of Gastroenterology & Hepatology*, 9, 149-153.
- Dajanta, K., Chukeatirote, E., Apichartsrangkoon, A. and Frazier, RA. (2009). Enhanced aglycone production of fermented soybean products by *Bacillus* species. *Acta Biologica Szegediensis* (2): pp.93-98. Retrieved from http://www.sci.u-szeged.hu/ABS 53
- Dalié, D.K.D., Deschamps, A.M. and Richard-Forget, F. (2010). Lactic acid bacteria Potential for control of mould growth and mycotoxins: A review. *Food control*, 21, 370-380.
- Dall'Asta, C., Galaverna, G., Dossena, A. and Marchelli, R. (2004). Reversed-phase liquid chromatographic method for the determination of Ochratoxin A in wine. *Journal of Chromatography A*, 1024, 275–279.
- Delcour, J., Ferain T., Deghorain M., Palumbo E. and Hols P. (1999). The biosynthesis and functionality of the cell-wall of lactic acid bacteria. *Antonie van Leeuwenhoek*. 76, 159–184.
- Dixit, A.K., Antony, J. I. X., Sharma, N.K. and Tiwari, R.K. (2011). Soybean constituents and their functional benefits. Opportunity. *Challenge and Scope of Natural Products in Medicinal Chemistry*, 367-383.
- Du-Toit, M., Franz, C., Schillinger, U., Warles, B. and Holzappfel, W. (1998).

  Characterization and selection of probiotic lactobacilli for a preliminary minipig-feeding trail and their effect on serum cholesterol level and faeces moisture contents. *International Journal of Food Microbiol*, 40, 93-104.

- El-Nezami, HS., Polychronaki, N., Salminen, S. and Mykkanen, H. (2002). Binding rather metabolism may explain the interaction of two food-grade *Lactobacillus* strains with zearalenone and its derivative a-zearalenol. *Applied and Environmental Microbiology*, 68, 3545–3549.
- EFSA (2004). Opinion of the Scientific Panel on Contaminants in the Food Chain on a request from the Commission related to Zearalenone as undesirable substance in animal feed, *The EFSA Journal*, 89, 1-35
- Ezekiel, C.N., Odebode, A.C. and Fapohunda, S.O. (2008) Zearalenone production by naturally occurring *Fusarium* species on maize, wheat and soybeans from Nigeria. *Journal of biology and environment. Science*, 2(6), 77-82.
- Fandohan, P., Hell, K., Marasas, W.F.O. and Wingfield, M.J. (2003). Infection of maize by Fusarium species and contamination with fumonisin in Africa. *African Journal of Biotechnology*, 2, 570-579.
- Farnworth, E.R., Mainville, I., Desjardins, M.P., Gardner, N., Fliss, I. Champagne, C. (2007). Growth of probiotic bacteria and bifidobacteria in a soy yogurt formulation. *International Journal of Food Microbiology*, 116,174–181.
- Faucet-Marquis, V., Pont, F., Stormer, F.C., Rizk, T., Castegnaro, M. and Pfohl-Leszkowicz, A. (2006). Evidence of a new dechlorinated ochratoxin A derivative formed in opossum kidney cell cultures after pretreatment by modulators of glutathione pathways: Correlation with DNA-adduct formation.

  \*Molecular Nutrition & Food Research\*, 50, 530-542
- Floch, M.H. (2010). Intestinal micro ecology. In Floch, M.H. and Kim, A.S. (Ed). Probiotics: A clinical guide. (3-12). New Jersey, US: SLACK Inc.
- Floch, M.H.and Hong-Curtiss, J. (2002). Probiotics and functional foods in gastrointestinal disorders. *Current Treatment Options in Gastroenterol*. 5, 311-321.
- Fooks, L.J., Fuller, R. and Gibson, G.R. (1999). Prebiotic, probiotic and human gut microbiology. *International Dairy Journal*, 9, 53-61.
- Franz, C.M.A.P., Cho, G., Holzapfel, W.H. and Gálvez, A. (2010). Safety of lactic acid bacteria. In: Mozzi, F., Raya, R., Vignolo, G. (Eds.), *Biotechnology of Lactic Acid Bacteria Novel Applications*. Wiley-Blackwell, Iowa, USA.

- Fuchs, S., Sontag, G., Stidl, R., Ehrlich, V., Kundi, M. and Knasmuller, S. (2008).
  Detoxification of patulin and ochratoxin A, two abundant mycotoxins, by lactic acid bacteria. *Food and Chemical Toxicology*. 46, 1398–1407.
- Fuller, R. (1989). Probiotics in man and animals. *Journal of Applied Bacteriology*. 66, 375-378.
- Garro, M.S., De Valdez, G.F., Oliver, G. and De Giori, G.S., (1999). Starter culture activity in refrigerated fermented soymilk. *Journal of Food Protection*, 62, 808–810.
- Garro, M.S., De Valdez, G.F. and De Giori, G.S. (2004). Temperature effect on the biological activity of *Bifidobacterium longum* CRL 849 and *Lactobacillus fermentum* CRL 251 in pure and mixed cultures grown in soymilk. *Food Microbiology*, 21, 511–518.
- Geraldo, M.R.F., Tessmann, D.J. and Kemmelmeier, C. (2006). Production of mycotoxins by *Fusarium graminaerum* isolated from small cereals (wheat, triticale and barley) affected with scab disease in Southern. *Brazilian Journal of Microbiology*, 37, 34-39.
- Gibson, G.R. and Fuller, R. (2000). Aspects of in vitro and in vivo research approaches directed toward identifying probiotics and probiotics for human use. *Journal of Nutrition*, 130, 391S-395S.
- Gilliland, SE. (1979). Beneficial interrelationships between certain microorganisms and human: Candidate microorganisms for use as dietary adjuncts. *Journal of Food protection*, 42(2): 164-167.
- Gilliand, S.E. and Walker, D.K. (1990). Factors to consider when selecting a culture of *L. acidophilus* as a dietary adjunct to produce a hypercholesterolemic effect in human. *Journal of Dairy Science*, 73, 905-911.
- Golinski, P., Waskiewicz, A. and Gromadzka, k. (2010a). Zearalenone and its derivatives: known toxins in new aspects. In Rai, M. and Verma, A. (Ed). *Mycotoxins in food, feed and bioweapons*. (pp.113-130). London: Springer Heidelberg Dordrecht publishing.

- Hadjeba-Medjdoub, K., Faucet-Marquis, V., Joannis-Cassan, C., Ballet, N. and Pfohl-Leszkowicz, A. (2009). Analysis of the factor influencing mycotoxin adsorption on yeast/development of a method allowing screening, p. 12. In Proceedings of the 2009 International Society for Mycotoxicology conference, Worldwide Mycotoxin Reduction in Food and Feed Chains. International Society for Mycotoxicology, University of Natural Resources and Applied Life Sciences, Vienna.
- Henningsson, I. and Nyman, M. (2001). Short-chain fatty acid formation at fermentation of indigestible carbohydrates. *Scandinavian Journal of Nutrition*, 145,165-168.
- Henningsson, A.M.; Björck, I.M. and Nyman, E.M. (2002). Combinations of indigestible carbohydrates affect short-chain fatty acid formation in the hindgut of rats. *Journal of Nutrition*, 132, 3098-3104.
- Hirayama, K. and Rafter, J. (2006). Probiotic in cancer prevent. In Goktepe, I., Juneja,
  V.K. and Ahmedna, M. (Ed). *Probiotics in food safety and human health*.
  (pp.365-382). London: CRC Press, an imprint of Taylor&Francis Group publishing.
- Holzapfel, W.H. and Schillinger, U. (2002). Introduction to pre- and probiotics. *Food Research International*, 35, 109-116.
- Huda-Faujan, N., Abdulamir, A.S., Fatimah, A.B., Muhammad, A.O., Shuhaimi, M., Yazid, A.M.and Loong, Y.Y. (2010). The impact of the level of the intestinal short chain fatty acids in inflammatory bowel disease patients versus healthy subjects. *The Open Biochemistry Journal*, 4, 53-58.
- Huebner, J., Wehling, R.L.and Hutkins, R.W., (2007). Functional activity of commercial prebiotics. *International of Dairy Journal*, 17, 770–775.
- Hyeong-Jun L., So-Young K. and Wan-Kyu L. (2004). Isolation of cholesterol-lowering lactic acid bacteria from human intestine for probiotic use. *Journal of veterinary science*, 5,391-395.
- International Agency for Research on Cancer (IARC). (1993). Some naturally occurring substances: Food items and constituents, heterocyclic aromatic amines, and mycotoxins. *Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans*, Vol. 56. Lyon (France): IARC.

- Itsaranuwat, P. (2003). *Aspects of the fermentation of soy milk*. Ph.D. Thesis in Food Biotechnology, University of Reading, England.
- Itsaranuwat, P. and Robinson, R.K. (2003). Production and use of dairy products in Thailand. *International Journal of Dairy Technology*, 56, 6-11.
- Janardhana, G.R., Rabeas, K.A. and Shekar Shetty, H., 1999. Mycotoxin contamination of maize grains grown in Karnataka (India). *Food and Chemical Toxicology*, 37, 863–868.
- Joannis-Cassan, C., Tozlovanu, M., Hadjeba-Medjdoub, K., Ballet, N. and Pfohl-Leszkowicz, A. (2011). Binding of Zearalenone, Aflatoxin B1, and Ochratoxin A by yeast-based products: A method for quantification of adsorption performance. *Journal of Food Protection*, 74, 1175-1185.
- Jooyandeh, H. (2011). Soy products as healthy and functional foods. *Middle-East Journal of Scientific Research*, 7, 71-80.
- Kailasapathy, K. and Chin, J. (2000). Survival and therapeutic potential of probiotic organisms with reference to *Lactobacillus acidophilus* and *Bifidobacterium* spp. *Immunology & Cell Biology*, 78, 80–88.
- Kailasapathy, K. (2010). Probiotic and prebiotic fermented foods. In Tamang, J.P. and Kailasapathy, K.(Ed). *Fermented foods and beverages of the world*. (Vol. 1, pp. 377-390). London: CRC Press, an imprint of Taylor&Francis Group publishing.
- Kamaly, K. M. (1997). Bifidobacteria fermentation of soybean milk. *Food Research International*, 30, 675-682.
- Kansandee, W. (2010). Screening and identification of probiotic Lactic Acid Bacteria from human colostrums. Master's thesis. Mahasarakham University.

  Mahasarakham, Thailand. 120 p.
- Kaur, I.P., Chopra, K. and Saini, A. (2002) Probiotics: potential pharmaceutical applications. *European Journal of Pharmaceutical Sciences*, 15, 1-9.
- Kiely, L.J. and Olson, N.F. (2000). The physicochemical surface characteristics of Lactobacillus casei. *Food Microbiology*, 17, 277-291.

- Kim, Y., Whang, JY., Whang, KY., Oh, S. and Kim, SH. (2008). Characterization of the cholesterol-reducing activity in a cell-free supernatant of *Lactobacillus* acidophilus ATCC 43121. Bioscience. Biotechnology. and Biochemistry, 72, 1483-90.
- Kimoto, H., Ohmomo, S. and Okamoto, T. (2002). Cholesterol Removal from Media by *Lactococci. Journal of Dairy Science*, 85, 3182-3188.
- Klaver, FA. and Van der Meer, R. (1993). The assumed assimilation of cholesterol by *Lactobacilli* and *Bifidobacterium bifidum* is due to their bile salt-deconjugating activity. *Applied and Environmental Microbiology*, 59, 1120-1124.
- Klayraung, S., Viernstein, H., Sirithunyalug, J. and Okonogi, S. (2008). Probiotic Properties of *Lactobacilli* Isolated from Thai Traditional Food. *Scientia Pharmaceutica*, 76, 485–503.
- Kurman, J.A. and Rasic, J.L. (1991). The health potential of products containing bifidobacteria. In: Therapeutic properties of functional milks. 117-158.
- Lannitti, T. and Palmieri, B. (2010). Therapeutical use of probiotic formulations in clinical practice. *Clinical Nutrition*. 29, 701-725.
- Lawless, H.T. and Heymann, H. (1999). *Sensory evaluation of food: Principles and practices*. Aspen Publishers, Inc., Gaitherburg, Maryland.
- Lilly, D.M.and Stillwell, R.H. (1965): Probiotics: Growth promoting factors produced by microorganisms. *Science*, 147,747-748.
- Liong M.T. (2006). *In-vivo and in-vitro cholesterol removal by Lactobacilli and Bifidobacteria*. PhD thesis. School of Molecular Sciences Victoria University, Werribee Campus, VIC, Australia.
- Liong, MT. and Shah, NP. (2005a). Acid and bile tolerance and cholesterol removal ability of lactobacilli strains. *Journal of Dairy Science*. 88, 55-66.
- Liong, M.T. and Shah, N.P. (2005b). Optimization of cholesterol removal, growth and fermentation patterns of Lactobacillus acidophilus ATCC 4962 in the presence of mannitol, fructooligosaccharide and inulin: A response surface methodology approach. *Journal of Applied Microbiology*, 98, 1115-1126.
- Lin, M.Y., and Chen, T.W. (2000). Reduction of cholesterol by *lactobacillus acidophilus* in culture broth. *Journal of Food Drug Analysis*, 8, 97-102.

- Liu, K. (1997). *Soybeans: chemistry, technology and utilization*. (535 p). New York: International Thomson publishing, an imprint of Chapman and Hall.
- Liu, S.N., Han, Y., and Zhou, Z.J. (2011). Lactic acid bacteria in traditional fermented Chinese foods: A review. *Food Research International*, 44: pp 643-651.
- Ly, C.M.H., Le, M.L., Thanh, M.L., Belin, J.M. and Wach, Y., (2010). Cell surface properties affect colonisation of raw milk by lactic acid bacteria at the microstructure level. *Food Research International*, 43,1594–1602.
- Lye, H.S., Rusul, G., and Liong, M.T. (2010). Removal of cholesterol by lactobacilli via incorporation of and conversion to coprostanol. *Journal of Dairy Science*, 2010, 93, 1383-1392.
- Lu, Q., Liang, X. and Chen, F. (2011). Detoxification of Zearalenone by viable and inactivated cells of Planococcus sp. *Food control*. 22, 191-195.
- Madani, G., Mirlohi, M., Yahay, M. and Hassanzadeh, A. (2013). Relationship between in vivo and in vitro cholesterol removal of *Lactobacillus spp*. *International Journal of Preventive Medicine*, 4 (4), 404-413.
- Mahrous, H. (2011). Probiotics bacteria from Egyptian infants cause cholesterol removal in media and survive in yoghurt. *Food and nutrition*. 2,150-155.
- Marona, D. and Pedrigon, G. (2004). Yoghurt feeding inhibits promotion and progression of cancer. *Medical Science Monitor*, 10, 96-104.
- Martı'nez-Villaluenga, C. and Go'mez, R. (2007). Characterization of bifidobacteria as starters in fermented milk containing raffinose family of oligosaccharides from lupin as prebiotic. *International Dairy Journal*, 17,116–122.
- Mathara J.M., Schillinger U., Guigas C., Franz C., Kutima P.M., Mbugua S.K., Shin H.K. and Holzapfel W.H. (2008): Functional characteristics of *Lactobacillus* spp. from traditional Maasai fermented milk products in Kenya. *International Journal of Food Microbiology*, 126, 57–64.
- Megharaj, M., Garthwaite, I. and Thiele, J.H. (1997). Total biodegradation of the oestrogenic mycotoxin zearalenone by a bacterial culture. *Letters in Applied Microbiology*, 24, 329–333.
- Metchnikoff, E. (1908). *The prolongation of life*. Optimistic studies. G.P. Putnam's sons, New York.

- Mirlohi, M., Madani, G., Hassanzadeh, A. and Yahay, M. (2012). On the colorimetric method for cholesterol determination in the laboratory media. *International of Journal Biology and Chemistry*, 6 (1), 37-42.
- Mital, B.K., Steinkraus, K.H. and Naylor, H.B. (1974). Growth of lactic acid bacteria in soy milks. *Journal of Food Science*, 39, 1018–1022.
- Mokoena, M.P., Chelule, P.K. and Gqalen, N. (2005). Reduction of fumonisin B1 and Zearalenone by lactic acid bacteria in fermented maize meal. *Journal of Food Protection*, 68, 2095–2099.
- Nakano, R., Hara M., Ishiguro, H., Yao, Y., Ochiai, T., Nakata, K., Murakami, T., Kajioka, J., Sunada, K., Hashimoto, K., Fujishima, A. and Kubota, Y. (2013). Broad spectrum microbicidal activity of photocatalysis by TiO<sub>2</sub>. *Catalysts*, 3(1), 310-323; Available online; http://www.mdpi.com/2073-344/3/1/310/htm# sthash.B4Z2UynD.dpuf
- Nuryono, N., Noviandi, C.T., Bo.hm, J. and Razzazi-Fazeli, E., (2005). A limited survey of zearalenone in Indonesian maize-based food and feed by ELISA and high performance liquid chromatography. *Food Control*, 16, 65–71.
- Oliveira, P.M., Zannini, E., and Arendt, E.K. (2014). Cereal fungal infection, mycotoxins, and lactic acid bacteria mediated bioprotection: From crop farming to cereal products. *Food Microbiology*, 37, 78-95.
- Ooi, L-G. and Liong, M.T. (2010). Cholesterol-lowering effects of probiotics and prebiotics: A review of in vivo and in vitro findings. *International Journal of Molecular Sciences*, 11, 2499-2522.
- Ouwehand, A.C., Kirjavainen, P.V., Shortt, C. and Salminen, S. (1999). Probiotics: mechanisms and established effects. *International Dairy Journal*, 9, 43-52.
- Ouwehand, A.C., Salminen, S., and Isolauri, E. (2002). Probiotics: an overview of beneficial effects. *Antonie van Leeuwenhoek*, 82, 279–289.
- Ott, L. (1988). An introduction to statistical methods and data analysis. Boston: PWS-Kent.
- Parker, R.B. (1974). Probiotics, the other half of the antibotics story. *Animal Nutrition* and *Health*. 29, 4-8.

- Park, J.W., Kim, E.K., Shon, D.H. and Kim, Y.B., (2002). Natural cooccurrence of aflatoxin B1, fumonisin B1 and ochratoxin A in barley and corn foods from Korea. *Food Additives & Contaminats*, 19, 1073–1080.
- Park, J.W., Shoi, S.Y., Hwang, H.J. and Kim, Y.B. (2005). Fungal mycoflora and mycotoxins in Korean polished rice destined for humans. *International Journal of Food Microbiololy*, 103, 305–314.
- Parvez, S., Malik, K.A., Kang, S.A. and Kim, H.Y. (2006). Probiotics and their fermented food products are beneficial for health. *Journal of Applied Microbiology*, 100, 1171–1185.
- Păucean, A., Mudura, E., Jimborean, M.A.M. and Man, S. (2011). The influence of honey addition on microbiological, physicochemical and sensory characteristics of a kefir-type product during shelf-life. *Bulletin UASVM Agriculture*, 68,393-399.
- Perdigeon, G., Valdez, J. and Rachid, M. (1998). Antitumour activity of yoghurt. Study of possible immune response. *Journal of Dairy Research*, 65, 129-138.
- Pereira, D.I.A. and Gibson, G.R. (2002). Cholesterol assimilation by lactic acid bacteria and bifidobacteria isolated from the human gut. *Applied and Environmental Microbiology*, 68, 4689-4693.
- Pereira, D.I.A., McCartney, A.L. and Gibsob, GR. (2003). An *In vitro* study of the probiotic potential of a bile-salt-hydrolyzing *Lactobacillus fermentum* strain, and determination of its cholesterol lowing properties. *Applied and Environmental Microbiology*. 69, 4743-4752.
- Pfohl-Leszkowicz, A., Chekir-Ghedir, L. and Bacha, L. (1995). Genotoxicity of zearalenone, an oestrogenic mycotoxin: DNA adduct formation in female mouse tissues. *Carcinogenesis*, 16, 2315–2320.
- Phillips, S.I., Wareing, P.W., Dutta, A., Panigrahi, S. and Medlock, V. (1996). The mycoflora and incidence of aflatoxin, zearalenone and sterigmatocystin in dairy feed and forage samples from eastern India and Bangladesh.

  Mycopathologia, 133, 15–21.
- Phommalth, S., Jeong, Y.S., Kim, Y.H. and Hwang, Y.H. (2008). Isoflavone composition within each structural part of soybean seeds and sprouts. *Journal Crop Science Biotechnology*, 11, 57-62.

- Placinta, C.M., D'Mello, J.P.F. and Macdonald, A.M.C. (1999). A review of worldwide contamination of cereal grains and animal feed with *Fusarium* mycotoxins. *Animal Feed Science and Technol*, 78, 21–37.
- Ram, C. and Chander, H. (2003). Optimization of culture conditions of probiotic bifidobacteria for maximal adhesion to hexadecane. *World Journal of Microbiology and Biotechnology*, 19, 407–410.
- Ramasamy, K., Abdullah, N., Wong, M.C.V.L., Karuth, C. and Hob, Y. W. (2009). Bile salt deconjugation and cholesterol removal from media by *Lactobacillus* strains used as probiotics in chickens. *Journal Science of Food Agriculture*, 90, 65–69.
- Ramos, A.J., Hernandez, E., Pla-Delfina, J.M. and Merino, M. (1996). Intestinal absorption of zearalenone and in vitro study of nonnutritive sorbent materials. *International Journal of Pharmaceutics*, 128, 129–137.
- Reddy, K.R.N., Salleh, B., Saad, B., Abbas, H.K., Abel, C.A. and Sheir, W.T. (2010). An overview of mycotoxin contamination in foods and its implications for human health. *Toxin Reviews*, 29, 3-26.
- Riazi, A. and Ziar, H. (2012). Effect of honey and starter culture on growth, acidification, sensory properties and bifidobacteria cell counts in fermented skimmed milk. *African Journal of Microbiology Research*, 6, 486-498.
- Richardson, K.E., Hagler, W.M. Jr., Haney, C.A., and Hamilton, R.B. (1985).

  Zearalenone and trichothecene production in soybeans by toxigenic Fusarium. *Journal of Food Protection*, 48, 240–243.
- Roberfroid, M. (2007). Prebiotics: The concept revisited. *Journal of Nutrition*, 137, 830S-837S.
- Rojas, M., Ascencio, F. and Conway, P.L. (2002). Purification and characterization of a surface protein from *Lactobacillus fermentum* 104R that binds to porcine small intestinal mucus and gastric mucin. *Applied and Environmental Microbiology*, 68, 2330-2336.
- Rosenberg, M., Gutnick, D. and Rosenberg, E. (1980). Adherence of bacteria to hydrocarbons: a simple method for measuring cell surface hydrophobicity. *FEMS Microbiology Letters*, 9, 29–33.

- Ross, S, and Jonsson, H (2002). A high-molecular mass cell surface protein from *Lactobacillus reuteri* 1063 adheres to mucus components. *Microbiology*,148, 433-442.
- Roumyan, N., Zapryanov, P. and Kondareva, S. (1996). On some aspects of a new fermented milk product medina. *Biotechnology and Biotechnological Equipment*, 10, 86-89.
- Rudel, L.L. and Morris, M.D. (1973). Determination of cholesterol using *o*-phthalaldehyde. *Journal of Lipid Research*, 14, 364-6.
- Salminen, S. (1996). Uniqueness of probiotic strains. *IDF Nutrition News Letter*, 5, 16-18.
- Sánzhes-Zapata, E., Fernández-López, J., Pérez-Alvarez, J.A., Soares, J., Sousa, S., Gomes, A.P.M. and Pintado, M.M.E. (2013). In vitro evaluation of "horchata" co-products as carbon source for probiotic bacteria growth. *Food and Bioproducts Processing*, 91, 279–286.
- Savage, DC. (1992). Growth phase, cellular hydrophobicity and adhesion in vitro of lactobacilli colonizing the keratinizing gastric epithelium in the mouse. *Apply Environment Microbiology*, 58, 1992–1995.
- Schaafsma, G. (1996). State of art concerning probiotics strains in milk products. *IDF*Nutrition News Letter, 5, 23-24.
- Scheppach, W. (1994). Effect of shorth chain fatty acids on gut morphology and function. Gut 1994; supplement1: s35-s38.
- Scheppach, W.; Bartram, H.P. and Richter, F. (1995). Role of short chain fatty acids
  I. prevention of colorectal cancer. *European Journal of Cancer*, 31, 1077-1080.
  Schillinger, U., Guigas, C. and Holzapfel, W.H. (2005). *In vitro* adherence and other properties of lactobacilli used in probiotic yoghurt-like products. *International Dairy Journal*, 15, 1289-1297.
- Schmid, K., Schlothauer, R.C., Friedrich, U., Staudt, C., Apajalahti, J. and Hansen, E.B. (2006). Development of probiotic food ingredients. In Goktepe, I., Juneja, V.K. and Ahmedna, M. (Ed). *Probiotics in food safety and human health*. (pp.35-66). London: CRC Press, an imprint of Taylor&Francis Group publishing.

- Seeling, K., Lebzien, P., Danicke, S., Spilke, J., Sudekum, K.H, and Flachowsky, G. (2006). Effects of level of feed intake and *Fusarium* toxin contaminated wheat on rumen fermentation as well as on blood and milk parameters in cows. *Journal of animal Physiology and Animal Nutrition*, 90, 103-110.
- Shah, N.P. (2007). Review: Functional cultures and health benefits. *Journal of International Dairy*, 17, 1262-1277.
- Shetty, P.H. and Jespersen, L. (2006). *Saccharomyces cerevisiae* and lactic acid bacteria as potential mycotoxin decontamination agents. *Trends in Food Science & Technology*, 17, 48-55.
- Sherbert, G.V. (1978). *The biophysical characterization of the cell surface*. London: Academic press.
- Sieladie, DV., Zambou, NF., Kaktcham, PM., Cresci, A. and Fonteh, F. (2011).

  Probiotic properties of lactobacilli strains isolated from raw cow milk in the western highlands of Cameroon. *Innovative Romanian Food Biotechnology*, 9, 12-28.
- Sirilun, S. (2012). Probiotic Lactobacillus plantarum LCC 150 isolated from Thai fermented food and cholesterol lowering property of its fermented soybean milk product. Ph.D. in pharmacy Thesis. Chiang Mai University.
- Sirilun, S., Chaiyasut, C., Kantachote, D. and Luxananil, P. (2010). Characterisation of non human origin probiotic *Lactobacillus plantarum* with cholesterol-lowering property. *African Journal of Microbiology Research*, 4, 994-1000.
- Smith, T.K., McMillian, E.G. and Castillo, J.B. (1997). Effects of feeding blends of Fusarium mycotoxin-contaminated grains containing deoxynivalenol and fusaric acid on growth and feed consumption of immature swine. *Journal of Animal Science*, 75, 2184-2191.
- Soccol, C.R., Vandenberghe, L.P.S., Spier, Medeiros, A.B.P., Yamaguishi, C.T., Lindner, J. D.D., Pandey, A. and Soccol, V.T. (2010): The potential of probiotics: A review. *Food Technology and Biotechnology*, 48, 413-434.
- Sodini, I., Lucas, A., Oliveira, MN., Remeuf, F. and Corrieu, G. (2002). Effect of milk base and starter culture on acidification, texture, and probiotic cell counts in fermented milk processing. *Journal of Dairy Science*, 85, 2479-2488.

- St-Onge, M.P., Farnworth, E.R. and Jones, P. (2000). Consumption of fermented and nonfermented dairy products: Effects on cholesterol concentrations and metabolism. *The American Journal of Clinical Nutrition*, 71, 674-681.
- Su, P., Henriksson, A. and Mitchell, H., (2007). Selected prebiotics support the growth of probiotic mono-cultures in vitro. *Anaerobe*,13, 134–139.
- Sumarna (2008). Changes of raffinose and stachyose in soy milk fermentation by lactic acid bacteria from local fermented foods of Indonesian. *Malaysian Journal of Microbiology*, 4(2), 26-34
- Szabo, N.J. Dolan, L.C., Burdock, G.A., Shibano, T. Sato, S.I., Suzuki, H., Uesugi, T., Yamahira, S., Toba, M. and Ueno, H. (2011). Safety evaluation of *Lactobacillus pentosus* strain b240. *Food and Chemical Toxicology*, 49, 251-258.
- Tamminen M, Joutsjoki T, Sjöblom M, Joutsen M, Palva A., Ryhänen E-L and Joutsjoki V. (2004); Screening of lactic acid bacteria from fermented vegetables by carbohydrate profiling and PCR-ELISA, *Letters in Applied Microbiology*. 39, 439-444.
- Tanasupawat, S. and K. Komagata. (1994). Lactic acid bacteria in fermented foods in Thailand. *World Journal of Microbiology and Biotechnology*, 11: 253–256.
- Tanasupawat, S., Okada, S., and Komagata, K. (1998). Lactic acid bacteria found in fermented fish in Thailand, *J Gen Appl Microbiol*.44 (3); 193-200.
- Tanasupawat, S., Pakdeeto, A., Thawai, C., Yukphan, P. and Okada, S. (2007).

  Identification of lactic acid bacteria from fermented tea leaves (miang) in

  Thailand and proposals of *Lactobacillus thailandensis* sp. nov., *Lactobacillus camelliae* sp. nov., and *Pediococcus siamensis* sp. nov. *Journal of Genetic Apply Microbiology*. 53(1): 7-15.
- Usman, H.A. (1999). Bile Tolerance, Taurocholate Deconjugation, and Binding of Cholesterol by *Lactobacillus gasseri* Strains. *Journal of Dairy Science*, 82, 243-248.
- Varga, L. (2006). Effect of acacia (*Robinia pseudo-acacia* L.) honey on the characteristic microflora of yoghurt during refrigerated storage. *International Journal of Food Microbiology*, 108, 272-275.

- Vasconcelos de Sá L. R., Oliveira, M. A. L., Cammarota M. C., Matos A. and Ferreira-Leita, V. S., (2011). Simultaneous analysis of carbohydrates and volatile fatty acids by HPLC for monitoring fermentative biohydrogen production. *International Journal of hydrogen energy*, 1, 646-650.
- Vasiljevic, T. and Shah, N.P. (2008). Probiotics from Metchnikoff to bioactives. *International Dairy Journal*, 18, 714–728.
- Venema, K. (2012). Intestinal fermentation of lactose and prebiotic lactose derivatives, including human milk oligosaccharides. *International Dairy Journal*, 22, 123-140.
- Villaluenga, M.C. and Gómez, R. (2007). Characterization of bifidobacteria as starters in fermented milk containing raffinose family of oligosaccharides from lupin as prebiotic. *International Dairy Journal*, 17, 116–122
- Vinderola, C.G. and Rrinheimer, J.A. (2003). Lactic acid starter and probiotic bacteria: a comparative "in vitro" study of probiotic characteristics and biological barrier resistance. *Food Research International*, 36, 895-904.
- Vijayaraghavan, K. and Yun, Y.S. (2008). Bacterial biosorbents and biosorption. *Biotechnology Advances*, 26, 266-291.
- Wang, Y.B. and Han, J.Z. (2007). The role of probiotic cell wall hydrophobicity in bioremediation of aquaculture. *Aquaculture*, 269, 349-354.
- Wang, Y.C., Yu, R.C. and Chou, C.C. (2002). Growth and survival of bifidobacteria and lactic acid bacteria during the fermentation and storage of cultured soymilk drinks. *Food Microbiology*, 19, 501–508.
- Wang, YC., Yu, RC., Yang, HY. and Chou, CC. (2003). Sugar and acid contents in soymilk fermented with lactic acid bacteria. *Food Microbiology*, 20, 333-338.
- Wright, A.V. and Axelsson, L. (2012). Lactic acid bacteria: An introduction. In Lahtinen, S., Salminen, S., Ouwehang, A., and Wright A.V. (Ed). *Lactic acid bacteria: microbiological and functional aspects*. (4<sup>th</sup> ed, pp.705-748). London: CRC Press, an imprint of Taylor&Francis Group publishing.
- Yamashita, A., Yoshizawa, T., Aiura, Y., Sanchez, P.C., Dizon, E.I., Arim, R.H., Sardjono. (1995). Fusarium mycotoxins (fumonisins, nivalenol and zearalenone) and aflatoxins in corn from southeast Asia. *Bioscience, Biotechnology, Biochemistry*, 59, 1804–1807.

- Yoawapa, S. (2013). Detoxification mycotoxin by microorganisms. *Research group of Agro-biotechnology*. Retrieved http://www.gpo.or.th/rdi/html/mycotoxin.html. January 20, 2013.
- Yoshizawa, T. (1997). Geographic difference in trichothecene occurrence in Japanese wheat and barley. *Bull. Inst. Compr. Agric. Sci. Kinki University*, 5, 23–30.
- Yoshizawa, T. and Jin, Y.Z. (1995). Natural occurrence of acetylated derivatives of deoxynivalenol and nivalenol in wheat and barley in Japan. *Food Additive & Contaminants*, 12, 689–694.
- Zinedine, A., Soriano, J.M., Moltó, J.C., and Mańes, J. (2007). Review on the toxicity, occurrence, metabolism, detoxification, regulations and intake of Zearalenone:

  An oestrogenic mycotoxin. *Food and Chemical Toxicology*, 45, 1–18.