

REFERENCES

- Aiking, H., & Tempest, D.W. (1976). Growth and physiology of *Candida utilis* HCYC 321 in potassium-limited chemostat culture. **Arch Microbiol**, **108**, 117-127.
- Albers, E., Larsson, C., Lidén, G., Niklasson, C., & Gustafsson, L. (1996). Influence of the nitrogen source on *Saccharomyces cerevisiae* anaerobic growth and product formation. **Appl Environ Microbiol**, **62**, 3187-3195.
- Alexander, H., & Charpentier, C. (1998). Biochemical aspects of stuck and sluggish fermentation in *grape must*. **J Ind Microbiol Biotech**, **20**, 20-27.
- Alfenore, S., Cameleyre, X., Benbadis, L., Bideaux, C., Uribe Larrea, J.L., & Goma, G., et al. (2004). Aeration strategy: a need for very high ethanol performance in *Saccharomyces cerevisiae* fed-batch process. **Appl Microbiol Biot**, **63**, 537-542.
- Bai, F.W., Anderson, W.A., & Moo-Young, M. (2008). Ethanol fermentation technologies from sugar and starch feedstocks. **Biotechnol Adv**, **26**, 89-105.
- _____, Chen, L.J., Zhang, Z., Anderson, W.A., & Moo-Young, M. (2004). Continuous ethanol production and evaluation of yeast cell lysis and viability loss under very high gravity medium conditions. **J Biotechnol**, **110**, 287-293.
- Balat, M. (2011). Production of bioethanol from lignocellulosic materials via the biochemical pathway: A review. **Energy Convers Manage**, **52**, 858-875.
- Banat, I.M., Nigam, P., Singh, D., Marchant, R., & McHale, A.P. (1998). Review: Ethanol production at elevated temperatures and alcohol concentrations: Part I- Yeasts in general. **World J Microb Biot**, **14**, 809-821.
- Benjaphokee, S., Hasegawa, D., Yokota, D., Asvarak, T., Auesukaree, C., & Sugiyama, M., et al. (2012). Highly efficient bioethanol production by a *Saccharomyces cerevisiae* strain with multiple stress tolerance to high temperature, acid and ethanol. **New Biotechnol**, **29**(3), 379-386.

- Beltran, G., Esteve-Zarzoso, B. Rozes, N., Mas, A., & Guillamoän, J.M. (2005). Influence of the timing of nitrogen additions during synthetic grape must fermentations on fermentation kinetics and nitrogen consumption. **J Agric Food Chem**, **53**, 996-1002.
- Berry, D.R., Stewart, G.G., & Russell, I. (1987). **Yeast Biotechnology**. London: Unwin Hyman.
- Bhima, B., Anjana, D.T., Sudhakara, R.M., Ramana, R.Y., & Venkateswar, R.L. (2010). Insights into the protein profiles of stress tolerant and mesophilic yeast by proteomics approach: a comparative study. **IJABPT**, **1**(2), 349-359.
- Birch, R.M., & Walker, G.M. (2000). Influence of magnesium ions on heat shock and ethanol stress responses of *Saccharomyces cerevisiae*. **Enzyme Microb Tech**, **26**, 678-687.
- Bradford, M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dry binding. **Annual Biochem**, **72**, 248-254.
- Burrows, S. (1970). Yeast Technology. In A.H. Rose & J.S. Harrison (Eds.). **The Yeast**. (pp. 349-420). New York: Academic Press.
- Casey, G.P. & Ingledew, W.M. (1986). Ethanol tolerance in yeasts. **Crit Rev Microbiol**, **13**, 219-280.
- Castro, R.G., Baigorí, D.M., & Siñeriz, F. (1995). A plate technique for screening of inulin degrading microorganisms. **J Microbiol Methods**, **22**, 51-56.
- Chi, Z.M., Chi, Z., Zhang, T., Liu, G.L., & Yue, L.X. (2009). Inulinase expressing microorganisms and applications of inulinases. **Appl Microbiol Biotechnol**, **82**, 211-220.
- Chubey, B.B., & Dorell, D.G. (1974). Jerusalem artichoke - a potential fructose crop for the prairies, **Can Institute Food Sci Technol J**, **7**, 98-106.
- Cooper, T.G. (1982). Nitrogen metabolism in *Saccharomyces cerevisiae*. In J.N. Strathren, E.W. Jone, & J.R. Broach (Eds.). **The molecular biology of the yeast *Saccharomyces***. (pp. 39-99). New York: Cold Spring Harbor Laboratories.

- Cooper, T.G., Britton, C., Brand, L., & Sumrada, R. (1979). Addition of basic amino acids prevents G-1 arrest of nitrogen starved cultures of *Saccharomyces cerevisiae*. **J Bacteriol**, **137**, 1447-1448.
- D'Amore, T. (1992). Improving yeast fermentation performance. **J Inst Brew**, **98**, 375-382.
- de Souza, C.J.A., Costa, D.A., Rodrigues, M.Q.R.B., dos Santos, A.F., Lopes, M.R., Abrantes, A.B.P., et al. (2012). The influence of presaccharification, fermentation temperature and yeast strain on ethanol production from sugarcane bagasse. **Bioresour Technol**, **109**, 63-69.
- Dhaliwal, S.S., Oberoi, H.S., Sandhu, S.K., Nanda, D., Kumar, D., & Uppal, S.K. (2011). Enhanced ethanol production from sugarcane juice by galactose adaptation of a newly isolated thermotolerant strain of *Pichia kudriavzevii*. **Bioresour Technol**, **102**, 5968-5975.
- Diaz M, Garcia A.I., & Garcia L.A. (1996). Mixing power, external convection, and effectiveness in bioreactors. **Biotechnol Bioeng**, **51**, 131-40.
- Dorell, D.G, & Chubey, B.B. (1977). Irrigation, fertilizer, harvest dates and storage effects on the reducing sugar and fructose concentration of Jerusalem artichoke tubers. **Can J Plant Sci**, **57**, 591-596.
- Dubois, M., Gilles, K.A., Hamilton, J.R., Robers, P.A., & Smith, F. (1956). Colourimetric method for determination of sugar and related substances. **Anal Chem**, **28**, 350-356.
- Edgardo, A., Carolina, P., Manuel, R., Juanita, F., & Baeza, J. (2008). Selection of thermotolerant yeast strains *Saccharomyces cerevisiae* for bioethanol production. **Enzyme Microb Tech**, **43**(2), 120-123.
- Fakruddin, M.D., Ariful Islam, M.D., Abdul Quayum, M.D., Morshed, M.A., & Chowdhury, N. (2013). Characterization of stress tolerant high potential ethanol producing yeast from agro-industrial waste. **American Journal of BioScience**, **1**(2), 24-34.
- Ge, X.Y., & Zhang, W.G. (2005). A shortcut to the production of high ethanol concentration from Jerusalem artichoke tubers. **Food Technol Biotechnol**, **43**(3), 241-246.

- Grubb, C.F., & Mawson, A.J. (1993). Effects of elevated solute concentrations on the fermentation of lactose by *Kluyveromyces marxianus* Y-113. **Biotechnol Lett**, **15**, 621-626.
- Hack, C.J., & Marchant, R. (1998). Characterisation of a novel thermotolerant yeast, *Kluyveromyces marxianus* var *marxianus*: development of an ethanol fermentation process. **J Ind Microbiol Biot**, **20**, 323-327.
- Harju, S., Fedosyuk, H., & Peterson, K.R. (2004). Rapid isolation of yeast genomic DNA: Bust n' Grab. **BMC Biotechnol**, **4**, 8.
- Hashem, M., Zohri, A.N.A., & Ali, M.M.A. (2013). Optimization of the fermentation conditions for ethanol production by new thermotolerant yeast strains of *Kluyveromyces* sp. **Afr J Microbiol Res**, **7**(37), 4550-4561.
- Hu, N., Yuan, B., Sun, J., Wang, S., & Li, F. (2012). Thermotolerant *Kluyveromyces marxianus* and *Saccharomyces cerevisiae* strains representing potentials for bioethanol production from Jerusalem artichoke by consolidated bioprocessing. **Appl Microbiol Biotechnol**, **95**, 1359-1368.
- Ingledew, W.M. (1999). Alcohol production by *Saccharomyces cerevisiae*: a yeast primer. In **The Alcohol Textbook**. 3rd ed. Nottingham: Nottingham University Press.
- Ingram, L.A., & Bukkte, T.M. (1984). Effect of alcohols on microorganisms. **Adv Microb Physiol**, **25**, 253-303.
- Jeffries, T.W., & Jin, Y.S. (2000). Ethanol and thermotolerance in the bioconversion of xylose by yeasts. **Adv Appl Microbiol**, **47**, 221-268.
- Johannesen, R. (1991). Energy Efficiency & Environmental News: Alcohol Production from Biomass. **Florida Energy Extension Service**. Retrieved May 14, 2012, from <http://www.p2pays.org/ref/13/12145.pdf>
- Jones, A.M., & Ingledew, W.M. (1994). Fuel alcohol fermentation: Optimization of temperature for efficiency very-high-gravity fermentation. **Appl Environ Microbiol**, **60**, 1048-1051.
- Jones, M., & Pierce, J.S. (1964). Absorption of amino acid from wort by yeast. **J Inst Brew**, **70**, 307-315.
- Kalmokoff, M.L., & Ingledew, W.M. (1985). Evolution of ethanol tolerance in selected *Sacharomyces cerevisiae* strains. **J Am Soc Brew Chem**, **43**, 189-196.

- Kalil, S.J., Maugeri-Filho, F., & Rodrigues, M.I. (2005). Ion exchange expanded bed chromatography for the purification of an extracellular inulinase from *Kluyveromyces marxianus*. **Proc Biochem**, **40**, 581-586.
- Khampa, S., Wangsomnuk, P.P., Jogloy, S., Wangsomnuk, P. & Pattanothai, A. (2008). Morphological and Anatomical Characteristics of Jerusalem Artichoke (*Helianthus tuberosus* L.) in Thailand. **Khon Kaen Agriculture Journal**, **36**(special), 205-212.
- Kim, S., & Kim, C.H. (2013). Evaluation of whole Jerusalem artichoke (*Helianthus tuberosus* L.) for consolidated bioprocessing ethanol production. **Renew Energ**, **65**, 83-91.
- Kiran, S.N., Sridhar, M., Suresh, K., Banat, I.M., & Venkateswar, R.L. (2000). Isolation of thermotolerant, osmotolerant, flocculating *Saccharomyces cerevisiae* for ethanol production. **Bioresour Technol**, **72**(1), 43-46.
- Koedrith, P., Dubois, E., Scherens, B., Jacobs, E., Boonchird, C., & Messenguy, F. (2008). Identification and characterization of a thermotolerant yeast strain isolated from banana leaves. **ScienceAsia**, **34**, 147-152.
- Kumar, S., Singh, S.P., Mishra, I.M., & Adhikari, D.K. (2009). Ethanol and xylitol production from glucose and xylose at high temperature by *Kluyveromyces* sp. IIPE453. **J Ind Microbiol Biotechnol**, **36**, 1483-1489.
- Kurtzman, C.P., & Robnett, C.J. (1998). Identification and phylogeny of ascomycetous yeasts from analysis of nuclear large subunit (26S) ribosomal {DNA} partial sequences. **Anton Leeuw Int J G**, **73**(4), 331-371.
- Kusch, H., Engelmann, S., Bode, R., Albrecht, D., Morschhäuser, J., & Hecker, M. (2008). A proteomic view of *Candida albicans* yeast cell metabolism in exponential and stationary growth phases. **Int J Med Microbiol**, **298**, 291-318.
- Kushnirov, V.V. (2000). Rapid and reliable protein extraction from yeast. **Yeast**, **16**, 857-860.
- Kwon, Y.J., Ma, A.Z., Li, Q., Wang, F., Zhuang, G.Q., & Liu, C.Z. (2011). Effect of lignocellulosic inhibitory compounds on growth and ethanol fermentation of newly-isolated thermotolerant *Issatchenkia orientalis*. **Bioresour Technol**, **102**, 8099-8104.

- Laopaiboon, L., Nuanpeng, S., Srinophakun, P., Klanrit, P., & Laopaiboon, P. (2009). Ethanol production from sweet sorghum juice using very high gravity technology: Effects of carbon and nitrogen supplementations. **Bioresour Technol**, **100**(18), 4176-4182.
- Lee, C., Yamakawa T., & Kodama, T. (1993). Rapid growth of a thermotolerant yeast on palm oil. **World J Microb Biot**, **9**, 187-190.
- Li, S.Z., & Chan-Halbrendt, C. (2009). Ethanol production in China: Potential and technologies. **Appl Energy**, **86**, S162-S169.
- Li,Y., Liu, G.L., & Chi, Z. (2013). Ethanol production from inulin and unsterilized meal of Jerusalem artichoke tubers by *Saccharomyces* sp. W0 expressing the endo-inulinase gene from *Arthrobacter* sp.. **Bioresour Technol**, **147**, 254-259.
- Lim, S.H., Ryu, J.M., Lee, H., Jeon, J.H., Sok, D.E., & Choi, E.S. (2010). Ethanol fermentation from Jerusalem artichoke powder using *Saccharomyces cerevisiae* KCCM50549 without pretreatment for inulin hydrolysis. **Bioresour Technol**, **102**, 2109-2111.
- Limtong, S., Sringiew, C., & Yongmanitchai, W. (2007). Production of fuel ethanol at high temperature from sugar cane juice by newly isolated *Kluyveromyces marxianus*. **Bioresour Technol**, **98**, 3367-3374.
- Lin, Y.H., Chien, W.S., Duan, K.J., & Chang, P.R. (2011). Effect of aeration timing and interval during very-high-gravity ethanol fermentation. **Process Biochem**, **46**, 1025-1028.
- Madigan, M.T., Martinko, J.M., & Parker, J. (2000). **Nutrition and metabolism, Brock Biology of Microbiology**. 9th ed. New Jersey: Prentice-Hall.
- Margaritis, A., & Bajpai, P. (1983). Effect of sugar concentration in Jerusalem artichoke extract on *Kluyveromyces marxianus* growth and ethanol production. **Appl environ microb**, **45**(2), 723-725.
- Malherbe, S., Bauer, F.F., & Du, T.M. (2007). Understanding problem fermentations - A review. **S Afr J Enol Vitic**, **28**(2), 169-186.
- Mauricio, J.C., & Salmon, J.M. (1992). Apparent loss of sugar transport activity in *Saccharomyces cerevisiae* may mainly account for maximum ethanol production during alcoholic fermentation. **Biotechnol Lett**, **14**, 577-582.

- McCracken, L.D. & Gong, C.S. (1982). Fermentation of cellulose and hemicellulose carbohydrates by thermotolerant yeasts. **Biotechnol Bioeng**, **12**, 91.
- Mehdikhani, P., Bari, M.R., & Hovsepyan, H. (2011). Screening of *Saccharomyces cerevisiae* for high tolerance of ethanol concentration and temperature. **Afr J Microbiol Res**, **5**(18), 2654-2660.
- Mota, M., Besie, J.M., Strehaiano, P., & Goma, G. (1987). A simple device for fed-batch control in alcoholic fermentation. **Biotechnol Bioeng**, **24**, 775-777.
- Nakamura, T., Ogata, Y., Hamada, S., & Ohta, K. (1996). Ethanol production from Jerusalem artichoke tubers by *Aspergillus niger* and *Saccharomyces cerevisiae*. **J Fert Eng**, **81**(6), 564-566.
- Narendranath, N.V., & Power, R. (2005). Relationship between pH and medium dissolved solids in terms of growth and metabolism of Lactobacilli and *Saccharomyces cerevisiae* during ethanol production. **Appl Environ Microbiol**, **71**, 2239-2243.
- Nolan, A.M., Barron, N., Brady, D., McAree, T., Smith, D., & McHale L., et al. (1994). Ethanol production at 45°C by an alginate-immobilized strain of *Kluyveromyces marxianus* following growth on glucose containing media. **Biotechnol Lett**, **16**, 849-852.
- Nuanpeng, S., Laopaiboon, L., Srinophakun, P., Klanrit, P., Jaisil, P., & Laopaiboon, P. (2011). Ethanol production from sweet sorghum juice under very high gravity conditions: Batch, repeated-batch and scale up fermentation. **Electron J Biotechnol**, **14**, 1-12.
- O'Conner-Cox, E.S.C., & Ingledew, W.M. (1989). Wort nitrogenous sources. Their use by brewing yeasts: a review. **J Am Soc Brew Chem**, **47**, 102-108.
- Oda, Y., Nakamura, K., Shinomiya, N., & Ohba, K. (2010). Ethanol fermentation of sugarbeet thick juice diluted with crude cheese whey by the flex yeast *Kluyveromyces marxianus* KD-15. **Biomass Bioenergy**, **34**, 1263-1266.
- O'Donnell, K. (1993). *Fusarium* and its near relatives. In D.R. Reynolds & J.W. Taylor (Eds.). **The fungal holomorph: mitotic, meiotic and pleomorphic speciation in fungal systematics**. (pp. 225-233). Wallingford: CAB International.

- Onsoy, T., Thanonkeo, P., Thanonkeo, S., & Yamada, M. (2007). Ethanol production from Jerusalem artichoke by *Zymomonas mobilis* in batch fermentation. **KMITL Sci Tech J**, 7, 55-60.
- Ozmichi, S., & Kargi, F. (2007). Ethanol fermentation of cheese whey powder solution by repeated fed-batch operation. **Enzyme Microb Technol**, 41, 169-174.
- Pandey, A., Soccol, C.R., Selvakumar, P., Soccol, V.T., Krieger, N., & Jose, D., et al. (1999). Recent developments in microbial inulinases, its production, properties and industrial applications. **Appl Biochem Biotechnol**, 81, 35-52.
- Pang, Z.W., Liang, J.J., Qin, X.J., Wang, J.R., Feng, J.X., & Haung, R.B. (2010). Multiple induced mutagenesis for improvement of ethanol production by *Kluyveromyces marxianus*. **Biotechnol Lett**, 32, 1847-1851.
- Patrascu, E., Rapeanu, G., & Hopulele, T. (2009). Current approaches to efficient biotechnological production of ethanol. **Innovat Rom food Biotechnol**, 4, 1-11.
- Pereira, F.B., Guimarães, P.M.R., Teixeira, J.A., & Domingues, L. (2010). Optimization of low-cost medium for very high gravity ethanol fermentations by *Saccharomyces cerevisiae* using statistical experimental designs. **Bioresour Technol**, 101, 7856-7863.
- Petrov, V.V., & Okorokov, L.A. (1990). Increase of the anion and proton permeability of *Saccharomyces carlsbergensis* plasmalemma by n-alcohols as a possible cause of its de-energization. **Yeast**, 6, 311-318.
- Razmovski, R.N., Šćiban, M.B., Vučurović, V.M. (2011). Bioethanol production from Jerusalem artichoke by acid hydrolysis. **Rom Biotech Lett**, 16(5), 6497-6503.
- Rubio-Arroyo, M.F., Vivanco-Loyo, P., Juárez, M., Martha, P., & Ramírez-Galicia, G. (2011). Bio-ethanol Obtained by Fermentation Process with Continuous Feeding of Yeast. **J Mex Chem Soc**, 55(4), 242-245.
- Sanger, F., Nicklen, S., & Coulson, A.R. (1977). DNA sequencing with chain terminating-inhibitor. **Proc Nalt Acad Sci USA**, 74, 5463-5467.
- Sansonetti, S., Hobley, T.J., Calabró, V., Villadsen, J., & Sin, G. (2011). A biochemically structured model for ethanol fermentation by *Kluyveromyces marxianus*: A batch fermentation and kinetic study. **Bioresour Technol**, 102, 7513-7520.

- Shakhashiri, B.Z. (2009). **Ethanol**. Retrieved May 10, 2012, from <http://scifun.chem.wisc.edu/chemweek/pdf/ethanol.pdf>
- Slapack, G.E., Russell, I., & Stewart, G.G. (1987). **Thermophilic microbes in ethanol production**. Boca Raton: CRC Press.
- Spayd, S.E., & Andersen-Bagge, J. (1996). Free amino acid composition of grape juice from 12 *Vitis vinifera* cultivars in Washington. **Am J Enol Vitic**, **47**, 389-402.
- Sprenger, G.A. (1996). Carbohydrate metabolism in *Zymomonas mobilis*: a catabolic highway with some scenic routes. **FEMS Microbiol Lett**, **145**, 301-307.
- Sumrada, R., & Cooper, T.G. (1978). Basic amino acid inhibition of cell division and macromolecular synthesis of *Saccharomyces cerevisiae*. **J Gen Microbiol**, **108**, 45-56.
- Swanton, C.J, Cavers, P.B, Clements, D.R, & Moore, M.J. (1992). The biology of Canadian weeds: 101 *Helianthus tuberosus* L. **Can J Plant Sci**, **72**, 1367-1382.
- Szambelan, K., & Chrapkowska, K.J. (2003). The influence of selected microorganisms on ethanol yield from Jerusalem artichoke (*Helianthus tuberosus* L.) tubers. **Pol J Food Nutr Sci**, **12**(53), 49-52.
- _____, Nowak, J., & Czarnecki, Z. (2004). Use of *Zymomonas mobilis* and *Saccharomyces cerevisiae* mixed with *Kluyveromyces fragilis* for improved ethanol production from Jerusalem artichoke tubers. **Biotechnol Lett**, **26**, 845-848.
- _____, Nowak, J., & Jelen, H. (2005). The composition of Jerusalem artichoke (*Helianthus tuberosus* L.) spirits obtained from fermentation with bacteria and yeasts. **Eng Life Sci**, **5**, 68-71.
- Thanonkeo, P., Laopaiboon, P., Sootsuwan, K., & Yamada, M. (2007). Magnesium ions improve growth and ethanol production of *Zymomonas mobilis* under heat or ethanol stress. **Biotechnology**, **6**(1), 112-119.
- _____, Thanonkeo, S., Charoensuk, K. & Yamada, M. (2011). Ethanol production from Jerusalem artichoke (*Helianthus tuberosus* L.) by *Zymomonas mobilis* TISTR548. **Afr J Biotechnol**, **10**(52), 10691-10697.
- Theobald, U., Mohns, J., & Rizzi, M. 1996. Dynamics of orthophosphate in yeast cytoplasm. **Biotechnol Lett**, **18**, 461-466.

- Thomas, K.C., & Ingledew, W.M. (1990). Fuel alcohol production: Effects of free amino nitrogen on fermentation of very-high-gravity wheat mash. **Appl Environ Microbiol**, **56**, 2046-2050.
- Thuesombat, P., Thanonkeo, P., Laopaiboon, L., Laopaiboon, P., Yunchalard, S., Kaewkannetra, P., & Thanonkeo, S. (2007). The batch ethanol fermentation of Jerusalem artichoke using *Saccharomyces cerevisiae*. **KMITL Sci Tech J**, **7**, 93-96.
- Todar, K. (2008). **Diversity of metabolism in prokaryotes**. Retrieved May 16, 2012, from http://textbookofbacteriology.net/metabolism_3.html
- Van Uden, N. (1984). Cardinal temperatures of yeast. In A.I. Laskin & H.H.A. Lechevalier (Eds.). **Handbook of Microbiology**. (pp. 1-8). 6th ed. Florida: CRC Press.
- Walker, G.M. (1994). The Roles of Magnesium in Biotechnology. **Crit Rev Biotechnol**, **14**(4), 311-354.
- Walker, G. (1998). Yeast growth. In G. Walker (Ed.). **Yeast: Physiology and Biotechnology**. (pp. 101-202). New York: Wiley.
- Watanabe, T., Srichuwong, S., Arakane, M., Tamiya, S., Yoshinaga, M., & Watanabe, I., et al. (2010). Selection of stress-tolerant yeasts for simultaneous saccharification and fermentation (SSF) of very high gravity (VHG) potato mash to ethanol. **Bioresour Technol**, **101**, 9710-9714.
- Winter, J.F., Loret, M.O., & Uribe Larrea, J.L. (1989). Inhibition and growth factor deficiencies in alcoholic fermentation by *Saccharomyces cerevisiae*. **Curr Microbiol**, **18**, 247-252.
- Yuan, W.J., Zhao, X.M., & Bai, F.W. (2008). Ethanol fermentation with *Kluyveromyces marxianus* from Jerusalem artichoke grown in salina and irrigated with a mixture of seawater and freshwater. **J Appl Microbiol**, **105**, 2076-2083.
- Yuangsaard, N., Yongmanitchai, W., Yamada, M., & Limtong, S. (2012). Selection and characterization of a newly isolated thermotolerant *Pichia kudriavzevii* strain for ethanol production at high temperature from cassava starch hydrolysate. **Antonie van Leeuwenhoek**, **103**(3), 577-588.

- Yue, G., Yu, J., Zhang, X., & Tan, T. (2010). The influence of nitrogen sources on ethanol production by yeast from concentrated sweet sorghum juice. **Biomass Bioenerg**, **39**, 48-52.
- Zafar, S., & Owais, M. (2006). Ethanol production from crude whey by *Kluyveromyces marxianus*. **Biochem Eng J**, **27**, 295-298.
- Zhang, T., Chi, Z., Zhao, C.H., Chi, Z.M., & Gong, F. (2010). Bioethanol production from hydrolysates of inulin and the tuber meal of Jerusalem artichoke by *Saccharomyces* sp. W0. **Bioresour Technol**, **101**, 8166-8170.
- Zhao, C.H., Zhang, T., Li, M., & Chi, C.M. (2010). Single cell oil production from hydrolysates of inulin and extract of tubers of Jerusalem artichoke by *Rhodotorula mucilaginosa* TJY15a. **Process Biochem**, **45**, 1121-1126.
- Zoecklein, B.W., Fugelsang, K.C., Gump, B.H., & Nury, F.S. (1995). **Wine analysis and production**. New York: Chapman & Hall.