

REFERENCES

- Adhirajan, N., Kumar, T.R., Shanmugasundaram, N., Babu, M. (2003). *In vivo* and *in vitro* evaluation of hair growth potential of *Hibiscus rosa-sinensis* Linn. **Journal of Ethnopharmacology**, 88, 235-239.
- Ames, B.N., Lee, F.D., Durston, W.F. (1973). An improved bacterial test system for the detection and classification of mutagens and carcinogens. **Proceedings of the National Academy of Sciences (USA)**. (pp. 782-786). California: Biochemistry Department, University of California.
- Asp, M.L., Collene, A.L., Norris, L.E., Cole, R.M., Cole, R.M., et al. (2011). Time-dependent effects of safflower oil to improve glycemia, inflammation and blood lipids in obese, post-menopausal women with type2 diabetes: A randomized, double-masked, crossover study. **Clinical Nutrition**, 30, 443-449.
- Balogh, P., & Engelmann, P. (Eds.). (2011). **Transdifferentiation and regenerative medicine**. Hungary: University of Pecs.
- Barahmani, N., Schabath, M.B., Duvic, M. (2009). History of atopy or autoimmunity increases risk of alopecia areata. **Journal of American Academy of dermatology**, 61 (4), 581-591.
- Barsh, G., Gunn, T., He, L., Schlossman, S., Duke, C.J. (2003). Biochemical and genetic studies of pigment-type switching. **Pigment Cell Research**, 13 (8), 48-53.
- Bergfeld, W.F. (1995). Androgenetic alopecia: an autosomal dominant disorder. **The American Journal of Medicine**, 98, 95-98.
- Bikle, D.D. (2004). Vitamin D regulated keratinocyte differentiation. **Journal of Cellular Biochemistry**, 92, 436-444.
- Botchkareva, N.V., Khlgatian, M., Longley, B.J., Botchkarev, V.A., Gilchrist, B.A. (2001). Scf/c-kit signaling is required for cyclic regeneration of the hair pigmentation unit. **Federation of American Societies for Experimental Biology**, 15, 645-658.

- Braun, S., Krampert, M., Bodo, E., Kumin, A., Berclaz, C.B., et al. (2006). Keratinocyte growth factor protects epidermis and hair follicles from cell death induced by UV irradiation, chemotherapeutic or cytotoxic agents. **Journal of Cell Science**, 119 (23), 4841-4849.
- Chen, C.C., Chuong, C.M. (2012). Multi-layered environmental regulation on the homeostasis of stem cells: The saga of hair growth and alopecia. **Journal of Dermatological Science**, 66, 3-11.
- Choi, H., Ahn, S., Lee, B.G., Chang, I., Hwang, J.S. (2005). Inhibition of skin pigmentation by an extract of *Lepidium apetalum* and its possible implication in IL-6 mediated signaling. **Pigment Cell Research**, 18, 439-446.
- Cilotti, A., Danza, G., Serio, M. (2001). Clinical applicaiton of 5 α -reductase inhibitor. **Journal of Endocrinological Investigation**, 24, 199-203.
- Datta, K., Singh, A.T., Mukherjee, A., Bhat, B., Ramesh, B., et al. (2009). *Eclipta alba* extract with potential for hair growth promoting activity. **Journal of Ethnopharmacology**, 124, 450-456.
- Ellis, J.A., Stebbing, M., Harrap, S.B. (2001). Polymorphism of the androgen receptor gene is associated with male pattern baldness. **Journal of Investigative Dermatology**, 116 (3), 452-455.
- Eswarakumar, V.P., Laz, I., Schiessinger, J. (2005). Cellular signaling by fibroblast growth factor receptors. **Cytokine & Growth Factor Reviews**, 16, 139-149.
- Eun, H.C., Kwon, O.S., Yeon, J.H., Shin, H.S., Shin, H.S., et al. (2010). Efficacy, safety, and tolerability of dutasteride 0.5 mg once daily in male patients with male pattern hair loss: A randomized double-blind placebo-controlled phase III study. **Journal of American Academy of Dermatology**, 63(2), 252-258.
- Fan, L., Zhao, H.Y., Xu, M., Zhou, L., Guo, H., et al. (2009). Qualitative evaluation and quantitative determination of 10 major active component in *Carthamus tinctorius* L. by high-performanc liquid chromatography couples with diode array detector. **Journal of Chromatography A**, 1216, 2063-2070.

- Faragalla, J., Bremner, J., Brown, D., Griffith, R., Heaton, A. (2003). Comparative pharmacophore development for inhibitors of human and rat 5- α -reductase. **Journal of Molecular Graphics and Modelling**, 22, 83-92.
- Fu, F., Su, C., Liu, K. (2005). Effect of safflower yellow A on the blood pressure in dog and man. **Antihypertensive Drugs and Pharmacology**, 18, 59A.
- Fujie, T., Katoh, S., Oura, H., Urano, Y., Arase, S. (2001). The chemotactic effect of a dermal papilla cell-derived factor on outer root sheath cells. **Journal of Dermatological Science**, 25, 206-212.
- Gahrton, G., Björkstrand, B. (2000). Progress in haematopoietic stem cell transplantation for multiple myeloma. **Journal of Internal Medicine**, 248 (3), 185-201.
- Goldman, S., Windream, M. (2006). Cell replacement therapy in neurological disease. **Philosophical Transactions of The Royal Society B Biological Sciences**, 361 (1473), 1463-1475.
- Harada, N., Okajima, K., Arai, M., Kurihara, H., Nakagata, N. (2007). Administration of capsaicin and isoflavone promotes hair growth by increasing insulin-like growth factor-I production in mice and in humans with alopecia. **Growth Hormone & IGF Research**, 17, 408-415.
- Hashimoto, K. (2000). Regulations of keratinocyte function by growth factors. **Journal of Dermatological Science 24 Suppl**, 1, S46-S50.
- Hoffmann, R., Happle, R. (2000). Current understanding of androgenetic alopecia. Part I. Aetiopathogenesis. **European Journal of Dermatology**, 10, 319-327.
- Hsia, C.W., Shui, H.A., Wang, C.Y., Yu, H.M., Ho, M.Y., et al., (2011). Proteomics demonstration that histone H4 is a colchicine-induced retro-modulator of growth and alkaline phosphatase activity in hair follicle dermal papilla culture. **Journal of Proteomics**, 74, 805-816.
- Huh, J.S., Kang, J.H., Yoo, Y.J., Kim, C.S., Cho, K.S., et al. (2001). The effect of safflower seed fraction extract on human periodontal ligament fibroblast and MC3T3-E1 cell *in vitro*. **Journal of Korean Academy of Periodontology**, 31, 833-846.

- Iehle, C., Radvanyi, F., Medina, S.G.D.D., Ouafik, L.H., Gerard, H., et al. (1999). Differences in steroid 5 α -reductase iso-enzyme expression between normal and pathological human prostate tissue. **The Journal of Steroid Biochemistry and Molecular Biology**, 68 (5-6), 189-195.
- Inamatsu, M., Tochio, T., Makabe, A., Endo, T., Oomizu, S., et al. (2006). Embryonic dermal condensation and adult dermal papilla induce hair follicles in adult glabrous epidermis through different mechanisms. **Development, Growth & Differentiation**, 48, 73-86.
- Inui, S., Fukuzato, Y., Nakajima, T., Yoshikawa, K., Itami, S. (2002). Androgen-inducible TGF- β 1 from balding dermal papilla cells inhibits epithelial cell growth: a clue to understand paradoxical effects of androgen on human hair growth. **FASEB Journal**, 16, 1967-1969.
- Itoh, T., Furuichi, Y. (2005). Hot-water extracts from Adzuki Beans (*Vigna angularis*) stimulate not only melanogenesis in cultured mouse B16 melanoma cells but also pigmentation of hair color in C3H mice. **Bioscience, Biotechnology, and Biochemistry**, 69 (5), 873-882.
- Jadhav, V.M., Thorat, R.M., Kadam, V.J., Gholve, S.B. (2009). Kesharaja: Hair vitalizing herbs. **International Journal of PharmTech Research**, 1, 454-467.
- Jahoda, C.A., Horne, K.A., Oliver, R.F. (1984). Induction of hair growth by implantation of cultured dermal papilla cells. **Nature**, 311(5986), 560-562.
- Jaks, V., Kasper, M., Toftgard, R. (2010). The hair follicle-a stem cell zoo. **Experimental Cell Research**, 316, 1422-1428.
- Jankovic, S.M., Jankovic, S.V. (1998). The control of hair growth. **Dermatology Online Journal**, 4(1), 2.
- Jiang, J.S., Xia, P.F., Feng, Z.M., Zhang, P.C. (2008). Chemical constituents from flowers of *Carthamus tinctorius*. **Zhongguo Zhong Yao Za Zhi**, 33(24), 2911-2913.
- Jun, S.M., Ha, Y.M., Kim, H.S., Jang, H.J., Kim, Y.M., et al. (2011). Anti-inflammatory action of methanol extract of *Carthamus tinctorius* involves in heme oxygenase-1 induction. **Journal of Ethnopharmacology**, 133, 524-530.

- Jung, G.D., Yang, J.Y., Song, E.S., Park, J.W. (2001). Stimulation of melanogenesis by glycyrrhizin in B16 melanoma cells. **Experimental and molecular medicine**, 33 (3), 131-135.
- Kanehira, T., Yakekoshi, S., Nagata, H., Matsuzaki, K., Kambayashi, Y., et al. (2003). A novel and potent biological antioxidant, Kinobeaon A, from cell culture of safflower. **Life Science**, 74, 87-97.
- Karahuseyinoglu, S., Cinar, O., Kilic, E., Kara, F., Gumus, G., et al. (2007). Biology of the stem cells in human umbilical cord stroma: *In situ* and *in vivo* surveys. **Stem Cells**, 25, 319-331.
- Karlsson, T., Vahlquist, A., Torma, H. (2010). Keratinocyte differentiation induced by calcium, phorbol ester or interferon- γ elicits distinct changes in the retinoid signaling pathways. **Journal of Dermatological Science**, 57, 207-213.
- Kaufman, K.D. (2002). Androgens and alopecia. **Molecular and Cellular Endocrinology**, 198, 89-95.
- Kauser, S., Thody, A.J., Schallreuter, K.U., et al. (2005). A fully functional proopiomelanocortin/melanocortin-1 receptor system regulates the differentiation of human scalp hair follicle melanocytes. **Endocrinology**, 146, 532-543.
- Kern, S., Eichler, H., Stoeve, J. (2006). Comparative analysis of mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue. **Stem Cells**, 24, 1294-1301.
- Kim, H.J., Bae, Y.C., Park, R.W., Choi, S.W., Cho, S.H., et al. (2002). Bone-protecting effect of safflower seeds in ovariectomized rats. **Calcified Tissue International**, 71, 88-94.
- _____, Cho, Y.D., Leem, K.H., Lee, D.N., Kim, E.H., et al., (2006). Effects of *Ephedrae herba* on melanogenesis and gene expression profiles using cDNA microarray in B16 melanocytes. **Phytotherapy research**, 20, 748-754.
- Krause, K., Foitzik, K. (2006). Biology of the hair follicle: The basics. **Seminars in Cutaneous Medicine and Surgery**, 25, 2-10.

- Kumar, N., Rungseevijitprapa, W., Narkkhong, N.A., Suttajit, M., Chaivasut, C. (2012). 5 α -reductase inhibition and hair growth promotion of some Thai plants traditionally used for hair treatment. **Journal of Ethnopharmacology**, 139 (3), 765-771.
- Lai-Cheong, J.E., McGrath, J.A., (2009). Structure and function of skin, hair and nails. **Medicine**, 37 (5), 223-226.
- Lee, G.S., Hong, E.J., Gwak, K.S., Park, M.J., Choi, K.C., et al. (2010). The essential oils of *Chamaecyparis obtusa* promote hair growth through the induction of vascular endothelial growth factor gene. **Fitoterapia**, 81 (1), 17-24.
- Lee, Y.C., Oh, S.W., Chang, J., Kim, I.H. (2004). Chemical composition and oxidative stability of safflower oil prepared from safflower seed roasted with different temperatures. **Food Chemistry**, 84, 1-6.
- Lei, T.C., Virador, V.M., Vieira, W.D., Hearing, V.J. (2002). A melanocyte-keratinocyte coculture model to assess regulators of pigmentation *in vitro*. **Analytical Biochemistry**, 305, 260-268.
- Lindvall, O. (2003). Stem cells for cell therapy in Parkinson's disease. **Pharmacological Research**, 47 (4), 279-287.
- Lovkvist, K., Sjolinder, H., Wehelie, R., Aro, H., Norrby-Teglund, A., et al. (2008). CD46 Contributes to the Severity of Group A Streptococcal Infection. **Infection and Immunity**, 76 (9), 3951-3958.
- Matsushima, T., Sawamura, M., Hara, K., Sugima, T. (1976). A safe substitute for polychlorinated biphenyls as an inducer of metabolic activation system. In F.J. de Serres, J.R. Fouts, J.R. Bend and R.M. Philpot (Eds.). **In vitro metabolic activation in mutagenesis testing**. (pp. 85-88). Amsterdam: North Holland.
- McElwee, K.J., Sinclair, R. (2008). Hair physiology and its disorders. **Drug Discovery Today: Disease Mechanisms**, 5, 163-171.
- Mitalipov, S., Wolf, D. (2009). Totipotency, pluripotency & nuclear reprogramming. **Advances in Biochemical Engineering/Biotechnology**, 114, 185-199.
- Moll, I. (1996). Differential epithelial outgrowth of plucked and microdissected human hair follicles in explant culture. **Archives of Dermatological Research**, 288 (10), 604-610.

- Moon, K.D., Back, S.S., Kim, J.H., Jeon, S.M., Lee, M.K., et al. (2001). Safflower seed extract lowers plasma and hepatic lipids in rats fed high-cholesterol diet. **Nutrition Research**, 21, 895-904.
- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival application to proliferation and cytotoxicity assays. **Journal of Immunological Methods**, 65, 55-63.
- Muller, R.S., Handjiski, B., Van, V.C., Eichmuller, S., Foitzik, K., et al. (2001). A comprehensive guide for the accurate classification of murine hair follicles in distinct hair cycle stages. **Journal of Investigative Dermatology**, 117, 3-15.
- Nagata, H., Takekoshi, S., Takeyama, R., Honma, T., Osamura, R. (2004). Quercetin enhances melanogenesis by increasing the activity and synthesis of tyrosinase in human melanoma cells and in normal human melanocytes. **Pigment Cell Research**, 17, 66-73.
- Neste, D.V., Tobin, D.J. (2004). Hair cycle and hair pigmentation: dynamic interactions and changes associated with aging. **Micron**, 35, 193-200.
- Nobakht, M., Fattahi, M., Hoormand, M., Milanian, I., Rahbar, N., et al. (2000). A study on the teratogenic and cytotoxic effects of safflower extract. **Journal of Ethnopharmacology**, 73, 453-459.
- Norwood, O.T. (2001). Incidence of female androgenetic alopecia (female pattern alopecia). **Dermatologic Surgery**, 27, 53-54.
- Occhiato, E.G., Cuarna, A., Danza, G., Serio, M. (2004). Selective non-steroidal inhibitors of 5 α -reductase type 1. **Journal of Steroid Biochemistry & Molecular Biology**, 88, 1-16.
- Oda, Y., Ishikawa, M.H., Hawker, N.P., Yun, Q.C., Bikle, D.D. (2007). Differential role of two VDR coactivators, DRIP205 and SRC-3, in keratinocyte proliferation and differentiation. **Journal of Steroid Biochemistry & Molecular Biology**, 103, 776-780.
- Oh, H.T., Kim, S.H., Choi, H.J., Chung, M.J., Ham, S.S. (2008). Antioxidative and antimutagenic activities of 70% ethanol extract from masou salmon (*Oncorhynchus masou*). **Toxicology**, 22, 1484-1488.

- Olsen, E.A., Hordinsky, M., Whiting, D., Stough, D., Hobbs, S., et al. (2006). The importance of dual 5 α -reductase inhibition in the treatment of male pattern hair loss: Results of a randomized placebo-controlled study of dutasteride versus finasteride. **Journal of American Academy of dermatology**, 55(6), 1014-1023.
- Ornelas, V.P., Cabeza, M., Bratoeff, E., Heuze, I., Sanchez, M., et al., (2005). New 5 α -reductase inhibitors: *In vitro* and *in vivo* effects. **Steroids**, 70, 217-224.
- Ozeki, M., Tabata, Y. (2003). *In vivo* promoted growth of mice hair follicles by the controlled release of growth factors. **Biology materials**, 24, 2387-2394.
- Park, W.S., Lee, C.H., Lee, B.G., Chang, I.S. (2003). The extract of *Thuja occidentalis* semen inhibited 5 α -reductase and androchronogenetic alopecia of B6CBAF1/j hybrid mouse. **Journal of Dermatological Science**, 31 (2), 91-98.
- Price, V.H. (2003). Androgenetic alopecia in women. **Journal of Investigative Dermatology Symposium**, 8, 24-27.
- Qiao, J., Turetsky, A., Kemp, P., Teumer, J. (2008). Hair morphogenesis *in vitro*: formation of hair structures suitable for implantation. **Future Medicine**, 3 (5), 683-692.
- Quinn, A.G. (2004). Biology of the skin and dermatological disease. **Medicine**, 32 (12), 1-3.
- Reid, E.E., Haley, A.C., Borovicka, J.H., Rademaker, A., West, D.P., et al. (2012). Clinical severity does not reliably predict quality of life in women with alopecia areata, telogen effluvium, or androgenic alopecia. **Journal of American Academy of dermatology**, 66 (3), 91-102.
- Ren, C., Fang, L., Ling, L., Wang, Q., Liu, S., et al. (2009). Design and *in vivo* evaluation of an indapamide transdermal patch. **International Journal of Pharmaceutics**, 370, 129-135.
- ReportLinker. (2012). Hair care products industry: Market research reports, statistics and analysis. **Industry reports, company data and country profiles**, Retrieved April 20, 2012, from www.reportlinker.com

- Reynolds, A.J., Jahoda, C.A. (1996). Hair matrix germinative epidermal cells confer follicle-inducing capabilities on dermal sheath and high passage papilla cells. **Development**, 122 (10), 3085-3094.
- Rho, S.S., Kim, C.D., Lee, M.H., Hwang, S.L., Rang, M.J., et al. (2002). The hair growth promoting effect of *Sophora flavescens* extract and its molecular regulation. **Journal of Dermatological Science**, 30, 43-49.
- Rho, S.S., Park, S.J., Hwang, S.L., Lee, M.H., Kim, C.D., et al. (2005). The hair growth promoting effect of *Asiasari radix* extract and its molecular regulation. **Journal of Dermatological Science**, 38, 89-97.
- Ro, B.I., Dawson, T.L. (2005). The role of sebaceous gland activity and scalp microfloral metabolism in the etiology of seborrheic dermatitis and dandruff. **Journal of investigative Dermatology Symposium Proceeding**, 10, 194-197.
- Rochat, A., Kobayashi, K., Barrandon, Y. (1994). Location of stem cells of human hair follicles by clonal analysis. **Cell**, 76, 1063-1073.
- Rushan, X., Fei, H., Zhirong, M., Yu-Zhang, W. (2007). Identification of proteins involved in aggregation of human dermal papilla cells by proteomics. **Journal of Dermatological Science**, 48, 189-197.
- Sadagurski, M., Yakar, S., Weingarten, G., Holzenberger, M., Rhodes, C.J., et al. (2006). Insulin-like growth factor1 receptor signaling regulates skin development and inhibits skin keratinocyte differentiation. **Molecular and Cellular Biology**, 26, 2675-2687.
- Sagiroglu, A., Ozcan, H.M., Satana, A. (2009). Functional and biochemical properties of proteins from safflower seed. **Preparative Biochemistry and Biotechnology**, 39(2), 159-169.
- Savin, R.C., Atton, A.V. (1993). Minoxidil: update on its clinical role. **Dermatologic Clinics**, 11, 55-64.
- Sato, E., Tsukimoto, M., Shimura, N., Awaya, A., Kojima, S. (2011). Mechanism of pigmentation by minocycline in murine B16 melanoma cells. **The Pharmaceutical Society of Japan**, 131 (5), 731-738.
- Sawaya, M.E., Shapiro, J. (2000). Androgenetic alopecia: new approve and unapproved treatments. **Dermatologic Clinics**, 18, 47-61.

- Schlabe, J., Johnen, C., Schwartlander, R., Moser, V., Hartmann, B., et al., (2008). Isolation and culture of different epidermal and dermal cell types from human scalp suitable for the development of a therapeutical cell spray. **Burns**, 34, 376-384.
- Secco, M., Zucconi, E., Vieira, N.M., Fogaca, L.L.Q., Cerqueira, A., et al. (2008). Multipotent stem cells from umbilical cord: cord is richer than blood. **Stem Cells**, 26, 146-150.
- Sennett, R., Rendl, M. (2012). Mesenchymal-epithelial interactions during hair follicle morphogenesis and cycling. **Seminars in Cell&Developmental Biology**, 23, 917-927.
- Shan, L.Q., Ma, S., Qiu, X.C., Zhou, Y., Zhang, Y., et al. (2010). Hydroxysafflor Yellow A protects spinal cords from ischemia/reperfusion injury in rabbits. **BMC Neuroscience**, 11, 1-8.
- Shirakata, Y. (2010). Regulation of epidermal keratinocytes by growth factors. **Journal of Dermatological Science**, 59, 73-80.
- _____, Ueno, H., Hanakawa, Y., Kameda, K., Yamasaki, K., et al. (2004). TGF-beta is not involved in early phase growth inhibition of keratinocytes by 1alpha, 25 (OH)₂vitaminD₃. **Journal of Dermatological Science**, 36, 41-50.
- Sinclair, R.D. (2004). Male androgenetic alopecia. **Journal of Men's Health & Gender**, 1, 319-327.
- Singleton, V.L., Orthofer, R., Lamuela-Raventos, R.M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. **Methods in Enzymology**, 299, 152-178.
- Sinha, V.R., Kaur, M.P. (2000). Permeation enhancers for transdermal drug delivery. **Drug Development and Industrial Pharmacy**, 26 (11), 1131-1140.
- Slominski, A., Paus, R., Plonka, P., Chakraborty, A., Maurer, M., et al. (1994). Melanogenesis during the anagen-catagen-telogen transformation of the murine hair cycle. **The Journal of Investigative Dermatology**, 102, 862-869.
- _____, Wortsman, J., Plonka, P.M., Schallreuter, K.U., Paus, R., et al. 2005. Hair follicle pigmentation. **The Journal of Investigative Dermatology**, 124 (1), 13-21.

- Smitinand, T. (2001). **Thai plant names**. 2nd ed. Thailand: The Forest Herbarium Royal Forest Department.
- Sogabe, Y., Abe, M., Yokoyama, Y., Ishikawa, O. (2006). Basic fibroblast growth factor stimulates human keratinocyte motility by Rac activation. **Wound Repair and Regeneration**, 14, 457-462.
- Soma, T., Tajima, M., Kishimoto, J. (2005). Hair cycle-specific expression of versican in human hair follicles. **Journal of Dermatological Science**, 39, 147-154.
- Sripanidkulchai, B., Tattawasart, U., Laupatarakasem, P., Sripanidkulchai, K. (2001). Antimutagenicity of eight indigenous medicinal plants. **Khon Kaen University Research Journal**, 6(1), 23-33.
- Stenn, K.S., Paus, R. (2001). Controls of hair follicle cycling. **Physiological Reviews**, 81, 449-494.
- Takahashi, H., Ibe, M., Kinouchi, M., Ishida-Yamanoto, A., Hahimoto, Y., et al. (2003). Similarly potent action of 1,25-dihydroxyvitamin D3 and its analogues, tacalcitol, calcipotriol, and maxacalcitol on normal human keratinocyte proliferation and differentiation. **Journal of Dermatological Science**, 31, 21-28.
- Thody, A.J., Graham, A. (1998). Does alpha-MSH have a role in regulating skin pigmentation in human?. **Pigment Cell Research**, 11 (5), 265-274.
- Tiede, S., Kloepper, J.E., Bodo, E., Tiwari, S., Kruse, C., et al. (2007). Hair follicle stem cells: Walking the maze. **European Journal of Cell Biology**, 86, 355-376.
- Tien, Y.C., Lin, J.Y., Lai, C.H., Kuo, C.H., Lin, W.Y. et al. (2010). *Carthamus tinctorius* L. prevents LPS-induced TNF- α signaling activation and cell apoptosis through JNK1/2-NFkB pathway inhibition in H9c2 cardiomyoblast cells. **Journal of Ethnopharmacology**, 130, 505-513.
- Tobin, D.J., Hagen, E., Botchkarev, V., Paus, R. (1998). Do hair bulb melanocytes undergo apoptosis during hair follicle regression (catagen)?. **Journal of Investigative Dermatology**, 111, 941-947.
- Trueb, R.M. (2002). Molecular mechanisms of androgenetic alopecia. **Experimental Gerontology**, 37, 981-990.

- Uno, H. (1991). Quantitative models for the study of hair growth *in vivo*. **Annals of the New York Academy of Sciences**, 642, 107-124.
- Whiting, D.A. (2001). Possible mechanisms of miniaturization during androgenetic alopecia or pattern hair loss. **Journal of American Academy of Dermatology**, 45, S81-S86.
- Wilson, C.L., Sun, T.T., Lavker, R.M. (1994). Cells in the bulge of the mouse telogen follicle give rise to the lower anagen follicle. **Skin Pharmacology**, 7, 8-11.
- Won, J.H., Im, H.T., Kim, Y.H., Yum, K.J., Park, H.J., et al. (2006). Anti-inflammatory effect of buddlejasaponin IV through the inhibition of iNOS and COX-2 expression in RAW264.7 macrophages *via* the NF-kB inactivation. **British Journal of Pharmacology**, 148, 216-225.
- Yang, C.C., Cotsarelis, G. (2010). Review of hair follicle dermal cells. **Journal of Dermatological Science**, 57, 2-11.
- Yang, J.Y., Koo, J.H., Song, Y.G., Kwon, K.B., Lee, J.H., et al. (2006). Stimulation of melanogenesis by scoparone in B16 melanoma cells. **Acta Pharmacologica Sinica**, 11, 1467-1473.
- Yoo, B.Y., Shin, Y.H., Yoon, H.H., Kim, Y.J., Seo, Y.K., et al. (2009). Evaluation of human umbilical cord-derived mesenchymal stem cells on *in vivo* hair inducing activity. **Tissue Engineering and Regenerative Medicine**, 6(1-3), 15-22.
- _____, Shin, Y.H., Yoon, H.H., Seo, Y.K., Park, J.K. (2010a). Hair follicular cell/organ culture in tissue engineering and regenerative medicine. **Biochemical Engineering Journal**, 48, 323-331.
- _____, Shin, Y.H., Yoon, H.H., Seo, Y.K., Song, K.Y., et al. (2010b). Application of mesenchymal stem cells derived from bone marrow and umbilical cord in human hair multiplication. **Journal of Dermatological Science**, 60, 74-83.
- Witner, A.N., Vrensen, G.F.J.M., Noorden, C.J.F.V., Schlingemann, R.O. (2003). Vascular endothelial growth factors and angiogenesis in eye disease. **Progress in Retinal and Eye Research**, 22, 1-29.