

REFERENCES

- Ahn, B.Y. (2009). Free radical scavenging effect of ethanol extract from *Areca catechu*. **J. Applied Biol. Chem.**, **52**, 92-95.
- Ando, H., Kondoh, H., Ichihashi, M. & Hearing, V. J. (2007). Approaches to identify inhibitors of melanin biosynthesis via the quality control of tyrosinase. **J. Invest. Dermatol.**, **127**, 751-761.
- Anthikat, R. R. N. & Michel, A. (2009). Study on the areca nut for its antimicrobial properties. **J. Young Pharmacists.**, **1**, 42-45.
- Anthikat, R. R. N. & Michel, A. (2012). Anti-inflammatory and antioxidant effect of *Areca catechu*. **Inter. J. Phamaceutic. Res. Scholars.**, **3**, 2031-2037.
- Arnao, M. B., Cano, A. & Acosta, M. (2001). The hydrophilic and lipophilic contribution to total antioxidant activity. **Food Chem.**, **73**, 239-244.
- Ashawat, M. S., Shailendra, S. & Swarnlata, S. (2007). *In vitro* antioxidant activity of ethanolic extracts of *Centella asiatica*, *Punica granatum*, *Glycyrrhiza glabra* and *Areca catechu*. **Res. J. Med. Plant.**, **1**, 13-16.
- Balasundram, N., Sundram, K. & Samman, S. (2006). Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence, and potential uses. **Food Chem.**, **99**, 191-203.

- Bauer, A. W., Kirby, W. M. M., Sheris, J. C. & Turck, M. (1996). Antibiotic susceptibility testing by a standardized single disk method. **Am. J. Clin. Pathol.**, **45**, 493-496.
- Baugh, S. and Ignelzi, S. (2000). Hydrolysis and redox factors affecting analysis of common phenolic marker compounds in botanical extracts and finished products. **J. AOAC Inter.**, **83**(5), 1135-1140.
- Benzie, I. F. F. & Strain, J. J. (1999). Ferric reducing/antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. **Meth. Enzymol.**, **299**, 15-27.
- Bhandare, A. M., Kshirsagar, A. D., Vyawahare, N. S., Hadambar, A. A. & Thorve, V. S. (2010). Potential analgesic, anti-inflammatory and antioxidant activities of hydroalcoholic extract of *Areca catechu* L. nut. **Food Chem. Toxicol.**, **48**, 3412-3417.
- Cai, Y., Luo, B., Sun, M. & Corke, H. (2004). Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. **Life Sciences**, **74**, 2157-2184.
- Chang, C. C., Yang, M. H., Wem, H. M. & Chern, J. C. (2002). Estimation of total flavonoid content in propolis by two complementary colorimetric methods. **J. Food Drug Anal.**, **10**(3), 178-182.
- Chaiwut, P., Nitsawang, S., Shank, L. & Kanasawud, P. (2007). A Comparative study on properties and proteolytic components of papaya peel and latex proteases. **Chiang Mai J. Sci.**, **34**(1), 109-118.
- Chavan, Y. & Singhal, R. S. (2013a). Separation of polyphenols and arecoline from areca nut (*Areca catechu* L.) by solvent extraction, its antioxidant activity, and identification of polyphenols. **J. Sci. Food Agric.** DOI: 10.1002/jsfa.6081

- Chavan, Y. & Singhal, R. S. (2013b). Ultrasound-assisted extraction (UAE) of bioactives from arecanut (*Areca catechu*L.) and optimization study using response surface methodology. **Innov. Food Sci. Emerg.**, **17**, 106-113
- Chen ,Y., Xie, M. Y. & Gong, X. F. (2007). Microwave-assisted extraction used for the isolation of total terpenoid saponins from *Ganoderma artrum*. **J. Food Engineer.**, **81**, 162-170.
- Cheng, H. Y., Lin, T. C., Yu, K. H., Yang, C. M. & Lin, C. C. (2003). Antioxidant and free radical scavenging activities of *Terminalia chebula*. **Biol. Pharm. Bull.**, **26**, 1331-1335.
- Choi, C. W., Kim, S. C., Hwang, S. S., Choi, B. K., Ahn, H. J., Lee, M.Y., Park, S.H. & Kim, S. K. (2002). Antioxidant activity and free radical scavenging capacity between Korean medicinal plants and flavonoids by assay-guided comparison. **Plant Science**, **163**, 1161-1168.
- Chulasiri, M., Wanasawas, P., Sriaum, D., Nakamat, S., Wongkrajang, Y., Kongsaktragoon, B., Phornchirasilp, S., Songchitsomboon, S. & Leelarungrayub, D. (2011). Utilizing hydroglycolic extract fro myrobalan fruits to counteract reactive oxygen species. **Inter. J. Cosmet. Sci.**, **33**, 371-376.
- Dashen, M. M., Chollom, P. F., Okechalu, J. N. & Ma-aji, J. A. (2011). Microbiological quality assessment of some brands of cosmetics powders sold within Jos Metropolis, Plateau State. **J. Microbiol. Biotech. Res.**, **1**(2), 101-106.
- Dearden, J. C. & Forbes, W. F. (1959). Light absorption studies, Part XIV: The Ultraviolet absorption spectra of phenols. **Can. J. Chem.**, **37**, 1294-1305.
- Devasagayam, T. P. A., Boloor, K. K. & Ramasarma. (2003). Methods for estimating lipid peroxidation: An analysis of merits and demerits. **Indian J. Biochem. Biophys.**, **40**, 300-308.

- Ernest, W. F. (2001). **Cosmetic and toiletry formulations** (2nd ed., vol 8).
New York: Noyes Publication and William Andrew Publishing.
- Farah, A. & Donangelo, C. M. (2006). Phenolic compounds in coffee. **Brazil. J. Plant Physiol.**, **18**(1), 23-36.
- Flora, S. J. S. (2009). Structural, chemical and biological aspects of antioxidants for strategies against metal and metalloids exposure. **Oxid. Med. Cell Longev.**, **2**(4), 191-206.
- Friedman, M. (1997). Chemistry, biochemistry, and dietary role of potato polyphenols. A review. **J. Agric. Food Chem.**, **45**, 1523-1540.
- Friedman, M. & Jurgens, H. S. (2000) Effect of pH on the stability of plant phenolic compounds. **J. Agric. Food Chem.**, **48**, 2101-2110.
- Gao, M., Huang, W., Moytri, R. C. & Lui, C. (2007). Microwave-assisted extraction of scutellarin from *Erigeron breviscapus* Hand-Mazz and the determination by high performance liquid chromatography. **Anal. Chem. Acta.**, **591**, 161-166.
- Global Insight. (2007). **A study of the European cosmetics industry: final report**. Retrieved March 17, 2013, from http://s3.amazonaws.com/zanran_storage/
- Gunaratne, A., Wu, K., Li, D., Bentota, A., Corke, H. & Cai, Y.Z. (2013). Antioxidant activity and nutritional quality of traditional red-grained rice varieties containing proanthocyanidins. **Food Chem.**, **138**, 1153–1161.
- Halliwell, B. (2005). Free radicals and other reactive species in disease. **eLS**.

- Hamsar, M.N., Ismail, S., Mordi, M.N., Ramanathan, S. & Mansor, S.M. (2011). Antioxidant activity and the effect of different parts of areca catechu extracts on Glutathione-S-Transferase activity *in vitro*. **Free Rad. Antiox.**, **1**(1), 28-31.
- Huang, D., Ou, B. & Prior, R. L. (2005). The chemistry behind antioxidant capacity assays. **J. Agric. Food Chem.**, **53**(6), 1841-1856.
- Huang, P. L., Chi, C. W. & Liu, T. Y. (2010). Effect of *Areca catechu* L. containing procyanidins on cyclooxygenase-2 expression *in vitro* and *in vivo*. **Food Chem. Toxicol.**, **48**, 306-313.
- Kasikorn Research. (2010). เครื่องสำอางนำเข้าขายตัวอย่างต่อเนื่อง ผู้ประกอบการไทยเร่งปรับตัว. Retrieved August 14, 2011, from <http://www.kasikornreserch.com/th>
- Kaur, G. J. & Arora, D. S. (2009). Antibacterial and phytochemical screening of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi*. **BMC Compl. Altern. Med.**, **9**, 30.
- Kim, B. J., Kim, J. H., Kim, H. P. & Heo, M. Y. (1997). Biological screening of 100 plant extracts for cosmetic use (II): anti-oxidative activity and free radical scavenging activity. **Inter. J. Cosmet. Sci.**, **19**, 299–307.
- Kim, J.H., Byun, J.C., Bandi, A.K.R., Hyun, C.G. & Lee, N.H. (2009). Compounds with elastase inhibition and free radical scavenging activities from *Callistemon lanceolatus*. **J. Med. Plant. Res.**, **3**(11), 914-920.
- Kim, J. S. J. & Kim, M. J. (2010). *In vitro* antioxidant activity of *Lespedeza cuneata* methanolic extracts. **J. Med. Plants Res.**, **4**, 674-679.

- Kim, K. H., Tsao, R., Yang, R. & Cui, S. W. (2006). Phenolic acid profiles and antioxidant activities of wheat bran extracts and the effect of hydrolysis conditions. **Food Chem.**, **95**, 466-473.
- Kim, Y., Uyama, H. & Kobayashi, S. (2004). Inhibition effects of (+)-catechin-aldehyde polycondensates on proteinases causing proteolytic degradation of extracellular matrix. **Biochem. Biophys. Res. Commun.**, **320**, 26-261.
- Kiss, G.A.C., Forgacs, E., Cserhati, T., Mota, T., Morais, H. & Ramos, A. (2000). Optimization of the microwave-assisted extraction of pigments from paprika (*Capsicum annum* L.) powders. **J. Chromatogr. A.**, **889**, 41-49.
- Kumar, S., Kumar, D. & Prakash, O. (2008). Evaluation of antioxidant potential phenolic and flavonoid content of *Hibiscus tiliacius* flowers. **Elec. J. Environ. Agric. Food. Chem.**, **7**, 2863-2871.
- Kumoro, A. C., Hasan, M. & Singh, H. (2009). Effects of solvent properties on the soxhlet extraction of diterpenoid lactones from *Andrographis paniculata* leaves. **Sci. Asia.**, **35**, 306-309.
- Lee, K. K., Cho, J. J., Park, E. J. & Cho, J. D. (2001). Anti-elastase and anti-hyaluronidase of phenolic Substance from *Areca catechu* as a new anti-ageing agent. **Inter. J. Cosmet. Sci.**, **23**, 341-346.
- Lee, K. K. & Choi, J. D. (1999a). The effects of *Areca catechu* L. extract in anti-inflammation and anti-melanogenesis. **Int. J. Cosmet. Sci.**, **21**, 275-284.
- Lee, K. K. & Choi, J. D. (1999b). The effects of *Areca catechu* L. extract on anti-aging. **Int. J. Cosmet. Sci.**, **21**, 285-295.
- Lee, S. E., Hwang, H. J., Ha, J. S., Jeong, H. S. & Kim, J. H. (2003). Screening of medicinal plant extracts for antioxidant activity. **Life Sciences**, **73**, 167-179.

- Li, C. C. & Lin, E. S. (2010). Antiradical capacity and reducing power of different extraction method of *Areca catechu* seed. **Afr. J. Biotechnol.**, **9**, 7831-7836.
- Li, Y. J., Chen, J., Li, Y., Li, Q., Zheng, Y. F., Fu, Y. & Li, P. (2011). The reactive rate and the ability of the radical scavenger depend on the rate and the peak value of disappearance of the DPPH. **J. Chromatogr., A**. **1218**, 8181-8191.
- Liu, Z. Y., Hu, X. L., Bu, F. Q., Ding, L. & Zhang, H. Q. (2007). Studies on the chemical change in the process of microwave-assisted extraction of flavonoids from *Acanthpanax senticosus* harms. **Chem. J. Chinese Univ.**, **28**, 431-435.
- Mandal, V., Mohan, Y. & Hemalatha, S. (2007). Microwave assisted extraction- An innovative and promising extraction tool for medicinal plant research. **J. Phamaceut. Biomed. Anal.**, **46**,322-327.
- Marketing Oops. (2008). ส่วนแบ่งตลาดผลิตภัณฑ์ดูแลเส้นผม ยาสีฟัน และโลชั่นบำรุงผิว.
Retrieved March 17, 2013, from [www. marketingoops.com/ reports/ matrix/market-share-2/](http://www.marketingoops.com/reports/matrix/market-share-2/)
- Moon, J. Y., Yim, E. Y., Song, G., Lee, N. H. & Hyun, C. G. (2010). Screening of elastase and tyrosinase inhibitory activity from Jeju Island plants. **EurAsia. J. BioSci.**, **4**, 41-53.
- Nelson, B. S. & Heischober, B. (1999). Betel nut: a common drug used by naturalized citizens from India, far east Asia, and the south Pacific islands. **Ann. Emerg. Med.**, **34**(2), 238-243.
- Norton, S. C. (1998). Betel: Consumption and consequences. **J. Am. Acad. Dermatol.**, **38**(1), 81-88.

- Onar, H. C., Yusufoglu, A., Turker, G. & Yanardag, R. (2012). Elastase, tyrosinase and lipoxygenase inhibition and antioxidant activity of an aqueous extract from *Epilobium angustifolium* L. leaves. **J. Med. Plants Res.**, **6**, 716-726.
- Oxenham, M. F., Locher, C., Cuong, N. L. & Thuy, N. K. (2002). Identification of *Areca catechu* (betel nut) residues on the dentitions of bronze age inhabitants of Nui Nap, Northern Vietnam. **J. Archaeol. Sci.**, **29**, 909–915.
- Patel, A., Patel, A., Patel, A. & Patel, N. M. (2010). Estimation of flavonoid, polyphenolics content and *in vitro* antioxidant capacity of leaves of *Tepharosia purpurea* Linn. (Leguminosae). **Int. J. Pharma Sci. Res.**, **1**(1), 66-77.
- Proestos, C. & Komaitis, M. (2008). Application of microwave-assisted extraction to the fast extraction of plant phenolic compounds. **LWT**, **41**, 652–659.
- Rice-Evans, C., Miller, N. & Paganga, G. (1996). Structure-antioxidant activity relationships of flavonoids and phenolic acids. **Free Radic. Biol. Med.**, **20**(7), 933-956.
- Robbins, R. J. (2003). Phenolic acids in foods: An overview of analytical methodology. **J. Agric. Food Chem.**, **51**, 2866-2887.
- Sani, I. M., Iqbal, S., Chan, K. W. & Ismail, M. (2012). Effect of acid and base catalyzed hydrolysis on the yield of phenolics and antioxidant activity of extracts from germinated brown rice (GBR). **Molecules.**, **17**, 7584-7594.
- Seo, S.Y., Sharma, V.K. & Sharma, N. (2003). Mushroom Tyrosinase: recent prospects. **J. Agric. Food Chem.**, **51**, 2837-2853.
- Shalini, S. (2012). Natural antioxidants- a review. **Inter. J. Phytotherapy.**, **2**, 7-15.

- Silva, M. C. A. & Paiva, S. R. (2012). Antioxidant activity and flavonoid content of *Clusia fluminensis* Planch. and Triana. **An. Acad. Bras. Cienc.**, **84**(3), 609-616.
- Singleton V. L., Orthofer R. & Lamune-Raventos R. M. (1999). Analysis of total phenols and other oxidation substance and antioxidants by means of Folin-Ciocalteu reagent. **Meth. Enzymol.**, **299**,152-17
- Staples, G. W. & Bevacqua, R. F. (2006). *Areca catehu* (**Betel nut palm**). Retrieved August 1, 2011, from <http://www.agroforestry.net/tti/Areca-catechu-betel-nut.pdf>
- Sultana, B., Anwar, F. & Ashraf, M. (2009). Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. **Molecules.**, **14**, 2167-2180.
- Sun, B., Ricardo-da-Silva, J. M., & Spranger, I. (1998). Critical factors of vanillin assay for catechins and proanthocyanidins. **J. Agric. Food Chem.**, **46**, 4267-4274.
- Taylor, W. P., Hamilton-Miller, M. T. J. & Stapleton, D. P. (2009). Antimicrobial properties of green tea catechins. **Food Sci. Technol. Bull.**, **2**, 71-81.
- Thai Bureau of Cosmetics and Hazardous Substance. (2004). นวัตกรรมผลิตภัณฑ์เครื่องสำอาง ศูนย์ข้อมูลเครื่องสำอาง. Retrieved August 14, 2011, from <http://web.dmsc.moph.go.th>
- Thai Department of Agriculture Extension. (2002). อนาคตหมากไทยจะเป็นอย่างไร. Retrieved August 10, 2011, from <http://www.doae.go.th/library/html/detail/>

- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L. & Byrne, D. H. (2006). Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. **J. Food. Compos. Anal.**, **19**, 669-675.
- Thai Ministry of Commerce. (2012). มูลค่าสินค้าออก สินค้าเข้า และดุลการค้าของไทย. Retrieved March 10, 2013, from www2.ops3.moc.go.th
- Thring, T. S. A., Hili, P. & Naughton, D. P. (2009). Anti-collagenase, anti-elastase and anti-oxidant activities of extracts from 21 plants. **BMC Complem. Altern. M.**, **9**(27), 1-11.
- Tomsone, L., Kruma, Z. & Galoburda, R. (2012). Comparison of different solvents and extraction methods for isolation of phenolic compounds from horseradish roots (*Armoracia rusticana*). **World Acad. Sci.**, **64**, 903-908.
- Uchino, K., Matsuo, T. & Iwamoto, M. (1988) New 5'-nucleotidase inhibitors, NPF-86IA, NPF-86IB, NPF-86IIA, and NPF-86IIB from *Areca catechu*. II. Anti-tumor effects. **Planta Med.**, **54**(5), 419-422.
- Vrhovsek, U., Palchetti, A., Reniero, F., Guillou, C., Masuero, D. & Mattivi, F. (2006). Concentration and mean degree of polymerization of *Rubus* ellagitannins evaluated by optimized acid methanolysis. **J. Agric. Food Chem.**, **54**, 4469-4475.
- Wang, C. K. & Lee, W. H. (1996). Separation, characteristics, and biological activities of phenolics in *Areca* fruit. **J. Agric. Food Chem.**, **44**, 2014-2019.
- Wang, C. K., Lee, W. H. & Peng, C. H. (1997). Contents of phenolics and alkaloids in *Areca catechu* Linn. during maturation. **J. Agric. Food Chem.**, **45**, 1185-1188.

- Wang, L. & Weller, C. L. (2006). Recent advances in extraction of nutraceuticals from plants. **Trends in Food Sci. Technol.**, **17**, 300-312.
- Wells, P. G., Bhuller, Y., Chen, C. S., Jeng, W., Kasapinovic, S., Kennedy, J. C., Kim, P. M., Laposa, R. R., McCallum, G. P., Nicol, C. J., Parman, T., Wiley, M. J. & Wong, A. W. (2005). Molecular and biochemical mechanisms in teratogenesis involving reactive oxygen species. **Toxicol. Applied Pharmacol.**, **207**(2), 354-366.
- Wetwitayaklung, P., Phaechamud, T., Limmatvapirat, C. & Keokitichai, S. (2006). The study of antioxidant capacity in various parts of *Areca catechu* L. **Naresuan Univ. J.**, **14**, 1-14.
- Wu, Q., Yang, Y. & Simon, J. E. (2007). Qualitative and quantitative HPLC/MS determination of proanthocyanidins in Areca nut (*Areca catechu*). **Chem. Biodiverse.**, **4**(12), 2817-2826.
- Yenjit, P., Issarakraisila, M., Intana, W. & Chantrapromma, K. (2010). Fungicidal activity of compounds extracted from the pericarp of *Areca catechu* against *Colletotrichum gloeosporioides in vitro* and in mango fruit. **Postharvest Biol. Technol.**, **55**, 129–132.
- Zhang, H. F., Yang, X. H. & Wang, Y. (2011). Microwave assisted extraction of secondary metabolites from plants: Current status and future direction. **Trends Food Sci. Technol.**, **21**, 672-688.
- Zhang, W. M., Li, B., Han, L. & Zhang, H. D. (2009). Antioxidant activities of extract from Areca (*Areca catechu* L.) flower, husk, and seed. **Elec. J. Environ. Agric. Food. Chem.**, **8**, 740-748.
- Zhang, X., Wu, J., Han, Z., Mei, W. L. & Dai, H. F. (2010). Antioxidant and cytotoxic phenolic compounds of areca nut (*Areca catechu*). **Chem. Res. Chinese Univer.**, **26**, 161-164.

Zhu, X., Su, Q., Cai, J. & Yang, J. (2006). Optimization of microwave-assisted solvent extraction for volatile organic acids in tobacco and its comparison with conventional extraction methods. **Analytica Chimica Acta.**, **579**, 88-94.