

เอกสารอ้างอิง

จารุณี เกสรพิกุล และสุรวัฒน์ ชลอสันติสุข. คู่มือปฏิบัติการจุลชีววิทยาสำหรับนักศึกษาวิทยาศาสตร์การเกษตร (700201). 2554;คณะสัตวศาสตร์และเทคโนโลยีการเกษตร มหาวิทยาลัยศิลปากร วิทยาเขตสารสนเทศเพชรบุรี, เพชรบุรี.

คันสนีย์ ชีระพันธ์. ความปลอดภัยทางชีวภาพในห้องปฏิบัติการจุลชีววิทยา. วารสารกรมวิทยาศาสตร์บริการ 2555;60:15-18

Alper H, Moxley J, Nevoigt E, Fink GR, Stephanopoulos G. Engineering yeast transcription machinery for improved ethanol tolerance and production. Science 2006;314:1565-1568.

Andersen G, Björnberg O, Polakova S, Pynyaha Y, Rasmussen A, Møller K, Hofer A, Moritz T, Sandrin MPB, Merico AM, Compagno C, Åkerlund HE, Gojkovic Z, Piškur J. A second pathway to degrade pyrimidine nucleic acid precursors in eukaryotes. J Mol Biol 2008;380:656-666.

Arasu MV, Kumar V, Ashok S, Song H, Rathnasingh C, Lee HJ, Seung D, Park S. "Isolation and characterization of the new *Klebsiella pneumonia* J2B strain showing improved growth characteristics with reduced lipopolysaccharide formation." Biotechnol Bioprocess Eng 2011;16:1134-1143.

Ashok S, Raj SM, Ko Y, Sankaranarayanan M, Zhou S, Kumar V, Park S. Effect of nitrate on glycerol metabolism and anaerobic production of 3-hydroxypropionic acid by recombinant *Klebsiella pneumoniae* overexpressing *puuC*. Metab Eng 2013;15:10-24.

Behr A, Botulinski A, Carduck F-j, Schneider M. Process for preparing 3-hydroxypropionic acid. 1996;Patent application no. EP0579617.

Biebl H, Menzel K, Zeng AP, Deckwer WD. Microbial production of 1,3-propanediol. Appl Microbiol Biotechnol 1999;52:289-297.

Booth IR, Cash P, O'Byrne C. Sensing and adapting to acid stress. *Antonie Van Leeuwenhoek Int J Gen Mol Microbiol* 2002;81:33-42.

Chang YH, Kim JK, Kim HJ, Kim WY, Kim YB, Park YH. Selection of a potential probiotic *Lactobacillus* strain and subsequent in vivo studies. *Antonie Van Leeuwenhoek*, 2001;80:193-199.

Dunlop MJ. Engineering microbes for tolerance to next generation biofuel. *Biotechnol Biofuels* 2011;4:32.

Fuchs G. Alternative pathways of carbon dioxide fixation: insights into the early evolution of life? *Annu Rev Microbiol* 2011;65:631-658.

Garai-Ibabe G, Ibarburu I, Berregi A, Claisse O, Lonvaud-Funel A, Irastorza A, Dueñas MT. Glycerol metabolism and bitterness producing lactic acid bacteria in cidermaking. *Int J Food Microbiol* 2008;121:253-261.

Haas T, Brossmer C, Meier M, Arntz D, Freund A. Process for preparing 3-hydroxypropionic acid or its salt. 2000;Patent application no. EP0819670.

Henry CS, Broadbelt LJ, Hatzimanikatis V. Discovery and analysis of novel metabolic pathways for the biosynthesis of industrial chemicals: 3-hydroxypropanoate. *Biotechnol Bioeng* 2010;106:462-473.

Holo H. *Chloroflexus aurantiacus* secretes 3-hydroxypropionate, a possible intermediate in the assimilation of CO₂ and acetate. *Arch Microbiol* 1989;151:252-256.

Huang Y, Li Z, Shimizu K, Ye Q. Co-production of 3-hydroxypropionic acid and 1,3-propanediol by *Klebsiella pneumoniae* expressing aldH under microaerobic conditions. *Bioresour Technol* 2013;128:505-512.

Hügler M, Huber H, Stetter KO, Fuchs G. Autotrophic CO₂ fixation pathways in archaea (Crenarchaeota). *Arch Microbiol* 2003;179:160-173.

Jarboe LR, Liu P, Royce LA. Engineering inhibitor tolerance for the production of biorenewable fuels and chemicals. *Curr Opin Chem Eng* 2011;1:38-42.

Jiang X, Meng X, Xian M. Biosynthetic pathways for 3-hydroxypropionic acid production. *Appl Microbiol Biotechnol* 2009;82:995-1003.

Johnson DT, Taconi KA. 2007. The glycerin glut: options for the value-added conversion of crude glycerol resulting from biodiesel production. *Environ Prog* 2007;26:338-348.

Krauter H, Willke T, Vorlop K-D. Production of high amounts of 3-hydroxypropionaldehyde from glycerol by *Lactobacillus reuteri* with strongly increased biocatalyst lifetime and productivity. *N Biotechnol* 2012;29(2):211-217.

Kumar V, Mugesh S, Jae K, Durgapal M, Ashok S, Ko Y, Sarkar R, Park S. Co-production of 3-hydroxypropionic acid and 1,3-propanediol from glycerol using resting cells of *Klebsiella pneumoniae* J2B strain with overexpression of KGSADH. *Appl Microbiol Biotechnol* 2012;96:373–83.

Kumar V, Ashok S, Park Sunghoon. Recent advances in biological production of 3-hydroxypropionic acid. *Biotechnol Adv* 2013;31:945-961.

Leal NA, Havemann GD, Bobik TA. PduP is a coenzyme-A-acylating propionaldehyde dehydrogenase associated with the polyhedral bodies involved in B12-dependent 1, 2-propanediol degradation by a *Salmonella enterica* serovar Typhimurium LT12. *Arch Microbiol* 2003;180:353-361.

Liu YN, Leal A, Sampson EM, Johnson CL, Havemann GD, Bobik TA. PduL is an evolutionarily distinct phosphotransacylase involved in B12-dependent 1,2-propanediol degradation by *Salmonella enterica* serovar Typhimurium LT2. *J Bacteriol* 2007;189:1589-1596.

Luo LH, Seo JW, Baek JO, Oh BR, Heo SY, Hong WK, Kim DH, Kim CH. Identification and characterization of the propanediol utilization protein PduP of *Lactobacillus reuteri* for 3-hydroxypropionic acid production from glycerol. *Appl Microbiol Biotechnol* 2011;89:697-703.

Luo LH, Kim CH, Heo SY, Oh BR, Hong WK, Kim S, Kim DH, Seo JW. Production of 3-hydroxypropionic acid through propionaldehyde dehydrogenase PduP mediated biosynthetic pathway in *Klebsiella pneumoniae*. *Bioresour Technol* 2012;103:1-6.

Mavrovouniotis ML. Duality theory for thermodynamic bottlenecks in bioreaction pathways. *Chem Eng Sci* 1996;51:1495-507.

Meng XS, Abraham T, Tsobanakis P. Process for preparing 3- hydroxycarboxylic acids. 2007;Patent application no. 20070015936.

Pina CD, Falletta E, Rossi M. A green approach to chemical building blocks. The case of 3-hydroxypropanoic acid. *Green Chem* 2011;13:1624-32.

Raj SM, Rathnasingh C, Jo JE, Park S. Production of 3-hydroxypropionic acid from glycerol by a novel recombinant *Escherichia coli* BL21 strain. *Process Biochem* 2008;43:1440-1446.

Rathnasingh C, Raj SM, Jo JE, Park SH. Development and evaluation of efficient recombinant *Escherichia coli* strains for the production of 3-hydroxypropionic acid from glycerol. *Biotechnol Bioeng* 2009;104:729-739.

Rathnasingh C, Raj SM, Lee Y, Catherine C, Ashok S, Park S. Production of 3-hydroxypropionic acid via malonyl-CoA pathway using recombinant *Escherichia coli* strains. *J Biotechnol* 2012;157(4):633-640.

Roe AJ, O'Byrne C, McLaggan D, Booth IR. Inhibition of *Escherichia coli* growth by acetic acid: a problem with methionine biosynthesis and homocysteine toxicity. *Microbiology* 2002;148:2215-2222.

Saini R, Kapoor R, Kumar R, Siddiqi TO, Kumar A. CO₂ utilizing microbes - a comprehensive review. *Biotechnol Adv* 2011;29:949-960.

Sneath PHA. Endospore-forming Gram-positive Rods and Cocci. In *Bergey's Manual of systematic Bacteriology*, Vol II, 1986;Williams & Wilkins, Baltimore.

Sobolov M, Smiley FA. Metabolism of glycerol by an acrolein-forming *Lactobacillus*. J Bacteriol 1960;79:261-266.

Tabuchi T, Sugisawa T, Ishidori T, Nakahara T, Sugiyama J. Itaconic acid fermentation by a yeast belonging to the genus *Candida*. Agric Biol Chem 1981;45:475-479.

Takamizawa K, Horitsu H, Ichikawa T, Kawai K, Suzuki T. β -hydroxypropionic acid production by *Byssoschlamys* sp. grown on acrylic acid. Appl Microbiol Biotechnol 1993;40:196-200.

Talarico TD, Casas IA, Chung TC, Dobrogosz WJ. Production and isolation of reuterin, a growth inhibitor produced by *Lactobacillus reuteri*. Antimicrob Agents Chemother 1988;32:1854-1858.

Valdehuesa KNG, Liu H, Nisola GM, Chung W-J, Lee SH, Park SJ. Recent advances in the metabolic engineering of microorganisms for the production of 3-hydroxypropionic acid as C3 platform chemical. Appl Microbiol Biotechnol 2013;97:3309–3321

van Maris AJA, Konings WN, Van Dijken JP, Pronk JT. Microbial export of lactic and 3-hydroxypropanoic acid: implications for industrial fermentation processes. Metab Eng 2004;6(4):245-255.

Warnecke T, Gill RT. Organic acid toxicity, tolerance, and production in *Escherichia coli* biorefining applications. Microb Cell Fact 2005;4:425.

Warnecke TE, Lynch MD, Lipscomb ML, Gill RT. Identification of a 21 amino acid peptide conferring 3-hydroxypropionic acid stress-tolerance to *Escherichia coli*. Biotechnol Bioeng 2012;109(5):1347-1352.

Werpy T, Petersen G. Top value added chemicals from biomass, vol 1: results of screening for potential candidates from sugars and synthesis gas. US Department of Energy; 2004 [<http://www.osti.gov/bridge>].

Xue J, Murrieta CM, Rule DC, Miller WK. Exogenous or L-rhamnose-derived 1,2-propanediol is metabolized via a pduD-dependent pathway in *Listeria innocua*. Appl Environ Microbiol 2008;74:7073-7079.

Yasuda S, Mukoyama M, Horikawa H, Toraya T, Morita H. Process for producing 1,3-propanediol and or/3-hydroxypropionic acid. US Patent 2007,0,148,749,2007.

Zeng AP, Biebl H. Bulk chemicals from biotechnology: the case of 1,3-propanediol production and the new trends. Adv Biochem Eng Biotechnol 2002;74:239-259.

Zhang D, Hillmyer MA, Tolman WB. A new synthetic route to poly[3-hydroxypropionic acid] (P[3-HP]): ring-opening polymerization of 3-HP macrocyclic esters. Macromolecules 2004;37:8198-8200.

Zhu B, Li J, He Y, Yamane H, Kimura Y, Nishida H, Inoue Y. Effect of steric hindrance on hydrogen-bonding interaction between polyesters and natural polyphenol catechin. J Appl Polym Sci 2004;91:3565-3573.