

REFERENCES

- Ahmed E.M. Abdalla , Saeid M. Darwish, Eman H.E. Ayad, Reham M. El-Hamahmy. (2007). Egyptian mango by-product 2. Antioxidant and antimicrobial activities of extract and oil from mango seed kernel. Food Chemistry. 103, 1141–1152.
- Ahmed E.M. Abdalla , Saeid M. Darwish, Eman H.E. Ayad, Reham M. El-Hamahmy. (2007). Egyptian mango by-product1. Compositional quality of mango seed kernel. Food Chemistry. 103, 1134-1140.
- Ahmed I, Aquil F, Owais M. (2006). Modern phytomedicine; turning medicinal plants into drugs. Weinheim, Wiley-VCH Verlag GmbH&co KGaA.
- Ajila,C.M., Naidu, K.A., Bhat, S.G., Prasada Rao, U.J.S. (2007). Bioactive compounds and antioxidant potential of mango peel extract. Food Chemistry. 105, 982–988.
- Anwar F., Jamil A., Iqbal S., Sheikh M.A. (2006). Antioxidant activity of various plant extracts under ambient and accelerated storage of sunflower oil. Grasas Aceites Sevilla, 57, 189-197.
- Arogba SS. (1997). Physical, chemical and functional properties of nigerian mango (*Mangifera indica*) kernel and its processed Flour. J Sci Food Agric. 73, 321-8.
- Arogba SS. (1999). The performance of processed mango (*Mangifera indica*) kernel flour in a model food system. Bioresource Technol. 70, 277-81.
- Arogba SS. (2000).Mango kernel: chromatographic analysis of the tannin and stability study of the associated polyphenols oxidase activity. J Food Compos Anal. 13, 149-56.
- Babbar, N., Oberoi H.S., Uppal, D.S., Patil, R.T. (2011). Total phenolic content and antioxidant capacity of extracts obtained from six important fruit residues. Food Research International. 44, 391-396.

- Blois, M. S. (1958). Antioxidant determinations by the use of a stable free radical. *Nature*. 181, 1199–1200.
- Campos, L.M.A.S., Leimann, F.V., Pedrosa, R.C., Ferreira, S.R.S. (2008). Free radical scavenging of grape pomace extracts from Cabernet Sauvignon (*Vitis vinifera*). *Bioresource Technology*. 99, 8413-8420.
- Chedea, V. S., Braicu, C. and Socaciu, C. (2010). Antioxidant/prooxidant activity of a polyphenolic grape seed extract. *Food Chemistry*. 121, 132–139.
- Cheng, Z.; Su, L.; Moore, J.; Zhou, K.; Luther, M.; Yin, J.J.; Yu, L.L. (2006). Effect of postharvest treatment and heat stress on availability of wheat antioxidants. *J. Agric. Food Chem.* 54, 5623-5629.
- Crews C. (2002). Authentication of cocoa butter; in oils and fats authentication. Michael J, editor. New York: Blackwell Publishing Ltd. p.66-89.
- Department of Agriculture. Thailand. (2008). [monograph on the Internet]. [cited 2011 Aug 10]. Available from: <http://it.doa.go.th/vichakan/news.php?newsid=37>
- Duangnak P. (2004). Antioxidant analysis. [monograph on the Internet]. [cited 2011 Mar 13]. Available from: <http://pirun.ku.ac.th/~b4755242/7.htm>
- Dubost, NJ., Ou, B., Beelman BR. (2007). Quantification of polyphenols and ergithioneine in cultivated mushrooms and correlation to total antioxidant capacity. *Food Chem.* (article in press)
- Fox, C. (1997). Advance in the cosmetics science and technology of topical bioactive materials. *Cosmet Toilet Magazine*. 112, 67-84.
- Gunstone, FD. (2002). Production and trade of vegetable oil. In: *Vegetable oils in food technology, composition, properties and uses*. Oxford: Blackwell Publishing Ltd. p.1-17.
- Gunstone, FD., Padley, FB. (1997). *Lipid technologies and applications*. New York: Marcel Dekker Inc.
- Gutteridge, JMC., Halliwell, B. (1996). *Antioxidants in nutrition, health, and disease*. Oxford: Oxford University Press.
- Halliwell B, Gutteridge JMC. (1984). Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem J*. 219, 1–14.

- Hatano, T., Edamatsu, R., Mori, A. (1989). Effect of interaction of tannins and related polyphenols on superoxide anion radical and on DPPH radical. *Chem Pharm Bull.* 37, 2016-2021.
- Hazra, B., Biswas, S., Mandal N. (2008). Antioxidant and free radical scavenging activity of *Spondias pinnata*. *BMC Complementary and Alternative Medicine.* 8:63.
- Hertog, MGL., Katan, MB. (1998). Quercetin in foods, cardiovascular disease, and cancer. New York: Marcel Dekker.
- Inprakhon P. (2009). Mango seed kernel. Science for Development; Academic articles collected for 50th anniversary of Faculty of Science. Bangkok: Mahidol University. 153-162.
- Inrod, P. (2008). Antioxidant activities and total phenolic contents of *Etlingera paviana* and *Amomum biflorum*. Chonburi : Faculty of science, Burapha University.
- Jamuna, S., Paulsamy, S., Karthika, K. (2012). Screening of in vitro antioxidant activity of methanolic leaf and root extracts of *Hypochaeris radicata* L. (Asteraceae). *Journal of Applied Pharmaceutical Science.* 02(07), 149-154.
- Joseph, JK., Abolaji, J. (1997). Effect of replacing maize with graded levels of cooked Nigerian mango seed kernels (*Mangifera indica*) on the performance, carcass yield and meat quality of broiler chickens. *Bioresource Technol.* 61, 99-102.
- Kähkönen, M.P., Hopia, A.I., Vuorela, H.J., Rauha, J., Pihlaja, P.K., Kujala T.S., Heinonen, M. (1999). Antioxidant Activity of Plant Extracts Containing Phenolic Compounds. *Journal of Agricultural and Food Chemistry.* 47, 3954-3962.
- Katalinic, V., Mozina, S.S., Skroza, D., Generalic, I., Abramovic, H., Milos, M., Ljubenkovic, I., Piskernik, S., Pezo, I., Terpin, P., Boban, M. (2010). Polyphenolic profile, antioxidant properties and antimicrobial activity of grape skin extracts of 14 *Vitis vinifera* varieties grown in dalmatia (Croatia). *Food Chemistry.* 119, 715-723.

- Khammuang, S., Sarnthima, R. (2011). Antioxidant and antibacterial activities of selected varieties of Thai mango seed extract. *Pak.J.Pharm.Sci.* 24(1), 37-42.
- Khunthajaroen, T., Kerdchoechuen, O., Laohakunjit, N. (2010). Phenolic Compounds from Peel and Seed Extracts of ‘White Malaka’ Grape. *Agricultural Sci. J.* 41(3/1)(Suppl.), 613-616.
- Lafka, T.I., Sinanoglou, V. and Lazos, E.S. (2007). On the extraction and antioxidant activity of phenolic compounds from winery waste. *Food Chemistry.* 104, 1206-1214.
- Laopatarakasem P. (2004). Roles of Natural products to cancer prevention and treatment. Khon Kaen: Department of Physiology, Faculty of Medicine, Khon Kaen University.
- Lee, J., Koo, N., & Min, D. B. (2004). Reactive oxygen species, aging and antioxidative nutraceuticals. *Comprehensive Reviews in Food Science and Food Safety.* 2, 21–33.
- Lercker, G., Rodriguez-Estrada, MT. (2000). Chromatographic analysis of unsaponifiable compound of olive oils and fat-containing foods. *J Chromat A.* 881, 105-29.
- Maier, T., Schieber, A., Kammerer, D.R. and Carle, R. (2009). Residues of grape (*Vitis vinifera* L.) seed oil production as a valuable source of phenolic antioxidants. *Food Chemistry.* 112, 551-559.
- Maisuthisakul, P., Gordon, M.H. (2009). Antioxidant and tyrosinase inhibitory activity of mango seed kernel by product. *Food Chemistry.* 117, 332–341.
- Manach, C., Scalbert, A., Monrand, C., Remesy, C., Jimenez, L. (2004). Polyphenol: food sources and bioavailability. *Am. J. Clin Nutr.* 79, 727-47.
- Manach, C., Williamson, G., Morand, C., Scalbert, A., & Remesy, C. (2005). Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *The American Journal of Clinical Nutrition.* 81(Suppl.), 230S–242S.
- Manmohan, S., Arindam, P., Hemendra, P.S., Sushil, K.D., Kalpesh, G. (2011). Evaluation of Reducing Power Assay of Chalcone Semicarbazones. *J. Chem. Pharm. Res.* 3(3), 639-645.

- Masibo, M., He, Q. (2008). Major mango polyphenols and their potential significance to human health. *Comp Rev Food Sci Food Safety*. 7, 309-19.
- Nakajima, H., Tadokoro, S., Hashiba, H., Ito, F., Furuya, H., Kabuki, T., Arai, M., Dosako, S. (2000). Bacteriostatic and antibacterial agent containing mango kernel component. U.S. Patent No. 6,063,382. Washington, DC: U.S. Patent and Trademark Office.
- Nilnond, S. (2001). Grape Production in Thailand. In *Grape Production in the Asia-Pacific Region*. Bangkok. RAP Publication. 70-79.
- Office of Agricultural Economics, Ministry of Agriculture and Cooperatives, (2013). Basic information of agricultural economics 2013. 402, 50-51.
- Omoregie, E., Ufodike, EBC., Umara, MS. (1991). Growth and feed utilization of *Oreochromis niloticus* fingerlings fed diets containing cassava peeling and mango seeds. *Aquabute*. 4, 6-7.
- Oyaizu, M. (1986). Studies on products of the browning reaction. Antioxidative activities of browning reaction products prepared from glucosamine. *Eiyogaku Zasshi*. 44, 307–315.
- Pedreschi, R., & Cisneros-Zevallos, L. (2006). Antimutagenic and antioxidant properties of phenolic fractions from Andean purple corn (*Zea mays* L.). *J Agric Food Chem*. 54(13), 4557-4567.
- Puravankara, D., Bohgra, V., & Sharma, R. S. (2000). Effect of antioxidant principles isolated from mango (*Mangifera indica* L.) seed kernels on oxidative stability of buffalo ghee (butter-fat). *Journal of the Science of Food and Agriculture*, 80, 522–526.
- Rao, YV., Das, BK., Pradhan, J., Chakrabarti, R. (2006). Effect of *Achyranthes aspera* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. *Fish Shellfish Immunol*. 20, 263-73.
- Ravindran, V., and Sivakanesan, R. (1995). The Nutritive Value of Mango Seed Kernels for Starting Chicks. *J Sci Food Agric*. 71, 245-250.
- Reddy, SY., Jeyarani, T. (2001). Trans-free bakery shortening from mango kernel and mahua fats by fractionation and blending. *J Am Oil Chem Soc*. 78, 635-40.

- Ribeiro, S. M. R., Queiroz, J. H., Queiroz, M. E. R. L., Campos, F. M., Pinheiro-Sant'Ana, H. M. (2007). Antioxidants in mango (*Mangifera indica*, L.) pulp. *Plant Foods for Human Nutrition*. 62, 13–17.
- Ribeiro, S.M.R., Barbosa, L.C.A., Queiroz, J.H., Knödler, M. Schieber, A. (2008). Phenolic compounds and antioxidant capacity of Brazilian mango (*Mangifera indica* L.) varieties. *Food Chemistry*. 110, 620–626.
- Ribéreau-Gayon, P., Glories, Y., Maujean, A. and Dubourdieu, D. (2000). The chemistry of wine stabilization and treatments. In: *Handbook of enology*. Vol. 2. 2nded. John Wiley & Sons, Chichester.
- Rukmini, C., Vijayaraghavan, M. (1984). Nutritional and toxicological evaluation of mango kernel.
- Sahu S, Kumar Das B, Pradhan J, Mohapatra BC, Mishra BK, Sarangi N. (2007). Effect of *Mangifera indica* kernel as feed additive on immunity and resistance to *Aeromonas hydrophila* in *Labeo rohita* fingerlings. *Fish Shellfish Immunol*. 23, 109-18.
- Scalzo, J., Politi, A., Pellegrini, N., Mezzetti, B., Battino, M. (2005). Plant genotype affects total antioxidant capacity and phenolic contents in fruit. *Nutrition*. 21, 207–213.
- Shahidi F, Naczki M. (1951). *Phenolics in food and nutraceuticals*. Florida: CRC press.
- Shih, PW., Lai, PL., Jen, HWK. (2006). Antioxidant activity of aqueous extracts of selected plants. *Food Chem*. 99, 775-783.
- Siddhuraju, P.; Becker, K. Antioxidant properties of various extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *J. Agric. Food Chem*. 2003, 51, 2144-2155.
- Singleton, VL., Rossi, JA. (1965). Colorimetry of total phenolics with phosphomolybdic acid-phosphotungstic acid reagents. *Am. J. Enol. Viticult*. 16, 144-158.
- Songchitsomboon, S. (2005). How importance of Antioxidant for health. Thailand: *Folk Doctor Magazine*. 316.
- Sultana, B., Anwar, F., Przybylski, R. (2009). Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. *Molecules*. 14, 2167-2180.

- Sultana, B.; Anwar, F.; Przybylski, R. (2007). Antioxidant activity of phenolic components present in barks of *Azadirachta indica*, *Terminalia arjuna*, *Acacia nilotica*, and *Eugenia jambolana* Lam. trees. *Food Chem.* 104, 1106-1114.
- Samavardhana, K., Supawitipattana, P., Jittrepotch, N., Rojsuntornkitti, K., Kongbangkerd, T. (2015). Effects of extracting conditions on phenolic compounds and antioxidant activity from different grape processing byproducts. *International Food Research Journal.* 22(3), 1169-1179.
- Suwanaalert R. (2009). Mango. [monograph on the Internet]. [cited 2011 Aug 10]. Available from: <http://clgc.rdi.ku.ac.th/index.php/resource/62-fruit/132-manaifera>
- The Society for Free Radical Research-Thai. (2013). Free Radicals and Antioxidants. Health Innovation Publisher. 450p.
- Velan, M., Krishanan, RV., Lakshmanan, CM. (1995). Conversion of mango kernel starch to glucose syrups by enzymatic hydrolysis. *Bioprocess Bioeng.* 12, 323-6.
- Villano, D., Fernandez-Pachon, MS., Moya, ML., Troncoso, AM., Garcia-Parrilla, MC. (2007). Radical scavenging ability of polyphenolic compounds towards DPPH free radical. *Talanta*: 71, 230-235.
- Wang, SY., Zheng W., Galletta, GJ. (2002). Cultural system affects fruit quality and antioxidant capacity in strawberries. *J. Agric. Food Chem.* 50, 6534-42.
- Xu, BJ., Chang, SKC. (2007). A comparative study on phenolic profiles and antioxidant activities of legumes as affected by extraction solvents. *J food sci.* 72, S159-S166.
- Yildirim, A., Mavi, A., Oktay, M., Kara, AA., Algur, OF., Bilaloglu, V. (2002). Comparison of antioxidant and antimicrobial activities of *Tilia* (*Tilia argentea* Desf Ex DC), *Sage* (*Savia triloba* L.), and *Black Tea* (*Camellia sinensis*). *J. Agric. Food Chem.* 48(10), 5030-5034.