

ACKNOWLEDGEMENT

I would like to give many thanks to my advisor, Dr. James True. He invested a lot of energy to improve my thesis. He gave me a lot of opportunities to attend both international and Thai conferences and also some international workshops. He was showing me how to be a good scientist and, crucially, he improved my scientific writing.

I am thankful to my committees — both proposal and thesis defense — who gave a lot of valuable suggestions and make my work reasonable. These include Dr. Wansuk Senanan from Department of Aquatic Sciences, Burapha University, who was my Thesis Defense Committee chairperson, Dr. Singtoe Boonrojpong from Department of biology, Faculty of Science, Prince of Songkla University, who was a member of my committee of Thesis Defense, Dr. Jaros Phinchongsakuldit from School of Science, Walailak University and Asst. Prof. Dr. Vachira Lhekchim from Department of Biology, Faculty of Science, Prince of Songkla University who were my proposal defense committee.

I am also grateful to Dr. Jaros Phinchongsakuldit, who permitted me to do molecular work at the Biomolecular Laboratory, Biodiversity Research Unit, Walailak University, Nakhon Si Thammarat province. She gave me a lot of experience in molecular work. I also thank all the staff and graduate students at Walailak University for their help and support for me during the period I worked at that laboratory.

I am many thankful to many graduate students, under graduate students and staff at Prince of Songkla University who helped me to collect samples during fieldwork and in specimen preparation. I am also grateful to those who gave me useful suggestions when I have seminar or I have some problem in research.

I am very grateful to the Development and Promotion of Science and Technology talents project (DPST) which is a good scholarship that supported me in all expenditure during my master degree program and provided research grants. I also gratefully acknowledge research grants from National Research Council of Thailand in 2012 and Thesis Financial Support fiscal year 2011 from graduate school, Prince of

Songkla University. In addition many thanks to the Institute for the Promotion of Teaching Science and Technology, Ministry of Education and Faculty of Science, Prince of Songkla University for supporting me to give an oral presentation at a major international conference.

Finally, I am very thankful to my father, my mother, my sister, brother all of my family for their encouragement and sustainment of me all the time.

Wanlada Klangnurak

CONTENTS

	Pages
Approval Page	III
Abstract	VI
Acknowledgements	IX
Content	XI
List of Tables	XIV
List of Figures	XV
List of paper and proceeding	XVIII
Chapter 1 Introduction	
1.1 Background and Rationale	1
1.2 Review of Literature	3
1.2.1 Moses perch <i>Lutjanus russelli</i> (Bleeker, 1849)	3
1.2. 2 The geography of Thailand: The area of interest	14
1.3 Objectives	18
1.4 Structure of the thesis	18
Chapter 2 External morphology study	
2.1 Introduction	20
2.2 Objectives	22
2.3 Research question	22
2.4 Materials and Methods	23
2.4.1. Sampling location and collection	23
2.4.2 Measurement of morphometric Characteristics	25
2.4.3. Statistic analysis	27
2.5 Result	28
2.5.1 General characters measurement and new characters measurement	29

CONTENTS (continued)

	Pages
2.6 Preliminary Discussion	35
Chapter 3 Genetic variation between populations of <i>Lutjanus russelli</i> in the Gulf of Thailand and Andaman Sea	
3.1 Introduction	38
3.2 Objective	41
3.3 Research questions	41
3.4 Materials and Methods	42
3.4.1. Study site	42
3.4.2. Sample collection and preservation	43
3.4.3. Sample analyses	44
3.4.4 Data analyses	46
3.5 Results	44
3.5.1. Microsatellite marker efficiency	50
3.5.2. Genetic diversity and tests of Hardy–Weinberg (HW) equilibrium	52
3.5.3. Population genetic differentiation	55
3.6 Preliminary Discussion	60
3.6.1 Genetic diversity and Hardy–Weinberg Equilibrium	60
3.6.2 Population genetic structure and Biogeographic Implications	62
Chapter 4 General Discussion	65
Suggestion	69
References	70
Appendix	79

CONTENTS (continued)

	Pages
Paper 1: Population structure and genetic connectivity of <i>Lutjanus russelli</i> (Lutjanidae) in Thailand.	105
Vitae	109

LIST OF TABLES

	Pages
Table 1 Summary of studies on population structure of lutjanid fishes	13
Table 2 Key characters differentiation of adult fish between Gulf of Thailand and Andaman Sea.	29
Table 3 Key characters differentiation of Juvenile fish between Gulf of Thailand and Andaman Sea	32
Table 4 Allele size (bp), missing genotype, and results from Microchecker.	51
Table 5 Descriptive data of genetic diversity of <i>L. russelli</i> population in Thailand	53
Table 6 AMOVA results testing population differentiation between the Gulf of Thailand and the Andaman Sea population groups	56
Table 7 Pairwise F_{ST} (below diagonal) and R_{ST} (upper diagonal) between 3 populations of <i>L. russelli</i> in Thailand	56
Table 8 Pairwise F_{ST} (below diagonal) and R_{ST} (upper diagonal) between 9 populations of <i>L. russelli</i> in Thailand	57
Table 9 Abbreviation of measurable characters in this study	79
Table 10 Pair-wise Correlation between adult's characters	74
Table 11 Pair-wise Correlation between juvenile's characters	82
Table 12 Characters differentiation of Adult fish between Gulf of Thailand and Andaman Sea	84
Table 13 Characters differentiation of juvenile fish between Gulf of Thailand and Andaman Sea	85
Table 14 Microsatellite markers of <i>L. russelli</i> developed by Guo <i>et al.</i> (2007) Table 15 F_{ST} of the original data (below diagonal) and F_{ST} after the ENA correction method (upper diagonal).	90

LIST OF FIGURES

	Pages
Figure 1 Diagnostic characters of Lutjanidae .	4
Figure 2 Identification character of <i>Lutjanus russelli</i>	5
Figure 3 Color variation of <i>Lutjanus russelli</i> between Pacific Ocean and Indian Ocean; left is adult and juvenile from Pacific region and right is adult from Indian Ocean region.	8
Figure 4 Two color pattern of snapper that often misidentification because of similarity morph; upper row is black spot group and lower row is both spot and stripe group.	11
Figure 5 The coastal of Thailand which face western Pacific Ocean on the eastern site and Andaman Sea on the western side.	14
Figure 6 Surface current in Malacca Straits, Andaman Sea and Indian Ocean in February (1a) and August (1b) (Wyrski 1961(cited in Rizal <i>et al.</i> ,2010)). Close up examination of surface current at the layer 30-50 m around Malacca Straights in February (2a) and August (2b).	15
Figure 7 Relative dissimilarity between fish composition between offshore and inshore and present of discriminated assemblages of fish between the southern part (blue line) and the northern part (red line) in each habitat within the Andaman Sea	16
Figure 8 Sites where fish samples for this study were obtained.	24
Figure 9 General morphology measurements in fish.	26
Figure 10 Outline drawing of <i>Lutjanus russelli</i> , 19 inter-distance from 10 Landmarks	22
Figure 11 Configuration compare the present of stripes of both adult (a) and juvenile (b) fishes from Gulf of Thailand (1) and the Andaman Sea (2).	28

LIST OF FIGURES (continued)

	Pages
Figure 12 Scatter plot from the first two principle component and convex hulls by some general characteristic of adult fish between the Andaman Sea (AN) and the Gulf of Thailand samples.	30
Figure 13 Eigenvector reveal effectiveness of each adult's character to distributions of data.	31
Figure 14 Scatter plot by the first two principle component from four general characters and ten new measurable characters of juvenile fish between the Andaman Sea (AN) and the Gulf of Thailand samples.	33
Figure 15 Eigenvector plots reveal contributions of each juvenile's character to the component axes.	34
Figure 16 Study sites; NA is northern Andaman Sea, SA is southern Andaman Sea and GT is Gulf of Thailand. Number was represent type of collected habitat; 1 is offshore coral reef, 2 is inshore coral reef and 3 is seagrass bed or mangrove	42
Figure 17 Visualization of Polyacrylamide gel of Lru43. Example allele band of twenty-one samples and four lanes of Low Molecular Weight DNA Ladder showed range of allele samples from 200 bp to upper 300 base-pair.	46
Figure 18 The probabilities of each sample in a single vertical line were defined to any given population. Red and green were the possibility of samples correctly assigned to Andaman Sea population and GT population, respectively. The numbers corresponds to the predefined population groups; 1is NA population; 2 is SA population and 3 is GT population.	59

LIST OF FIGURES (continued)

	Pages
Figure 19 Mean likelihood $L(K)$ of data over 20 runs for each K obtained from STRUCTURE software reveals the possible number of clusters.	59
Figure 20 An ad hoc statistic ΔK reveals the most likely number of clusters.	60
Figure 21 Scree plot of adult samples	87
Figure 22 Scree plot of juvenile samples	87
Figure 23 Percentage of statistical power of the data set at different sample sizes and F_{ST} .	100
Figure 24 Relative between the mean likelihood $L(K)$ over 9 hypothetical K	101
Figure 25 Relative of the mean difference between successive likelihood values $L'(K)$ over 9 hypothetical K	102
Figure 25 Relative of the mean difference between successive of $L''(K)$ over 9 hypothetical K	102
Figure 26 Relative of ΔK over 9 hypothetical K	103

LIST OF PAPER AND PROCEEDING

- 1) Population structure and genetic connectivity of *Lutjanus russelli* (Lutjanidae) in Thailand.