

REFERENCES

REFERENCES

- [1] Kobata, K. and Watanabe, T. (1998). Novel capsaicinoid-like substances, capsiate and dihydrocapsiate from the fruits of a nonpungent cultivar, CH-19 Sweet of pepper. **J. Agric. Food Chem**, 46, 1695-1697.
- [2] Walpole, C.S.J., Wrigglesworth R., Bevan S., Campbell E.A. and Winter J. (1993). Analogues of capsaicin with agonist activity as novel analgesic agents: structure-activity studies.1.The aromatic, A-region. **J. Med. Chem**, 36, 2362-2372.
- [3] Walpole, C.S.J., Wrigglesworth R., Bevan S., Campbell E.A. and James I.F. (1993). Analogues of capsaicin with agonist activity as novel analgesic agents: structure-activity studies.2.The amide bond, B-region. **J. Med. Chem**, 36, 2373-2380.
- [4] Walpole, C.S.J., Wrigglesworth R., Bevan S., Campbell E.A. and Perkins M.N. (1993). Analogues of capsaicin with agonist activity as novel analgesic agents: structure-activity studies.3.The hydrophobic side-chain, C-region. **J. Med. Chem**, 36, 2381-2389.
- [5] Hosseini, M., Maddalena, D.J. and Spence, I. (1997). Using artificial neural networks to classify the activity of capsaicin and its analogues. **J. Chem. Inf. Comput. Sci**, 37, 1129-1137.
- [6] Klopman, G. and Li, J. (1995). Quantitative structure-agonist activity relationship of capsaicin analogues. **J. Comput. Aided. Mol. Des**, 9, 283-294.
- [7] Virus, R. M., and Gebhart, G. F. (1979). Pharmacologic actions of capsaicin: apparent involvement of substance P and serotonin. **Life Sci**, 25, 1273-1284.
- [8] Wood, J.N., Winter, J., James, I.F., Rang, H.P. and Bevan, S. (1988). Capsaicin-induced ion fluxes in dorsal root ganglion cells in culture. **J. Neurosci**, 8, 3208-3220.
- [9] Winter, J., Bevan, S. and Campbell, E.A. (1995). Capsaicin and pain mechanisms. **Brit. J. Anaesth**, 75, 157-168.
- [10] Fitzgerald, M. (1983). Capsaicin and sensory neurones-A review. **Pain**, 15, 109-130.

- [11] Sancho, R., Lucena, C., Macho, A., Calzado, M.A. and Blanco-Molina, M. (2002). Immunosuppressive activity of capsaicinoids: capsiate derived from sweet peppers inhibits NF- κ B activation and is a potent anti-inflammatory compound in vivo. **Eur. J. Immunol**, 32, 1753–1763.
- [12] Anish, S. and Arpad, S. (2007). The emerging role of TRPV1 in diabetes and obesity. **Pharmacological Sciences**, 29, 29-36.
- [13] Koichiro, O., Shoji, N., Satoshi, M., Susumu, Y. and Tohru, F. (2001). CH-19 sweet, a non-pungent cultivar of red pepper, Increased body temperature and oxygen consumption in humans. **Biosci. Biotechnol. Biochem**, 65, 2033-2036.
- [14] Naohiko, I., Yoshiko, M., Hitoshi, S. and Michio, T. (2007). Enhanced energy expenditure and fat oxidation in humans with high BMI scores by the ingestion of novel and non-pungent capsaicin analogues (Capsinoids). **Biosci. Biotechnol. Biochem**, 71, 380-389.
- [15] Andrea, R. J., Scott, S. S., Andrew, M. H., Richard, A., Aaron, W. S. and Stuart M. P. (2010). Effects of capsinoid ingestion on energy expenditure and lipid oxidation at rest and during exercise. **Nutr. Metab**, 65, 1-10.
- [16] Satoshi, H., Fuminori, K., Yoriko, M., Koichiro, O., Tatsuo, W. and Susumu, Y. (2011). Capsinoids, non-pungent capsaicin analogs, reduce body fat accumulation without weight rebound unlike dietary restriction in mice. **Biosci. Biotechnol. Biochem**, 75, 95-99.
- [17] Jose, E. G., Donna, H. R. and Eric, R. (2010). Effect of capsinoids on energy metabolism in human subjects. **Brit. J. Nutr**, 103, 38–42.
- [18] Sutoh, K., Kobata, K. and Watanabe, T. (2001). Stability of capsinoid in various solvents. **J. Agric. Food. Chem**, 49, 4026-4030.
- [19] Yoko, I., Fumiko, W. and Kentaro, M. (2007). Process of producing a capsinoid-containing food and drink. **United States Patent Application Publication**, Pub. No. US/0264411 A1.
- [20] Premkumar, L.S., (2009). Treatment of Pain with Resiniferatoxin and Related Analogues. **United States Patent Application Publication**, Pub. No. US/0209633 A1.

- [21] Riteshi, K. B., Dhirendra, P. S., Siddhartha, M. S., Jaspreet, K., Rajat, S. and Ravneet, K. B. (2014). Capsaicin induces “Brite” phenotype in differentiating 3T3-L1 preadipocytes. **PLOS ONE**, 9, 1371-1383.
- [22] Kobata, K., Sutoh, K. and Todo, T. (1999). Nordihydrocapsiate, a new capsinoid from the fruits of a nonpungent pepper, *Capsicum annuum*. **J. Nat. Prod.**, 62, 335-336.
- [23] **Ajinomoto to Market New Dietary Supplement in the U.S.A. CAPSIATE NATURA™.** (2007, July 12). Retrieved July 12, 2007, from http://www.ajinomoto.com/en/ir/ir_library/g2007_07_12.html
- [24] Iwai, K. and et al. (2003). Roles as metabolic regulators for the non-nutrients capsaicin and capsiate, supplemented to diet. **Proc. Japan Acad**, 79, 207-212.
- [25] Bernard, B.K., Tsubuku, T., Kayahara, K. M., Hamada, T., Nakamura, Y. and Shirai, A. (2008). Studies of the toxicological potential of capsinoids: X safety assessment and pharmacokinetics of capsinoids in healthy male volunteers after a single oral ingestion of CH-19 sweet extract. **J. Toxicol**, 27, 137-147.
- [26] Hiroaki, K., Kawasaki, S., Yaqin, L., Ryuji, S. and Tetsuya, M. (2011). Genetically modified plant capable of biosynthesizing capsinoid. **United States Patent Application Publication**, US 0166371 A1.
- [27] Sutoh, K., Kobata, K. and Yazawa, S. (2006). Capsinoid is biosynthesized from phenylalanine and valine in a non-pungent pepper, *Capsicum annuum* L. cv. CH-19 Sweet. **Biosci. Biotechnol. Biochem**, 70, 1513-1516.
- [28] Lang, Y.Q., Kisaka, H. and Sugiyama, R. (2009). Functional loss of pAMT results in biosynthesis of capsinoids, capsaicinoid analogs, in *Capsicum annuum* cv. CH-19 Sweet. **Plant J**, 59, 953-961.
- [29] Tanaka, Y., Hosokawa, M. and Miwa, T. (2010). Newly mutated putative aminotransferase in nonpungent pepper (*Capsicum annuum*) results in biosynthesis of capsinoids, capsaicinoid analogues. **J. Agric. Food. Chem**, 58, 1761-1767.
- [30] Yusuke Amino and Yoshinobu Takino. (2011). Substituted benzyl ester derivative and use thereof. **United States Patent Application Publication**, US 7,981,460 B2.

- [31] Yosuke, A., Wataru, K., Takashi, N. and Kazuko, H. (2010). Production method of capsinoid by dehydrating condensation, stabilizing method of capsinoid, and capsinoid composition. **United States Patent Application Publication**, US 0152291 A1.
- [32] Watanabe, T., Ohnuki, K. and Kobata, K., (2011). Studies on the metabolism and toxicology of emerging capsinoids. **Expert Opin. Drug Metab. Toxicol**, 7(5), 533-542.
- [33] Kodama, T., Masuyama, T., Kayahara, T., Tsubuku, S., Ohishi, T. and Wagner, B.M. (2010). Studies of the toxicological potential of capsinoids XIV: A 26-week gavage toxicity study of dihydrocapsiate in rats. **Int. J. Toxicol**, 29(2S), 27S-54S.
- [34] Powell, A.G., Apovian, C.M. and Aronne, L.J. (2011). New drug targets for the treatment of obesity. **Clin. Pharmacol. Ther**, 90, 40–51.
- [35] Meydani, M. and Hasan, S.T. (2010). Dietary polyphenols and obesity. **Nutrients**, 2, 737–751.
- [36] Kee-Hong, K. and Yeonhwa, Park. (2011). Food components with anti-obesity effect. **Annu. Rev. Food. Sci. Technol**, 2, 237-257.
- [37] Williams, D.J., Edward, D., Hamernig, I., Jian, L., James A.P., Johnson S.K., et al. (2013). Vegetables containing phytochemicals with potential anti-obesity properties a review. **Food. Res. Int**, 52, 323–333.
- [38] Andrea, Z. (2013). Vanilloid receptors-do they have a role in whole body metabolism? Evidence from TRPV1. **J. Diabetes. Complicat**, 27, 287–292.
- [39] Fuminori, K., Naohiko, I., Susumu, Y., Teruo, K., Kazuo, I. and Tohru, F. (2006). Effects of CH-19 sweet, a non-pungent cultivar of red pepper, in decreasing the body weight and suppressing body fat accumulation by sympathetic nerve activation in humans. **Biosci. Biotechnol. Biochem**, 65, 2735–2740.
- [40] Eiamsa-sd, P. (2014). **Synthesis of stable capsiate analogues**. Thesis M.S., Naresuan University, Phitsanulok.
- [41] Liu, Y., Peterson, D.A., Kimura, H. and Schubert, D. (1997). Mechanism of cellular 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction. **J. Neurochem**, 69, 581-593.

- [42] Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. **J. Immunol. Meth**, 65, 55-63.

BIOGRAPHY

BIOGRAPHY

Name - Surname	Arpon Pongkasetkam
Date of Birth	August 11, 1984
Address	25 Moo 5, Salokbat, Amphor Khanuworaluksaburi, Kamphaengphet, Thailand, 62140
Education Background	
2006	B.S. (Chemistry) Naresuan University, Phitsanulok, Thailand