LIST OF TABLES

TABLES		PAGE
2.1	Sign and symptom of musculoskeletal pain from musculoskeletal	
	system disease and degenerative change	10
2.2	Thai mental measurement tools	33
3.1	The character of subjects in no pain (NP), acute musculoskeletal pain	
	(AMSP) and chronic musculoskeletal pain (CMSP) groups	50
3.2	The comparison of the mean of the pain intensity score (PI score),	
	disability point (DP) and pain dysfunction score (PD score)	
	between acute musculoskeletal pain (AMSP) and chronic	
	musculoskeletal pain (CMSP) groups	51
3.3	The number of samples by center, age and gender for the total 3,000	
	samples	54
4.1	The percentage of depression screening by risk factors in 194	
	chronic musculoskeletal pain subjects	64
4.2	Coefficient of selected variables in multiple logistic regression of	
	probable depression	67
4.3	Predicted Logistic regression of probable depression and risk factors	
	in 194 CMSP	68
4.4	The percentage of the no pain (NP) and chronic musculoskeletal	
	pain (CMSP) subjects by demographic factors at the beginning of	
	phase 2: follow-up phase	69
4.5	Percent increase of high risk depression (D score ≥ of 20) in no pain	74
	and CMSP subjects during six months follow-up	71
4.6	The number of occurrences of high risk depression (D score of \geq 20)	74
4 7	in the no pain and CMSP groups in the follow up period	71
4.7	General characteristics of 34 CMSP with high risk depression	70
	(D score of \geq 20).	73

4.8	The percentage of the group A and group B subjects by demographic	
	factors at the beginning of period 2	77
4.9	Pain score and D score in group A and B	78
4.10	The change of pain score, D score, and the events entire the treatment	
	periods (from interview) in group A	79
4.11	The change of pain score, D score, and the events entire the treatment	
	periods (from interview) in group B	80