

บรรณานุกรม

บรรณานุกรม

- [1] World Heart Federation. (2008). Awareness world heart day.
Retrieved March 10, 2013, from <http://www.world-heart-federation.org/what-wedo/awareness/>
- [2] สำนักงานนโยบายและยุทธศาสตร์ กระทรวงสาธารณสุข. (2551). ข้อมูลสถิติ.
สืบค้นเมื่อ 10 มีนาคม 2556, จาก <http://bps.ops.moph.go.th/index.php?mod=bps&doc=54>.
- [3] World Heart Organization. Media centre. (update 2008).
Retrieved March 10, 2013, from http://www.who.int/mediacentre/events/annual/world_heart_day/en/
- [4] สำนักโรคไม่ติดต่อกระทรวงสาธารณสุข. (2555). รายงานประจำปี 2554. กรุงเทพฯ: สำนักงานกิจการโรงพยาบาลองค์การสงเคราะห์ทหารผ่านศึกในพระบรมราชูปถัมภ์.
- [5] Eckel, R.H., Alberti, K.G., Grundy, S.M. and Zimmet, P.Z. (2010). The metabolic syndrome. *Lancet*, 375(9710), 181-183.
- [6] Laaksonen, D.E., Niskanen, L., Lakka, H.M., Lakka, T.A. and Uusitupa, M. (2004). Epidemiology and treatment of the metabolic syndrome. *Annals of Internal Medicine*, 36(5), 332-346.
- [7] Laaksonen, D.E., Lakka, H.M., Niskanen, L.K., Kaplan, G.A., Salonen, J.T. and Lakka, T.A. (2002). Metabolic syndrome and development of diabetes mellitus: application and validation of recently suggested definitions of the metabolic syndrome in a prospective cohort study. *American Journal of Epidemiology*, 156(11), 1070-1077.
- [8] Facchini, F.S., Hua, N., Abbasi, F. and Reaven, G.M. (2001). Insulin resistance as a predictor of age-related diseases. *The Journal of Clinical Endocrinology & Metabolism*, 86(8), 3574-3758.
- [9] Thomas, J.D. and Popovic, Z.B. (2006). Assessment of left ventricular function by cardiacultrasound. *Journal of the American College of Cardiology*, 48(10), 2012-2025.

- [10] Jae, K., James, B., Seward, A. and Tajik, J. (2006). **The Echo Manual**. Rochester, USA: Lippincott Williams & Wilkins.
- [11] Mondillo, S., Galderisi, M., Mele D., Cameli M., Lomoriello VS., Zaca V, and Ballo P. (2011). Speckle-Tracking Echocardiography A New Technique for Assessing of myocardial function. **Journal of Ultrasound in Medicine**, 30(1), 71-83.
- [12] Saito, K., Okura, H., Watanabe, N., Watanabe, N., Hayashida, A., Obase, k., et al. (2009). Comprehensive evaluation of left ventricular strain using speckle echocardiography in normal adults: comparison of three-dimensional and two-dimensional approaches. **Journal of the American Society of Echocardiography**, 22(9), 1025-1030.
- [13] Biswas M., Sudhakar S., Nanda N.C., Buckberg G., Pradhan M., Roomi AU, et al. (2013). Two- and three-Dimensional speckle tracking echocardiography: clinical applications and future directions. **Echocardiography**, 30(1), 88-105.
- [14] Dalen, H., Thorstensen, A., Aase, S.A., Ingul, CB., Torp, H., Vatten, LJ., et al. (2010). Segmental and global longitudinal strain and Strain rate based on echocardiography of 1266 healthy individuals: the HUNT study in Norway. **European Journal of Echocardiography**, 11(2), 176-183.
- [15] Grundy, S.M., Brewer, H.B.Jr., Cleeman, J.I., Smith, SC Jr. and Lenfant, C. (2004). Definition of metabolic syndrome: report of the National Heart, Lung and Blood Institute/American Heart Association Conference on scientific issues related to definition. **Circulation**, 109(3), 433-438.
- [16] Ervin, R.B. (2009). Prevalence of metabolic syndrome among adults 20 years of age and over, by sex, age, race and ethnicity, and body mass index: United States, 2003-2006. **National Health Statistics Reports**, 13(1), 1-8.
- [17] Lakka, H.M., Laaksonen, D.E., Lakka, T.A., Niskanen, L.A., Kumpusalo, E., Tuomilehto, J., et al. (2002). The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. **Journal of the American Medical Association**, 288(21), 2709-2716.

- [18] McNeill, A.M., Rosamond, W.D. Girman, C.J., Golden, S.H., Schmidt, M.I., East, H.E., et al. (2005). The metabolic syndrome and 11-year risk of incident cardiovascular disease in the atherosclerosis risk in communities study. *Diabetes Care*, 28(2), 385-390.
- [19] World Health Organization. (1999). **Definition, diagnosis and classification of diabetes mellitus and its complications: report of a WHO Consultation. Part1: diagnosis and classification of diabetes mellitus.** Geneva: World Health Organization.
- [20] Balkau, B. and Charles, M.A. (1999). Comment of the provisional report from the WHO consultation: European Group for the Study of Insulin Resistance (EGIR). *Diabetes Medicine*, 16(5), 442-443.
- [21] National Cholesterol Education Program Expert Panel. (2001). Executive summary of the third report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III): Final report. *The Journal of the American Medical Association*, 285(19), 2486-2497.
- [22] Einhorn, D., Reaven, G.M., Cobin, R.H., Ford, E., Ganda, O.P., Handelsman. Y. (2002). American College of Endocrinology position statement of the insulin resistance syndrome. *Endocrine Practice*, 9(3), 237-252.
- [23] Alberti, K.G., Zimmet, P. and Shaw J. (2006). The metabolic syndrome-a new worldwide definition. *Lancet*, 366(9491), 1059-1062.
- [24] Grundy, S.M., Cleeman, J.I., Daniels, S.R., Donato, K.A., Krauss, R.M., Savage, P.J., et al. (2005). Diagnosis and management of the metabolic syndrome. An American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. Executive Summary. *Cardiology in Review*, 13(6), 322-327.
- [25] Ko, G.T., Chan, J.C., Cockram, C.S. and Woo, J. (1999). Prediction of hypertension, diabetes, dyslipidemia or albuminuria using simple anthropometric indexes in Hong Kong Chinese. *International Journal of Obesity*, 23(11), 1136-1142.

- [26] Deurenberg-Yap, M., Yian, T.B., Kai, C.S., Deurenberg, P. and Van Staveren, W.A. (1999). Manifestation of cardiovascular risk factors at low levels of body mass index and waist-to-hip ratio in Singaporean Chinese. **Asia Pacific Journal of Clinical Nutrition**, 8(3), 177-183.
- [27] Snehalatha, C., Viswanathan, V. and Ramachandran, A. (2004). Cutoff values for normal anthropometric variables in Asian Indian adults. **Diabetes Care**, 26(5), 1380-1384.
- [28] Choo, V. (2002). WHO reassesses appropriate body-mass index for Asian populations. **Lancet**, 360(9328), 235.
- [29] Mckeigue, P.M., Shah, B. and Marmott, M.G. (1991). Relationship of central Obesity and insulin resistance with high diabetes prevalence and cardiovascular risk in South Asians. **Lancet**, 337(8738), 382-386.
- [30] DeFronzo, R.A. and Ferrannini, E. (1991). Insulin resistance: a multifaceted Syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia and atherosclerotic cardiovascular disease. **Diabetes Care**, 14(3), 173-194.
- [31] Tan, C.E, Ma, S., Wai, D., Chew, S.K. and Tai, E.S. (2004). Can we apply the National Cholesterol Education Program Adult Treatment Panel definition of the metabolic syndrome to Asians?. **Diabetes Care**, 27(5), 1728-1734.
- [32] Alberti, G., Zimmet, P. and Shaw, J., (2006). The IDF consensus worldwide definition of the metabolic syndrome. **International Diabetes Federation**, 23(5), 469-480.
- [33] กลุ่มงานอายุรกรรม โรงพยาบาลราชวิถี (2006). **Metabolic Syndrome**. Retrieved March 15, 2013, from http://hpe4.anamai.moph.go.th/hpe/data/ms/ms_chanchai.htm
- [34] Ford, E.S. (2003). The metabolic syndrome and C-reactive protein, fibrinogen, and leukocyte count: findings from the third national health and nutrition examination survey. **Atherosclerosis**, 168(2), 351-358.

- [35] Abbasi, F., Brown, B.W.Jr., Lamendola, C., McLaughlin, T. and Reaven G.M. (2002). Relationship between obesity, insulin resistance and coronary heart diseases risk. **Journal of the American College of Cardiology**, 40(5), 937-943.
- [36] Alberti, K.G. (1998). Impaired glucosetolerance: what are the clinical implications?. **Diabetes Research and Clinical Practice**, 40(1), s3-s8
- [37] ExecutiveSummary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection. (2001). Evaluation, And Treatment of High Blood CholesterolIn Adults (Adult Treatment Panel III). **The Journal of the American Medical Association**, 285(19), 2486-2497.
- [38] Nabipour, I., Amiri, M., Imami, S.R., Jahfari, S.M., Shafeiaie, E., Nosrati, A., et al. (2007). The metabolic syndrome and nonfatalischemic heart disease; a population-based study. **International Journal of Cardiology**, 118(1), 48-53.
- [39] Ingelsson, E., Pencina, M.J., Tofler, G.H., Benjamin, E.J., Lanier, K.J., Jacques, P.F., et al. (2007). Multimarker approach to evaluate theincidence of the metabolic syndrome and longitudinal changes in metabolic riskfactors: the Framingham Offspring Study. **Circulation**, 116(9), 984-992.
- [40] Jeppesen, J., Hansen, T.W., Rasmussen, S., Ibsen, H., Torp-Pedersen, C. and Madsbad, S. (2007). Insulin resistance, the metabolic syndrome,and risk of incident cardiovascular disease: a population-based study. **Journal of the American College of Cardiology**, 2007, 49(21), 2112-2119.
- [41] Lakka, H.M., Laaksonen, D.E., Lakka, T.A., Niskanen, L.K., Kumpusalo, E. and Tuomilehto, J. (2002). The metabolic syndrome and totaland cardiovascular disease mortality in middle-aged men. **The Journal of the American Medical Association**, 288(21), 2709-2716.
- [42] Darvall, K.A., Sam, R.C., Silverman, S.H., Bradbury, A.W. and Adam, D.J. (2007). Obesity and thrombosis. **European Journal of Vascular and Endovascular Surgery**, 33(2), 223-233.

- [43] Ritchie, S.A. and Connell, J.M. (2007). The link between abdominal obesity, Metabolic syndrome and cardiovascular disease. **Nutrition, Metabolism & Cardiovascular Diseases**, 17(4), 319-326.
- [44] Diamant, M., Lamb, H.J., Van de Ree, M.A., Endert, E.L., Groeneveld, Y. and Bots, M.L. (2005). The association between abdominal visceral fat and carotid stiffness is mediated by circulating inflammatory markers in uncomplicated type 2 diabetes. **The Journal of Clinical Endocrinology and Metabolism**, 90(3), 1495-1501.
- [45] Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults-the evidence report. (1998). **National Institutes of Health**. 2(6), 51S-209S.
- [46] Lindstrom, J., Louheranta, A. and Mannelin, M. Rastas M, Salminen V, Eriksson J, Uusitupa M, Tuomilehto J. (2003). The Finnish Diabetes Prevention Study (DPS): lifestyle intervention and 3-year results on diet and physical activity. **Diabetes Care**, 26(12), 3230-3236.
- [47] Thompson, P.D., Buchner, D., Pina, I.L., Balady, G.J., Williams M.A., Marcus B.H., et al. (2003). Exercise and physical activity in the prevention and treatment of atherosclerotic cardiovascular disease: a statement from the American Heart Association council on clinical cardiology and the council on nutrition, physical activity, and metabolism. **Circulation**, 107(24), 3109-3116.
- [48] Grundy, S.M., Hansen, B., Smith, S.C., Cleeman J.I. and Kahn R.A. (2004). Clinical management of metabolic syndrome: Report of the National Heart, Lung and Blood Institute/American Heart Association/National Heart, Lung, and Blood Institute/American Diabetes Association conference on scientific issues related to management. **Circulation**, 109(4), 551-556

- [49] National Cholesterol Education Program Expert Panel. (2001). Executive summary of the third report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III): Final report. **The Journal of the American Association**, 285(19), 2486-2497.
- [50] He, J., Whelton, P.K., Appel, U., Charleston, J. and Klag, M.J. (2000). Long-term effects of weightloss and dietary sodium reduction on incidence of hypertension. **Hypertension**, 35(2), 544-549.
- [51] The ALLHAT Officers and Coordinators for the ALLHAT Collaborative Reserch Group. (2002). Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium channel blocker vs diuretic. **The Journal of the American Association**, 288(23), 2981-2987.
- [52] The Heart Outcomes Prevention Evaluation Study Investigators. (2000). Effects of an angiotensin-converting enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. **The New England Journal of Medicine**, 342(3), 145-153.
- [53] Dahlof, B., Devereux, R.B. and Kjeldsen, S.E. (2002). Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomized trial against atenolol. **Lancet**, 362(9395), 1527-35.
- [54] Julius, S., Kjeldsen, S.E., Weber, M., Brunner, H.R., Ekman, S., Hansson, L., et al. (2004). Outcomes in hypertensive patients at high cardiovascular risk treated with regimens based on valsartan or amlodipine: the VALUE randomized trial. **Lancet**, 363(9426), 2022-2031.
- [55] Hansson, L., Lindholm, L.H., Ekblom, T., Dahlof, B., Lanke, J., Schersten, B., et al. (1999). Randomised trial of old and new antihypertensive drugs in elderly patients: cardiovascular mortality and morbidity the Swedish Trial in Old Patients with hypertension-2 study. **Lancet**, 354(9192), 1751-1756.

- [56] The Diabetes Prevention Program Research Group. (2002). Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. **The New England Journal of Medicine**, 346(6), 393-403.
- [57] Chiasson, J.L., Josse, R.G., Gomis, R., Hanefeld, M., Karasik, A. and Laakso, M. (2002). Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomized trial. **Lancet**, 359(9323), 2072-2077.
- [58] Gaede, P., Vedel, P., Larsen, N., Jensen, G.V., Parving, H.H. and Pedersen, O. (2003). Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. **The New England Journal of Medicine**, 348(5), 383-393.
- [59] Douglas, P.S., Garcia, M.J., Lai, W.W., Manning, W.J., Patel, A.R., Picard, M.H., et al. (2011). Appropriate Use Criteria for Echocardiography. **Journal of the American College of Cardiology**, 57(9), 1126–1166.
- [60] Lang, R. M., Bierig, M., Devereux, R.B., Lang, R.M., Bierig, M., Devereux, R.B., et al. (2005). Recommendations for chamber quantification: quantitative assessment of left ventricular volume and ejection fraction using two-dimensional speckle tracking echocardiography **Cardiovascular Imaging. European Society of Cardiology**, 7(11), 79-108.
- [61] Pislaru, C., Abraham, T.P. and Belohlavek, M. (2002). Strain and strain rate echocardiography. **Current Opinion in Cardiology**, 17(5), 443-454.
- [62] Stoylen A. (November, 2013) **Myocardial deformation imaging by ultrasound/ echocardiography: Tissue Doppler and Speckle tracking**. Retrieved October 5, 2013, from <http://folk.ntnu.no/stoylen/strainrate/>
- [63] Applegarth, D.A., Besley, G.T.N. and Clarke, L.A. (1995). Biochemical and Molecular Genetics. In "G.B. Reed, A.E. Claireaux, F. Cockburn (Eds.). **Diseases of the fetus and Newborn: Pathology, imaging, genetics and Management** (2nd ed. pp. 1131-1163). London: Chapman & Hall.
- [64] Disorders of lipoprotein metabolism. (2008). In E. Braunwald, A.S. Fauci, D.L. Kasper, S.L. Hauser, D.L. Longo, J.L. Jameson (Eds.). **Harrison's Principles of Internal Medicine** (17th Ed., pp. 2286-2303). New York: McGrawHill.

- [65] Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults. (2001). Executive summary of the third report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel III). *Journal of the American Medical Association*, 28(19), 2486-2497.
- [66] Feigenbaum, H., Armstrong, W.F. and Thomas, R. (2004). **Feigenbaum's Echocardiography**. Lippincott Williams & Wilkins.
- [67] Gorcsan, J. and Tanaka, H. (2011). Echocardiographic assessment of myocardial strain. *Journal of the American College of Cardiology*, 58(14), 1401-1413.
- [68] Marcus, K., Mavinkurve-Groothuis, A.M., Barends, M., Dijk, A.V., Feuth, T., Korte, K.D., et al. (2011). Reference values for myocardial two-dimensional strain echocardiography in a healthy pediatric and young adult cohort. *Journal of the American Society of Echocardiography*, 24(6), 625-636.
- [69] Marwick, T.H. (2006). Measurement of strain and strain rate by echocardiography: ready for prime time?. *Journal of the American College of Cardiology*, 47(7), 1313-1327.
- [70] Mesas, A.E., León-Muñoz, L.M., Rodríguez-Artalejo, F. and Lopez-Garcia, E. (2011). The effect of coffee on blood pressure and cardiovascular disease in hypertensive individuals: a systematic review and meta-analysis. *The American Journal of Clinical Nutrition*, 94(4), 1113-1126.
- [71] Nesser, H.J. and Winter, S. (2009). Speckle tracking in the evaluation of left ventricular dyssynchrony. *Echocardiography*, 26(3), 324-336.
- [72] Oh, J.K., Seward, J.B. and Tajik A.J. (2007). **The Echo manual** (2nd ed.). Philadelphia: Lippincott Williams & Wilkins.

- [73] Lang, R.M., Bierig, M., Devereux, R.B., Flachskampf, F.A., Foster, E., Pellikka, P.A., et al. (2003). Recommendations for chamber Quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, Developed in Conjunction with the European Association of Echocardiography, a Branch of the European Society of Cardiology. *Journal of the American Society of Echocardiogram*, 18(12), 1440-1463.
- [74] Sen Gupta, P.P., Korinek, J., Belohlavek, M., Narula, J., Vannan, M.A., Jahangir, A., et al. (2006). Left ventricular structure and function: basic science for cardiac imaging. *Journal of the American College of Cardiology*, 48(10), 1988-2001.
- [75] Takeuchi, M., Nishikage, T., Nakai, H., Kokumai, M., Otani, S. and Lang, R.M. (2007). The Assess of Left Ventricular Twist in Anterior Wall Myocardial Infarction Using Two-dimensional Speckle Tracking Imaging. *Journal of the American College of Cardiology*, 20(1), 36-45.
- [76] Vasan, R.S., Beiser, A., Seshadri, S., Larson, M.G., Kannel, W.B., D'Agostino, R.B., et al. (2002). Residual lifetime risk for developing hypertension in middle-aged women and men: The Framingham Heart Study. *Journal of the American Medical Association*, 287(8), 1003-1010.
- [77] Riediger, N.D. and Clara, L. (2011). Prevalence of metabolic syndrome in the Canadian adult population. *Canadian Medical Association Journal*, 183(15), 1127-1134.
- [78] Bassin, S.L., Luo, Y., Li, A., Perez, A. and Wong, N. (2013). Prevalence of and trends in metabolic syndrome and associated cardiovascular risk factor among US adolescents between 1999 and 2008. *Cardiovascular Endocrinology*, 2(2), 23-30.
- [79] Ervin, R.B. (2009). Prevalence of metabolic syndrome among adults 20 years of age and over, by sex, age, race and ethnicity, and body mass index: United States, 2003-2006. *National Health Statistics Reports*, 5(13), 1-8.

- [80] Lakka, H.M., Laaksonen, D.E., Lakka, T.A., Niskanen, L.K., Kumpusalo, E., Tuomilehto, J., et al. (2002). The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. **Journal of the American Medical Association**, 288(21), 2709-2716.
- [81] McNeill, A.M., Rosamond, W.D., Girman, C.J., Golden, S.H., Schmidt, M.I., East, H.E., et al. (2005). The metabolic syndrome and 11-year risk of incident cardiovascular disease in the atherosclerosis risk in communities study. **Diabetes Care**, 28(2), 385-390.
- [82] Yu, C.M., Sanderson, J.E., Marwick, T.H. and Oh, J.K. (2007). Tissue Doppler imaging a new prognosticator for cardiovascular diseases. **Journal of the American College of Cardiology**, 49(19), 1903-1914.
- [83] Winter, R., Jussila R., Nowak, J. and Brodin, L.A. (2007). Speckle tracking echocardiography is a sensitive tool for detection of myocardial ischemia: a pilot study from the catheterization laboratory during percutaneous coronary intervention. **Journal of the American Society of Echocardiography**, 20(8), 974-981.
- [84] Artis, N.J., Oxborough, D.L., Williams, G., Pepper, C.B. and Tan, L.B. (2008). Two-dimensional strain imaging: a new echocardiographic advance with research and clinical applications. **International Journal of Cardiology**, 123(3), 140-148.
- [85] Marwick, T.H. (2006). Measurement of strain and strain rate by echocardiography: ready for prime time?. **Journal of the American College of Cardiology**, 47(7), 1313-1327.
- [86] Leitman, M., Lysyansky, P., Sidenko, S., Shir, V., Peleg, E., Binenbaum, M., et al. (2004). Two-dimensional strain – a novel software for real-time quantitative echocardiographic assessment of myocardial function. **Journal of the American Society of Echocardiography**, 17(10), 1021-1029.

- [87] Marwick, T.H., Leano, R.L., Brown, J., Sun, J.P., Hoffmann, R., Lysyansky, P., et al. (2009). Myocardial strain measurement with 2-dimensional speckle-tracking echocardiography. *Journal of the American College of Cardiology*, 2(1), 80-84.
- [88] Cuspidi, C., Vaccarella, A., Negri, F. and Sala, C. (2010). Resistant hypertension and left ventricular hypertrophy: an overview. *Journal of the American Society of Hypertension*, 4(6), 319-314.
- [89] Kouzu, H., Yuda, S. Muranaka, A., Doi, T., Yamamoto, H., Shimoshige, S., et al. (2011). Left Ventricular Hypertrophy Causes Different Changes in Longitudinal, Radial, and Circumferential Mechanics in Patients with Hypertension: A Two-Dimensional Speckle Tracking Study. *Journal of the American Society of Echocardiography*, 24(2), 192-199.
- [90] Tamaki S., Sakata Y., Mano T., Ohtani, T., Takeda, Y., Kamimura, D., et al. (2010). Long-term -blocker therapy improves diastolic function even without the therapeutic effect on systolic function in patients with reduced ejection fraction. *Journal of Cardiology*, 56(2), 176-182.
- [91] El-refai M., Edward L., Wells K., Swadia, T., Sabbah, H.N., Spertus, J.A., et al. (2013). Comparison of Beta-Blocker Effectiveness in Heart Failure Patients with Preserved Ejection Fraction Versus Those With Reduced Ejection Fraction. *Journal of Cardiac Failure*, 19(2), 73-79.
- [92] Gong, H.P, Tan, H.W., Fang, N.N., Song, T., Li, S.H, Zhong M, et al. (2008).. Impaired left ventricular systolic and diastolic function in patients with metabolic syndrome as assessed by strain and strain rate. *Diabetes Research and Clinical Practice*. 83(3), 300-307.
- [93] Katz, D.H., Beussink, L., Sauer, A.J., Freed, B.H., Burke, M.A. and Shah, S.J. (2013). Prevalence, Clinical Characteristics, and Outcomes Associated with Eccentric Versus Concentric Left Ventricular Hypertrophy in Heart Failure With Preserved Ejection Fraction. *Journal of the American College of Cardiology*, 112(8), 1158-1164.

- [94] Nishikage, T., Nakal, H., Mor-Avi, V., Lang, R.M., Salgo, I.S., Settlemier, S.H., et al. (2009). Quantitative assessment of left ventricular volume and ejection fraction using two-dimensional speckle tracking echocardiography. **European Journal of Echocardiography**. 10(1), 82-88.
- [95] Saaïd, I., Fehri, W., Ben Mansour, N., Amor, B., Dahmani, R., Haggui, A., et al. (2011). Left ejection fraction evaluation by speckle tracking. **Archives of Cardiovascular Diseases**, 104(4), 271-271.
- [96] Di Salvo, G., Di Bello, V. and Salustri, A. (2011). Early left ventricular longitudinal systolic dysfunction and cardiovascular risk factors in 1,371 asymptomatic subjects with normal ejection fraction: a tissue Doppler study. **Echocardiography (Mount Kisco, NY)**, 28(3), 268-75.
- [97] Clinton, W. and Edward, W. (2005). **Age-Related Changes in the Cardiovascular System Hypertension in elderly Humana Press Inc.** United States: Totowa, N.J.
- [98] Govindaraju, D.R., Pecina, K.M., Raj, D.S., Massaro, J.M., Carnes, B.A. and D'Agostino, R.B. (2014). A systems analysis of age-related changes in some cardioaging traits. **Biogerontology**, 15(2), 139-152.

อภิธานศัพท์

อภิธานศัพท์

Speckle tracking echocardiogram	=	เทคนิคการวิเคราะห์การเคลื่อนที่ของเนื้อเยื่อในหัวใจที่ได้จากการภาพคลื่นเสียงสะท้อนความถี่สูง
Strain	=	การเคลื่อนที่ของกล้ามเนื้อหัวใจ
Longitudinal strain	=	การเคลื่อนที่ของกล้ามเนื้อหัวใจตามแนวยาว
Radial strain	=	การเคลื่อนที่ของกล้ามเนื้อหัวใจตามแนวเส้นผ่านศูนย์กลาง
Circumference strain	=	การเคลื่อนที่ของกล้ามเนื้อหัวใจตามแนวเส้นรอบวง
Metabolic syndrome	=	ภาวะอ้วนลงพุง, กลุ่มอาการที่มีกระบวนการเผาผลาญอาหารที่ผิดปกติ
Ejection fraction	=	ค่าการประเมินทำงานของหัวใจ

ประวัติผู้วิจัย

ประวัติผู้วิจัย

ชื่อ-ชื่อสกุล	สมฤทัย วัฒนชัย
วัน เดือน ปีเกิด	12 พฤษภาคม 2531
ที่อยู่ปัจจุบัน	9/1 หมู่ 5 ตำบลท่าหลวง อำเภอเมือง จังหวัดพิจิตร 66000
ที่ทำงานปัจจุบัน	โรงพยาบาลรามาริบัติ คณะแพทยศาสตร์ มหาวิทยาลัยมหิดล เขตราชเทวี กรุงเทพฯ 10400
ประวัติการศึกษา	พ.ศ. 2552 วท.บ. (เทคโนโลยีหัวใจและทรวงอก) มหาวิทยาลัยนเรศวร

ผลงานการตีพิมพ์

สมฤทัย วัฒนชัย. (2557). การเปรียบเทียบการเคลื่อนที่ของกล้ามเนื้อหัวใจจากภาพคลื่นเสียงสะท้อนความถี่สูง 2 มิติด้วยวิธีสเปกเทรกกิง ในกลุ่มผู้สูงอายุที่มีภาวะอ้วนลงพุงกับกลุ่มคนสุขภาพดี. ใน นเรศวรวิจัย ครั้งที่ 10 เครือข่ายวิจัย สร้างความรู้สู่อาเซียน (หน้า 1,271-1,280). พิษณุโลก: มหาวิทยาลัยนเรศวร.