

REFERENCES

REFERENCES

- Abate, N. and Garg, A. (1995). Heterogeneity in adipose tissue metabolism: causes, implications and management of regional adiposity. **Prog Lipid Res**, 34, 53-70.
- Aekplakorn, W., Chongsuvivatwong, V., Tatsanavivat, P. and Suriyawongpaisal, P. (2011). Prevalence of metabolic syndrome defined by the International Diabetes Federation and National Cholesterol Education Program criteria among Thai adults. **Asia Pac J Public Health**, 23, 792-800.
- Akase, T., Shimada, T., Terabayashi, S., Ikeya, Y., Sanada, H. and Aburada, M. (2011). Antiobesity effects of *Kaempferia parviflora* in spontaneously obese type II diabetic mice. **J Nat Med**, 65, 73-80.
- Alberti, K.G., Zimmet, P. and Shaw, J. (2005). The metabolic syndrome--a new worldwide definition. **Lancet**, 366, 1059-1062.
- Alberti, K.G., Zimmer, P., Shaw, J. and Grundy, S.M. (2006). The IDF consensus worldwide definition of the metabolic syndrome. **Brussels: International Diabetes Federation**, 16, 1-23.
- Alberti, K.G. and Zimmet, P.Z. (1998). Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. **Diabet Med**, 15, 539-553.
- Allison, D.B. and Heshka, S. (1991). Toward an empirically derived typology of obese persons. **Int J Obes**, 15, 741-754.
- Angela, M., Michael, A., Matthew, R. and Edward, A. (2007). Diet-induced metabolic syndrome in rodent models. **animal lab new**, 12, 1-7.
- Anjana, M., Sandeep, S., Deepa, R., Vimalaswaran, K.S., Farooq, S. and Mohan, V. (2004). Visceral and central abdominal fat and anthropometry in relation to diabetes in Asian Indians. **Diabetes Care**, 27, 2948-2953.
- Arai, H., Yamamoto, A., Matsuzawa, Y., Saito, Y., Yamada, N., Oikawa, S., et al. (2006). Prevalence of metabolic syndrome in the general Japanese population in 2000. **J Atheroscler Thromb**, 13, 202-208.

- Balkau, B. and Charles, M.A. (1999). Comment on the provisional report from the WHO consultation. European Group for the Study of Insulin Resistance (EGIR). **Diabet Med**, 16, 442-443.
- Ball, S.D., Keller, K.R., Moyer-Mileur, L.J., Ding, Y.W., Donaldson, D. and Jackson, W.D. (2003). Prolongation of satiety after low versus moderately high glycemic index meals in obese adolescents. **Pediatrics**, 111, 488-494.
- Bartoli, E., Fra, G.P. and Carnevale Schianca, G.P. (2011). The oral glucose tolerance test (OGTT) revisited. **Eur J Intern Med**, 22, 8-12.
- Ble-Castillo, J.L., Aparicio-Trapala, M.A., Juarez-Rojop, I.E., Torres-Lopez, J.E., Mendez, J.D., Aguilar-Mariscal, H., et al. (2012). Differential effects of high-carbohydrate and high-fat diet composition on metabolic control and insulin resistance in normal rats. **Int J Environ Res Public Health**, 9, 1663-1676.
- Buettner, R., Parhofer, K.G., Woenckhaus, M., Wrede, C.E., Kunz-Schughart, L.A., Scholmerich, J., et al. (2006). Defining high-fat-diet rat models: metabolic and molecular effects of different fat types. **J Mol Endocrinol**, 36, 485-501.
- Cameron, A.J., Magliano, D.J., Zimmet, P.Z., Welborn, T. and Shaw, J.E. (2007). The metabolic syndrome in Australia: prevalence using four definitions. **Diabetes Res Clin Pract**, 77, 471-478.
- Castro, J.P., El-Atat, F.A., McFarlane, S.I., Aneja, A. and Sowers, J.R. (2003). Cardiometabolic syndrome: pathophysiology and treatment. **Curr Hypertens Rep**, 5, 393-401.
- Cawston, E.E. and Miller, L.J. (2010). Therapeutic potential for novel drugs targeting the type 1 cholecystokinin receptor. **Br J Pharmacol**, 159, 1009-1021.
- Chien, T.Y., Chen, L.G., Lee, C.J., Lee, F.Y. and Wang, C.C. (2008). Anti-inflammatory constituents of *Zingiber zerumbet*. **Food Chemistry**, 110, 584-589.
- Chivapat, S., Chavalittumrong, P., Attawish, A. and Rungsipipat, A. (2010). Chronic toxicity study of *Kaempferia parviflora* Wall ex. Extract. **Thai J. Vet.Med**, 40(4), 377-383.

- Chomchalow, N., Bansiddh, I.J. and MacBaine, C. (2003). Amazing Thai Medicinal Plants. **Horticultural Research Institute (HRI), Department of Agriculture and Horticultural Science Society of Thailand (HSST)**, Bangkok: HSST, The institute.
- Coelho, M.S., Lopes, K.L., Freitas Rde, A., de Oliveira-Sales, E.B., Bergasmaschi, C.T., Campos, R.R., et al. (2010). High sucrose intake in rats is associated with increased ACE2 and angiotensin-(1-7) levels in the adipose tissue. **Regul Pept**, 162, 61-67.
- Coletta, D.K. and Mandarino, L.J. (2011). Mitochondrial dysfunction and insulin resistance from the outside in: extracellular matrix, the cytoskeleton, and mitochondria. **Am J Physiol Endocrinol Metab**, 301, E749-755.
- Cong, W.N., Tao, R.Y., Tian, J.Y., Liu, G.T. and Ye, F. (2008). The establishment of a novel non-alcoholic steatohepatitis model accompanied with obesity and insulin resistance in mice. **Life Sci**, 82, 983-990.
- Department of Agriculture, Thailand. (2009). **Medicinal properties of *Kaempferia parviflora***. Retrieved December 3, 2013, from <http://it.doa.go.th/vichakan/news.php?newsid=22>
- DeFronzo, R.A. and Ferrannini, E. (1991). Insulin resistance. A multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia, and atherosclerotic cardiovascular disease. **Diabetes Care**, 14, 173-194.
- Dissard, R., Klein, J., Caubet, C., Breuil, B., Siwy, J., Hoffman, J., et al. (2013). Long term metabolic syndrome induced by a high fat high fructose diet leads to minimal renal injury in C57BL/6 mice. **PLoS One**, 8, e76703.
- Doucet, E., Almeras, N., White, M.D., Despres, J.P., Bouchard, C. and Tremblay, A. (1998). Dietary fat composition and human adiposity. **Eur J Clin Nutr**, 52, 2-6.
- Einhorn, D., Reaven, G.M., Cobin, R.H., Ford, E., Ganda, O.P., Handelsman, Y., et al. (2003). American College of Endocrinology position statement on the insulin resistance syndrome. **Endocr Pract**, 9, 237-252.
- Enser, M. (1972). Clearing-factor lipase in obese hyperglycaemic mice (ob-ob). **Biochem J**, 129, 447-453.

- Ervin, R.B. (2009). Prevalence of metabolic syndrome among adults 20 years of age and over, by sex, age, race and ethnicity, and body mass index: United States, 2003-2006. **Natl Health Stat Report**, 1-7.
- Folkow, B. (1982). Physiological aspects of primary hypertension. **Physiol Rev**, 62, 347-504.
- Ford, E.S. (2005a). Prevalence of the metabolic syndrome defined by the International Diabetes Federation among adults in the U.S. **Diabetes Care**, 28, 2745-2749.
- Ford, E.S. (2005b). Risks for all-cause mortality, cardiovascular disease, and diabetes associated with the metabolic syndrome: a summary of the evidence. **Diabetes Care**, 28, 1769-1778.
- Ford, E.S., Giles, W.H. and Dietz, W.H. (2002). Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. **JAMA**, 287, 356-359.
- Fraulob, J.C., Ogg-Diamantino, R., Fernandes-Santos, C., Aguila, M.B. and Mandarim-de-Lacerda, C.A. (2010). A mouse model of metabolic syndrome: insulin resistance, fatty liver and non-alcoholic fatty pancreas disease (NAFPD) in C57BL/6 mice fed a high fat diet. **J Clin Biochem Nutr**, 46, 212-223.
- Fung, T.T., Rimm, E.B., Spiegelman, D., Rifai, N., Tofler, G.H., Willett, W.C., et al. (2001). Association between dietary patterns and plasma biomarkers of obesity and cardiovascular disease risk. **Am J Clin Nutr**, 73, 61-67.
- Gallou-Kabani, C., Vige, A., Gross, M.S., Rabes, J.P., Boileau, C., Larue-Achagiotis, C., et al. (2007). C57BL/6J and A/J mice fed a high-fat diet delineate components of metabolic syndrome. **Obesity (Silver Spring)**, 15, 1996-2005.
- Gordon, D.J., Probstfield, J.L., Garrison, R.J., Neaton, J.D., Castelli, W.P., Knoke, J.D., et al. (1989). High-density lipoprotein cholesterol and cardiovascular disease. Four prospective American studies. **Circulation**, 79, 8-15.
- Grundy, S.M., Brewer, H.B., Jr., Cleeman, J.I., Smith, S.C. and Jr., Lenfant, C. (2004). Definition of metabolic syndrome: report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. **Arterioscler Thromb Vasc Biol**, 24, e13-18.

- Grundy, S.M., Cleeman, J.I., Daniels, S.R., Donato, K.A., Eckel, R.H., Franklin, B.A., et al. (2005). Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. **Circulation**, 112, 2735-2752.
- Halton, T.L., Willett, W.C., Liu, S., Manson, J.E., Stampfer, M.J. and Hu, F.B. (2006). Potato and french fry consumption and risk of type 2 diabetes in women. **Am J Clin Nutr**, 83, 284-290.
- Hirayama, I., Yi, Z., Izumi, S., Arai, I., Suzuki, W., Nagamachi, Y., et al. (1999). Genetic analysis of obese diabetes in the TSOD mouse. **Diabetes**, 48, 1183-1191.
- Huang, W., Metlakunta, A., Dedousis, N., Zhang, P., Sipula, I., Dube, J.J., et al. (2010). Depletion of liver Kupffer cells prevents the development of diet-induced hepatic steatosis and insulin resistance. **Diabetes**, 59, 347-357.
- Jensen, M.D. (2008). Role of body fat distribution and the metabolic complications of obesity. **J Clin Endocrinol Metab**, 93, S57-63.
- Kaplan, N.M. (1989). The deadly quartet. Upper-body obesity, glucose intolerance, hypertriglyceridemia, and hypertension. **Arch Intern Med**, 149, 1514-1520.
- Karmelic, I., Lovric, J., Bozina, T., Ljubic, H., Vogrinc, Z., Bozina, N., et al. (2012). Adiponectin level and gene variability are obesity and metabolic syndrome markers in a young population. **Arch Med Res**, 43, 145-153.
- Kim, M.J. and Kim, H.K. (2009). Perilla leaf extract ameliorates obesity and dyslipidemia induced by high-fat diet. **Phytother Res**, 23, 1685-1690.
- Kopelman, P.G. (2000). Obesity as a medical problem. **Nature**, 404, 635-643.
- Koruk, M., Savas, M.C., Yilmaz, O., Taysi, S., Karakok, M., Gundogdu, C., et al. (2003). Serum lipids, lipoproteins and apolipoproteins levels in patients with nonalcoholic steatohepatitis. **J Clin Gastroenterol**, 37, 177-182.
- Lakka, H.M., Laaksonen, D.E., Lakka, T.A., Niskanen, L.K., Kumpusalo, E., Tuomilehto, J., et al. (2002). The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. **JAMA**, 288, 2709-2716.
- Lam, J.C. and Ip, M.S. (2009). Obstructive sleep apnea and the metabolic syndrome. **Expert Rev Respir Med**, 3, 177-186.

- Leardkamolkarn, V., Tiemyuyen, S. and Sripanidkulchai, B.O. (2009). Pharmacological activity of *Kaempferia parviflora* extract against human bile duct cancer cell lines. **Asian Pac J Cancer Prev**, 10, 695-698.
- Lin, S., Thomas, T.C., Storlien, L.H. and Huang, X.F. (2000). Development of high fat diet-induced obesity and leptin resistance in C57Bl/6J mice. **Int J Obes Relat Metab Disord**, 24, 639-646.
- Lohsoonthorn, V., Lertmaharit, S. and Williams, M.A. (2007). Prevalence of metabolic syndrome among professional and office workers in Bangkok, Thailand. **J Med Assoc Thai**, 90, 1908-1915.
- Lombardo, Y.B., Drago, S., Chicco, A., Fainstein-Day, P., Gutman, R., Gagliardino, J.J., et al. (1996). Long-term administration of a sucrose-rich diet to normal rats: relationship between metabolic and hormonal profiles and morphological changes in the endocrine pancreas. **Metabolism**, 45, 1527-1532.
- Lutsey, P.L., Steffen, L.M. and Stevens, J. (2008). Dietary intake and the development of the metabolic syndrome: the Atherosclerosis Risk in Communities study. **Circulation**, 117, 754-761.
- Maachi, M., Pieroni, L., Bruckert, E., Jardel, C., Fellahi, S., Hainque, B., et al. (2004). Systemic low-grade inflammation is related to both circulating and adipose tissue TNF α , leptin and IL-6 levels in obese women. **Int J Obes Relat Metab Disord**, 28, 993-997.
- Malakul, W., Ingkaninan, K., Sawasdee, P. and Woodman, O.L. (2011a). The ethanolic extract of *Kaempferia parviflora* reduces ischaemic injury in rat isolated hearts. **J Ethnopharmacol**, 137, 184-191.
- Malakul, W., Thirawarapan, S., Ingkaninan, K. and Sawasdee, P. (2011b). Effects of *Kaempferia parviflora* Wall. Ex Baker on endothelial dysfunction in streptozotocin-induced diabetic rats. **J Ethnopharmacol**, 133, 371-377.
- Marques-Lopes, I., Ansorena, D., Astiasaran, I., Forga, L. and Martinez, J.A. (2001). Postprandial de novo lipogenesis and metabolic changes induced by a high-carbohydrate, low-fat meal in lean and overweight men. **Am J Clin Nutr**, 73, 253-261.

- Matthews, D.R., Hosker, J.P., Rudenski, A.S., Naylor, B.A., Treacher, D.F. and Turner, R.C. (1985). Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. **Diabetologia**, 28, 412-419.
- McKeown, N.M., Meigs, J.B., Liu, S., Saltzman, E., Wilson, P.W. and Jacques, P.F. (2004). Carbohydrate nutrition, insulin resistance, and the prevalence of the metabolic syndrome in the Framingham Offspring Cohort. **Diabetes Care**, 27, 538-546.
- Mekjaruskul, C., Jay, M. and Sripanidkulchai, B. (2012). Modulatory effects of Kaempferia parviflora extract on mouse hepatic cytochrome P450 enzymes. **J Ethnopharmacol**, 141, 831-839.
- Melancon, S., Bachelard, H., Badeau, M., Bourgoin, F., Pitre, M., Lariviere, R., et al. (2006). Effects of high-sucrose feeding on insulin resistance and hemodynamic responses to insulin in spontaneously hypertensive rats. **Am J Physiol Heart Circ Physiol**, 290, H2571-2581.
- Messier, C., Awad, N. and Gagnon, M. (2004). The relationships between atherosclerosis, heart disease, type 2 diabetes and dementia. **Neurol Res**, 26, 567-572.
- Ministry of Public Health, Thailand. (2013). **Public Health Statistics: The number and rate of deaths from major diseases**. Retrieved December 1, 2013, from http://bps.ops.moph.go.th/Healthinformation/statistic54/2.3.2_54.pdf
- Misra, A. and Khurana, L. (2009). The metabolic syndrome in South Asians: epidemiology, determinants, and prevention. **Metab Syndr Relat Disord**, 7, 497-514.
- Modan, M., Halkin, H., Almog, S., Lusky, A., Eshkol, A., Shefi, M., et al. (1985). Hyperinsulinemia. A link between hypertension obesity and glucose intolerance. **J Clin Invest**, 75, 809-817.
- Muller-Wieland, D., Kotzka, J., Knebel, B. and Krone, W. (1998). Metabolic syndrome and hypertension: pathophysiology and molecular basis of insulin resistance. **Basic Res Cardiol** 93 Suppl, 2, 131-134.

- Murase, T., Mizuno, T., Omachi, T., Onizawa, K., Komine, Y., Kondo, H., et al. (2001). Dietary diacylglycerol suppresses high fat and high sucrose diet-induced body fat accumulation in C57BL/6J mice. **J Lipid Res**, 42, 372-378.
- Murata, K., Deguchi, T., Fujita, T. and Matsuda, H. (2013). Improvement in blood fluidity by *Kaempferia parviflora* rhizome. **J Nat Med**, 67, 719-724.
- NCEP. (2001). Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). **JAMA**, 285, 2486-2497.
- Nicol, C.J., Adachi, M., Akiyama, T.E. and Gonzalez, F.J. (2005). PPARgamma in endothelial cells influences high fat diet-induced hypertension. **Am J Hypertens**, 18, 549-556.
- Ogden, C.L., Carroll, M.D., Curtin, L.R., McDowell, M.A., Tabak, C.J. and Flegal, K.M. (2006). Prevalence of overweight and obesity in the United States, 1999-2004. **JAMA**, 295, 1549-1555.
- Oka, R., Yagi, K., Sakurai, M., Nakamura, K., Moriuchi, T., Miyamoto, S., et al. (2012). Insulin secretion and insulin sensitivity on the oral glucose tolerance test (OGTT) in middle-aged Japanese. **Endocr J**, 59, 55-64.
- Palanisamy, N., Viswanathan, P., Ravichandran, M.K. and Anuradha, C.V. (2010). Renoprotective and blood pressure-lowering effect of dietary soy protein via protein kinase C beta II inhibition in a rat model of metabolic syndrome. **Can J Physiol Pharmacol**, 88, 28-37.
- Parekh, P.I., Petro, A.E., Tiller, J.M., Feinglos, M.N. and Surwit, R.S. (1998). Reversal of diet-induced obesity and diabetes in C57BL/6J mice. **Metabolism**, 47, 1089-1096.
- Patanasethanont, D., Nagai, J., Matsuura, C., Fukui, K., Sutthanut, K., Sripanidkulchai, B.O., et al. (2007). Modulation of function of multidrug resistance associated-proteins by *Kaempferia parviflora* extracts and their components. **Eur J Pharmacol**, 566, 67-74.
- Patel, P. and Macerollo, A. (2010). Diabetes mellitus: diagnosis and screening. **Am Fam Physician**, 81, 863-870.

- Pietilainen, K.H., Kaprio, J., Borg, P., Plasqui, G., Yki-Jarvinen, H., Kujala, U.M., et al. (2008). Physical inactivity and obesity: a vicious circle. **Obesity (Silver Spring)**, 16, 409-414.
- Pitsavos, C., Panagiotakos, D., Weinem, M. and Stefanadis, C. (2006). Diet, exercise and the metabolic syndrome. **Rev Diabet Stud**, 3, 118-126.
- Rasgon, N.L., Altshuler, L.L., Fairbanks, L., Elman, S., Bitran, J., Labarca, R., et al. (2005). Reproductive function and risk for PCOS in women treated for bipolar disorder. **Bipolar Disord**, 7, 246-259.
- Reaven, G.M. (1988). Banting lecture 1988. Role of insulin resistance in human disease. **Diabetes**, 37, 1595-1607.
- Reaven, G.M. (2003). The insulin resistance syndrome. **Curr Atheroscler Rep**, 5, 364-371.
- Reaven, G.M. (2005). The metabolic syndrome: requiescat in pace. **Clin Chem**, 51, 931-938.
- Rebuffe-Scrive, M., Surwit, R., Feinglos, M., Kuhn, C. and Rodin, J. (1993). Regional fat distribution and metabolism in a new mouse model (C57BL/6J) of non-insulin-dependent diabetes mellitus. **Metabolism**, 42, 1405-1409.
- Romaguera, J., Ortiz, A.P., Roca, F.J., Colon, G. and Suarez, E. (2010). Factors associated with metabolic syndrome in a sample of women in Puerto Rico. **Menopause**, 17, 388-392.
- Rosenson, R.S. (2005). New approaches in the intensive management of cardiovascular risk in the metabolic syndrome. **Curr Probl Cardiol**, 30, 241-279.
- Rossmesl, M., Rim, J.S., Koza, R.A. and Kozak, L.P. (2003). Variation in type 2 diabetes--related traits in mouse strains susceptible to diet-induced obesity. **Diabetes**, 52, 1958-1966.
- Rujjanawate, C., Kanjanapothi, D., Amornlerdpison, D. and Pojanagaroon, S. (2005). Anti-gastric ulcer effect of Kaempferia parviflora. **J Ethnopharmacol**, 102, 120-122.
- Sae-wong, C., Tansakul, P. and Tewtrakul, S. (2009). Anti-inflammatory mechanism of Kaempferia parviflora in murine macrophage cells (RAW 264.7) and in experimental animals. **J Ethnopharmacol**, 124, 576-580.

- Sahin, K., Onderci, M., Tuzcu, M., Ustundag, B., Cikim, G., Ozercan, I.H., et al. (2007). Effect of chromium on carbohydrate and lipid metabolism in a rat model of type 2 diabetes mellitus: the fat-fed, streptozotocin-treated rat. **Metabolism**, 56, 1233-1240.
- Sairam, M.R., Wang, M., Danilovich, N., Javeshghani, D. and Maysinger, D. (2006). Early obesity and age-related mimicry of metabolic syndrome in female mice with sex hormonal imbalances. **Obesity (Silver Spring)**, 14, 1142-1154.
- Samuelsson, A.M., Matthews, P.A., Argenton, M., Christie, M.R., McConnell, J.M., Jansen, E.H., et al. (2008). Diet-induced obesity in female mice leads to offspring hyperphagia, adiposity, hypertension, and insulin resistance: a novel murine model of developmental programming. **Hypertension**, 51, 383-392.
- Sarkar, S., Das, M., Mukhopadhyay, B., Chakrabarti, C.S. and Majumder, P.P. (2006). High prevalence of metabolic syndrome and its correlates in two tribal populations of India and the impact of urbanization. **Indian J Med Res**, 123, 679-686.
- Sato, A., Kawano, H., Notsu, T., Ohta, M., Nakakuki, M., Mizuguchi, K., et al. (2010). Antiobesity effect of eicosapentaenoic acid in high-fat/high-sucrose diet-induced obesity: importance of hepatic lipogenesis. **Diabetes**, 59, 2495-2504.
- Sawasdee, P., Sabphon, C., Sitthiwongwanit, D. and Kokpol, U. (2009). Anticholinesterase activity of 7-methoxyflavones isolated from *Kaempferia parviflora*. **Phytother Res**, 23, 1792-1794.
- Schmidt, R.E., Dorsey, D.A., Beaudet, L.N. and Peterson, R.G. (2003). Analysis of the Zucker Diabetic Fatty (ZDF) type 2 diabetic rat model suggests a neurotrophic role for insulin/IGF-I in diabetic autonomic neuropathy. **Am J Pathol**, 163, 21-28.
- Shima, K., Zhu, M. and Mizuno, A. (1999). Pathoetiology and prevention of NIDDM lessons from the OLETF rat. **J Med Invest**, 46, 121-129.
- Shimada, T., Horikawa, T., Ikeya, Y., Matsuo, H., Kinoshita, K., Taguchi, T., et al. (2011). Preventive effect of *Kaempferia parviflora* ethyl acetate extract and its major components polymethoxyflavonoid on metabolic diseases. **Fitoterapia**, 82, 1272-1278.

- Shoelson, S.E., Herrero, L. and Naaz, A. (2007). Obesity, inflammation, and insulin resistance. **Gastroenterology**, 132, 2169-2180.
- Shulman, G.I. (2000). Cellular mechanisms of insulin resistance. **J Clin Invest**, 106, 171-176.
- Simmons, R.K., Alberti, K.G., Gale, E.A., Colagiuri, S., Tuomilehto, J., Qiao, Q., et al. (2010). The metabolic syndrome: useful concept or clinical tool? Report of a WHO Expert Consultation. **Diabetologia**, 53, 600-605.
- Sookkongwaree, K., Geitmann, M., Roengsumran, S., Petsom, A. and Danielson, U.H. (2006). Inhibition of viral proteases by Zingiberaceae extracts and flavones isolated from *Kaempferia parviflora*. **Pharmazie**, 61, 717-721.
- Sorrentino, M.J. (2005). Implications of the metabolic syndrome: the new epidemic. **Am J Cardiol**, 96, 3E-7E.
- Spranger, J., Kroke, A., Mohlig, M., Hoffmann, K., Bergmann, M.M., Ristow, M., et al. (2003). Inflammatory cytokines and the risk to develop type 2 diabetes: results of the prospective population-based European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam Study. **Diabetes**, 52, 812-817.
- Stanhope, K.L. and Havel, P.J. (2010). Fructose consumption: recent results and their potential implications. **Ann N Y Acad Sci**, 1190, 15-24.
- Storgaard, H., Poulsen, P., Ling, C., Groop, L. and Vaag, A.A. (2006). Genetic and nongenetic determinants of skeletal muscle glucose transporter 4 messenger ribonucleic acid levels and insulin action in twins. **J Clin Endocrinol Metab**, 91, 702-708.
- Sudwan, P., Saenphet, K., Saenphet, S. and Suwansirikul, S. (2006). Effect of *Kaempferia parviflora* Wall. ex. Baker on sexual activity of male rats and its toxicity. **Southeast Asian J Trop Med Public Health**, 37 (3), 210-215.
- Sumiyoshi, M., Sakanaka, M. and Kimura, Y. (2006). Chronic intake of high-fat and high-sucrose diets differentially affects glucose intolerance in mice. **J Nutr**, 136, 582-587.
- Surwit, R.S., Feinglos, M.N., Rodin, J., Sutherland, A., Petro, A.E., Opara, E.C., et al. (1995). Differential effects of fat and sucrose on the development of obesity and diabetes in C57BL/6J and A/J mice. **Metabolism**, 44, 645-651.

- Surwit, R.S., Kuhn, C.M., Cochrane, C., McCubbin, J.A. and Feinglos, M.N. (1988). Diet-induced type II diabetes in C57BL/6J mice. **Diabetes**, 37, 1163-1167.
- Sutthanut, K., Sripanidkulchai, B., Yenjai, C. and Jay, M. (2007). Simultaneous identification and quantitation of 11 flavonoid constituents in *Kaempferia parviflora* by gas chromatography. **J Chromatogr**, 1143, 227-233.
- Tao, R., Ye, F., He, Y., Tian, J., Liu, G., Ji, T., et al. (2009). Improvement of high-fat-diet-induced metabolic syndrome by a compound from *Balanophora polyandra* Griff in mice. **Eur J Pharmacol**, 616, 328-333.
- Tappy, L. and Le, K.A. (2010). Metabolic effects of fructose and the worldwide increase in obesity. **Physiol Rev**, 90, 23-46.
- Termizy, H.M. and Mafauzy, M. (2009). Metabolic syndrome and its characteristics among obese patients attending an obesity clinic. **Singapore Med J**, 50, 390-394.
- Tewtrakul, S. and Subhadhirasakul, S. (2007). Anti-allergic activity of some selected plants in the Zingiberaceae family. **J Ethnopharmacol**, 109, 535-538.
- Tewtrakul, S. and Subhadhirasakul, S. (2008). Effects of compounds from *Kaempferia parviflora* on nitric oxide, prostaglandin E2 and tumor necrosis factor- α productions in RAW264.7 macrophage cells. **J Ethnopharmacol**, 120, 81-84.
- Tilg, H. and Moschen, A.R. (2006). Adipocytokines: mediators linking adipose tissue, inflammation and immunity. **Nat Rev Immunol**, 6, 772-783.
- Vanni, E., Bugianesi, E., Kotronen, A., De Minicis, S., Yki-Jarvinen, H. and Svegliati-Baroni, G. (2010). From the metabolic syndrome to NAFLD or vice versa? **Dig Liver Dis**, 42, 320-330.
- Pudata, V. and Konduru, J. (2011). Metabolic syndrome in endocrine system. **J Diabetes Metab**, 2, 163-168.
- VanSaun, M.N., Lee, I.K., Washington, M.K., Matrisian, L. and Gorden, D.L. (2009). High fat diet induced hepatic steatosis establishes a permissive microenvironment for colorectal metastases and promotes primary dysplasia in a murine model. **Am J Pathol**, 175, 355-364.

- Voshol, P.J., Rensen, P.C., van Dijk, K.W., Romijn, J.A. and Havekes, L.M. (2009). Effect of plasma triglyceride metabolism on lipid storage in adipose tissue: studies using genetically engineered mouse models. **Biochim Biophys Acta**, 1791, 479-485.
- Wannamethee, S.G., Shaper, A.G. and Whincup, P.H. (2006). Modifiable lifestyle factors and the metabolic syndrome in older men: Effects of lifestyle changes. **J Am Geriatr Soc**, 54, 1909-1914.
- Warren, J.M., Henry, C.J. and Simonite, V. (2003). Low glycemic index breakfasts and reduced food intake in preadolescent children. **Pediatrics**, 112, 414-416.
- Wattanapitayakul, S.K., Suwatronnakorn, M., Chularojmontri, L., Herunsalee, A., Niumsakul, S., Charuchongkolwongse, S., et al. (2007). Kaempferia parviflora ethanolic extract promoted nitric oxide production in human umbilical vein endothelial cells. **J Ethnopharmacol**, 110, 559-562.
- Winzell, M.S. and Ahren, B. (2004). The high-fat diet-fed mouse: a model for studying mechanisms and treatment of impaired glucose tolerance and type 2 diabetes. **Diabetes**, 53 (3), 215-219.
- World Health Organization (2008). **Global Health Observatory: Deaths from Cardiovascular disease and diabetes**. Retrieved October 5, 2013, from http://www.who.int/gho/ncd/mortality_morbidity/cvd/en/index.html
- Wu, F., Wang, H., Li, J., Liang, J. and Ma, S. (2012). Homoplantaginins modulates insulin sensitivity in endothelial cells by inhibiting inflammation. **Biol Pharm Bull**, 35, 1171-1177.
- Yasuda, K., Nishikawa, W., Iwanaka, N., Nakamura, E., Seino, Y. and Tsuda, K., Ishihara A (2002). Abnormality in fibre type distribution of soleus and plantaris muscles in non-obese diabetic Goto-Kakizaki rats. **Clin Exp Pharmacol Physiol**, 29, 1001-1008.
- Yenjai, C., Prasanphen, K., Daodee, S., Wongpanich, V. and Kittakoop, P. (2004). Bioactive flavonoids from Kaempferia parviflora. **Fitoterapia**, 75, 89-92.
- Yun, S.N., Ko, S.K., Lee, K.H. and Chung, S.H. (2007). Vinegar-processed ginseng radix improves metabolic syndrome induced by a high fat diet in ICR mice. **Arch Pharm Res**, 30, 587-595.

- Yusuf, S., Hawken, S., Ounpuu, S., Dans, T., Avezum, A., Lanas, F., et al. (2004). Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. **Lancet**, 364, 937-952.
- Zhang, H., Chen, X., Aravindakshan, J. and Sairam, M.R. (2007). Changes in adiponectin and inflammatory genes in response to hormonal imbalances in female mice and exacerbation of depot selective visceral adiposity by high-fat diet: implications for insulin resistance. **Endocrinology**, 148, 5667-5679.
- Zimmet, P., Magliano, D., Matsuzawa, Y., Alberti, G. and Shaw, J. (2005). The metabolic syndrome: a global public health problem and a new definition. **J Atheroscler Thromb**, 12, 295-300.
- Zivkovic, A.M., German, J.B. and Sanyal, A.J. (2007). Comparative review of diets for the metabolic syndrome: implications for nonalcoholic fatty liver disease. **Am J Clin Nutr**, 86, 285-300.

BIOGRAPHY

BIOGRAPHY

Name –Surname	Nittaya Promson
Date of Birth	March 06, 1982
Address	16/1 Moo 5 Bansan, Srisamrong District, Sukhothai Province Thailand 64120
Work Experiences	
2005 - 2011	Kaengkhoi Hospital 107 Moo 8 Taldieo, Kaengkhoi District, Saraburi Province Thailand 18110
Education Background	
2005	B.SC. (Physical Therapy), Naresuan University, Thailand