

LIST OF FIGURES

		Page
Figure 1.1	A conceptual framework of this thesis	4
Figure 2.1	Representation patterns of growth and regression of individual follicles during the estrous cycle in goats with two (A), three (B), four (C), and five (D) waves of follicular development. Arrows indicate the emergence of follicular waves, and different symbols indicate a different follicle in each follicular wave (Ov = ovulation)	10
Figure 2.2	Schematic representation of the different physiological events Occurring during estrous cycle in goat: pattern of follicle development, ovarian cycle and endocrine regulations *Ovulatory follicle(s).	13
Figure 2.3	Evolution of sexual behavior during estrous cycle in Japanese dwarf goat	17
Figure 2.4	Control of compensatory growth	27
Figure 2.5	One stair-step feeding model in rats	40
Figure 2.6	Northern blot analysis of the β -casein gene in mammary tissues collected during the first and second lactation cycles from female rats on a CNR during gestation. The data shown are arbitrary densitometric unit means with standard errors represented by vertical bars (n = 4). *Different from the control (P < 0.05)	41
Figure 2.7	A stair-step compensatory nutrition regimen for dairy heifers	44
Figure 3.1	Body weights gain of goat during 12 weeks (4, 5, 6 months of age)	61
Figure 3.2	Body weights of goat during 24 weeks (7-12 months of age)	63
Figure 3.3	Plasma P4 concentrations of Thai-native female goat during estrous cycle	67

LIST OF FIGURES (CONT.)

		Page
Figure 4.1	Description of timing of injections for the two treatments FSH protocols, laparotomy for detect to corpora hemorrhagica (CH) at 24 h and at 72 h to count corpora lutea (CL)	71
Figure 4.2	Mean (\pm SE) serum P4 concentrations (ng/ml) of goats on days prior injected 2D-FSH (day18) 3D-FSH (day17), hCG (day 20), laparotomy at 24 h (day 21) and 72 h (day 23).a, b Proportion was differ ($P < 0.05$)	76
Figure 5.1	Description of Stair-step feeding regimen, day 0 = heat, B = blood collection	81
Figure 5.2	Means (\pm SEM) serum P4 concentrations (ng/ml) of goats on days 0, 6, 12, 18, and day 21 during 1 st estrous cycle	93
Figure 5.3	Means (\pm SEM) serum P4 concentrations (ng/ml) of goats on days 0, 6, 12, 18, 20, and day 24 (2 nd laparotomy) during 2 nd estrous cycle. ^{a, b} Proportion differ ($P < 0.05$)	94
Figure 5.4	Immunolocalization of PCNA; (A), in a section of the negative staining of follicle; (B), in a section of the positive staining of follicle; (C) and (D), in a section of follicular size 1-3 mm; (E) and (F), in a section of follicular size 4-6 mm; G = granulosa layer; T = Theca layer; size bar indicates 100 μ m; arrows indicate stained PCNA	97
Figure 5.5	The standard curves obtained by plotting the logarithm of mRNA concentration for Bcl-2 versus threshold cycle (Ct) for population quantification by using real time PCR	99
Figure 5.6	The expression of Bcl-2 mRNA in thecal tissue of follicular size 1-6 mm during the 2 nd estrous cycle. Different superscripts denote statistically different values ($P < 0.05$)	100

LIST OF FIGURES (CONT.)

		Page
Figure 5.7	The expression of Bcl-2 mRNA in thecal tissue of follicular size 1-3 mm during the 2 nd estrous cycle	100
Figure 5.8	The expression of Bcl-2 mRNA in thecal tissue of follicular size 4-6 mm during the 2 nd estrous cycle	101
Figure 5.9	The expression of Bcl-2 mRNA in granulosa tissue of follicular size 1-6 mm during the 2 nd estrous cycle. Different superscripts denote statistically different values (P<0.01)	101
Figure 5.10	The expression of Bcl-2 mRNA in granulosa tissue of follicular size 1-3 mm during the 2 nd estrous cycle. Different superscripts denote statistically different values (P<0.01)	102
Figure 5.11	The expression of Bcl-2 mRNA in granulosa tissue of follicular size 4-6 mm during the 2 nd estrous cycle	102
Figure 6	Laparotomy preparation in goat; (A-B), The instruments and inventory for surgery; (C-D), The arrangement for goat on the cradle; (E), The slitting cuts to abdomen; (F), The uterus of goat	136
Figure 7	The procedure of ovarian aspiration and collection in this study; (A-B), Measurement of the ovary; (C-D), Oocyte aspiration from follicle; (E), The ovarian clamp by arteries forceps; (F), Cutting the ovary	137
Figure 8	Oocyte quality assessment by microscope; (A-B), Microscopy for oocyte quality assessment; (C-D), Oocyte image under the 40x objective lens	138

LIST OF FIGURES (CONT.)

		Page
Figure 9	Ovary collection and preparation was used for PCNA detection and Bcl-2 gene expression; (A-B), Preparation of ovarian tissue sections; (C), Deparaffinization; (D-E), Block paraffin embedded; (F), Preparation tissue sections by microtome	139
Figure 10	Immunohistochemical detection of proliferative cell nuclear antigen; (A-B), Preparation before immunostaining; (C-D), Ovarian tissue sections after immunostaining	140
Figure 11	Overview of real time PCR procedures	141
Figure 12	Overview of cDNA synthesis procedures	143
Figure 13	The Graphs screen of the Quantitation window of the mRNA levels of Bcl-2 in this study; (A), Plate diagram; (B), Graphed samples list; (C) Data graph; (D) Standard graph	149
Figure 14	Preparation for detect of Bcl-2 gene expression by real time RT-PCR technique; (A-B), PCR master mix procedure for reaction; (C), Program setting for Real time PCR machine; (D), Plate diagram and Graphed samples list	150