

REFERENCES

- Abbasi T, Tauseef SM and Abbasi SA. Anaerobic digestion for global warming control and energy generation – An overview. *Renewable and Sustainable Energy Reviews*. 2012; 16: 3228-3242.
- Abdulkarim BI and Abdullahi ME. Effect of buffer (NaHCO_3) and waste type in high solid thermophilic anaerobic digestion. *International Journal of ChemTech Research*. 2010; 2: 980-984.
- AOAC International. AOAC official method 2001.11 protein (crude) in animal feed, forage (plant tissue), grain and oilseeds. In official methods of analysis of AOAC International. 18th ed. Gaithersburg, MD. 2005.
- APHA/AWWA/WPCF. Standard Methods for the Examination of Water and Wastewater. 21st ed. Washington, D.C. USA; 2005.
- Argun H, Kargi F, Kapdan IK and Oztekin R. Biohydrogen production by dark fermentation of wheat powder solution: Effect of C/N and C/P ratio on hydrogen yield and formation rate. *International Journal of Hydrogen Energy*. 2008; 33: 1813-1819.
- Bureau of Environment and Health Nonthaburi municipality. Sewage disposal in Nonthaburi municipality [Online]. 2013 [cited 2013 Dec 15]; Available from: <http://www.behn.go.th/dirt.php>
- Chaganti SR, Kim DH and Lalman JA. Dark fermentative hydrogen production by mixed anaerobic cultures: Effect of inoculums treatment methods on hydrogen yield. *Renewable Energy*. 2012; 48: 117-121.
- Chandra R, Takeuchi H and Hasegawa T. Methane production from lignocellulosic agricultural crop wastes: A review in context to second generation of biofuel production. *Renewable and Sustainable Energy Reviews*. 2012; 16: 1462-1476.

- Chen X, Romano RT and Zhang R. Anaerobic digestion of food wastes for biogas production. *International Journal of Agricultural and Biological Engineering*. 2010; 3: 61-72.
- Chen Y, Cheng JJ and Creamer KS. Inhibition of anaerobic digestion process: A review. *Bioresource Technology*. 2008; 99: 4044-4064.
- Chong ML, Sabaratnam V, Shirai Y and Hassan MA. Biohydrogen production from biomass and industrial wastes by dark fermentation. *International Journal of Hydrogen Energy*. 2009; 34: 3277-3287.
- Chu CF, Li YY, Xu KQ, Ebie Y, Inamori Y and Kong HN. A pH- and temperature-phased two stage process for hydrogen and methane production from food waste. *International Journal of Hydrogen Energy*. 2008; 33: 4739-4746.
- Chu CF, Xu KQ, Li YY and Inamori Y. Hydrogen and methane potential based on the nature of food waste materials in a two-stage thermophilic fermentation process. *International Journal of Hydrogen Energy*. 2012; 37: 10611-10618.
- Deublein D and Steinhauser A. *Biogas from waste and renewable resources*. Weinheim: Wiley-VCH; 2008.
- Elbeshbishy E and Nakhla G. Batch anaerobic co-digestion of proteins and carbohydrates. *Bioresource Technology*. 2012; 116: 170-178.
- Escamilla-Alvarado C, Ponce-Noyola T, Rios-Leal E and Poggi-Varaldo HM. A multivariable evaluation of biohydrogen production by solid substrate fermentation of organic municipal wastes in semi-continuous and batch operation. *International Journal of Hydrogen Energy* 2013; 38: 12527-12538.
- Forster-Carneiro T, Perez M and Romero LI. Thermophilic anaerobic digestion of source-sorted organic fraction of municipal solid waste. *Bioresource Technology*. 2008; 99: 6763-6770.
- Gioannis GD, Muntoni A, Polettini A and Pomi R. A review of dark fermentative hydrogen production from biodegradable municipal waste fractions. *Waste Management*. 2013; 33: 1345-1361.

- Giordano A, Cantu C and Spagni A. Monitoring the biochemical hydrogen and methane potential of the two-stage dark-fermentative process. *Bioresource Technology*. 2011; 102: 4474-4479.
- Gomez-Romero J, Gonzalez-Garcia A, Chairez I and Torres L. Selective adaptation of anaerobic microbial community: Biohydrogen production by co-digestion of cheese whey and vegetables fruit waste. *International Journal of Hydrogen Energy*. 2014; 39: 12541-12550.
- Guo YC, Dai Y, Bai YX, Li YH, Fan YT and Hou HW. Co-producing hydrogen and methane from higher-concentration of corn stalk by combining hydrogen fermentation and anaerobic digestion. *International Journal of Hydrogen Energy*. 2014; 39: 14204-14211.
- Han SK and Shin HS. Biohydrogen production by anaerobic fermentation of food waste. *International Journal of Hydrogen Energy*. 2004; 29: 569-577.
- Holladay JD, Hu J, King DL and Wang Y. An overview of hydrogen production technologies. *Catalysis Today*. 2009; 139: 244-260.
- Kalil MS, Alshiyab HS and Yusoff WMW. Effect of nitrogen source and carbon to nitrogen ratio on hydrogen production using *C. acetobutylicum*. *American Journal of Biochemistry and Biotechnology*. 2008; 4: 393-401.
- Kargi F, Eren NS and Ozmihci S. Bio-hydrogen production from cheese whey powder (CWP) solution: Comparison of thermophilic and mesophilic dark fermentations. *International Journal of Hydrogen Energy*. 2012; 37: 8338-8342.
- Kim DH, Kim SH, Jung KW, Kim MS and Shin HS. Effect of initial pH independent of operational pH on hydrogen fermentation of food waste. *Bioresource Technology*. 2011; 102: 8646-8652.
- Kim JK, Nhat L, Chun YN and Kim SW. Hydrogen production conditions from food waste by dark fermentation with *Clostridium beijerinckii* KCTC 1785. *Biotechnology and Bioprocess Engineering*. 2008; 13: 499-504.
- Kim MS, Cha J, Kim DH. Fermentative biohydrogen production from solid wastes. In: Pandey A, Chang JS, Hallenbeck C, Larroche C, editors. *Biohydrogen*. Oxford: Elsevier Science; 2013. p. 259-283.

- Lee KS, Hsu YF, Lo YC, Lin PJ, Lin CY and Chang JS. Exploring optimal environmental factors for fermentative hydrogen production from starch using mixed anaerobic microflora. *International Journal of Hydrogen Energy*. 2008a; 33: 1565-1572.
- Lee ZK, Li SL, Lin JS, Wang YH, Kuo PC and Cheng SS. Effect of pH in fermentation of vegetable kitchen wastes on hydrogen production under a thermophilic condition. *International Journal of Hydrogen Energy*. 2008b; 33: 5234-5241.
- Lin CY and Lay CH. Carbon/nitrogen ratio effect on fermentative hydrogen production by mixed microflora. *International Journal of Hydrogen Energy*. 2004; 29: 41-45.
- Lin Y, Lu F, Shao L and He P. Influence of bicarbonate buffer on the methanogenetic pathway during thermophilic anaerobic digestion. *Bioresource Technology*. 2013; 137: 245-253.
- Liu D, Liu D, Zeng RJ and Angelidaki I. Hydrogen and methane production from household solid waste in the two-stage fermentation process. *Water Research*. 2006; 40: 2230-2236.
- Liu G, Zhang R, El-Mashad HM and Dong R. Effect of feed to inoculum ratios on biogas yields of food and green wastes. *Bioresource Technology*. 2009; 100: 5103-5108.
- Liu X, Li R, Ji M and Han L. Hydrogen and methane production by co-digestion of waste activated sludge and food waste in the two-stage fermentation process: Substrate conversion and energy yield. *Bioresource Technology*. 2013; 146: 317-323.
- Luo G, Xie L, Zhou Q and Angelidaki I. Enhancement of bioenergy production from organic wastes by two-stage anaerobic hydrogen and methane production process. *Bioresource Technology*. 2011; 102: 8700-8706.
- Mecozzi M. Estimation of total carbohydrate amount in environmental samples by the phenol – sulphuric acid method assisted by multivariate calibration. *Chemometrics and Intelligent Laboratory Systems* 2005; 79: 84-90.

- Nathao C, Sirisukpoka U and Pisutpaisal N. Production of hydrogen and methane by one and two stage fermentation of food waste. *International Journal of Hydrogen Energy*. 2013; 38: 15764-15769.
- Nazlina HMY, Nor Aini AR, Ismail F, Yusof MZM and Hassan MA. Effect of different temperature, initial pH and substrate composition on biohydrogen production from food waste in batch fermentation. *Asian Journal of Biotechnology*. 2009; 1: 42-50.
- Ni M, Leung DY, Leung MKH and Sumathy K. An overview of hydrogen production from biomass. *Fuel Processing Technology*. 2006; 87: 461-472.
- O-Thong S, Mamimin C and Prasertsan P. Effect of temperature and initial pH on biohydrogen production from palm oil mill effluent: long term evaluation and microbial community analysis. *Electronic Journal of Biotechnology*. 2011 [cited 2014 Sep 12]; 14. Available from: <http://dx.doi.org/10.2225/vol14-issue5-fulltext-9>
- Oztekin R, Kapdan IK, Kargi F and Argun H. Optimization of media composition for hydrogen gas production from hydrolyzed wheat starch by dark fermentation. *International Journal of Hydrogen Energy*. 2008; 33: 4083-4090.
- Ramos C, Buitron G, Moreno-Andrade I and Chamy R. Effect of the initial total solids concentration and initial pH on the bio-hydrogen production from cafeteria food waste. *International Journal of Hydrogen Energy*. 2012; 37: 13288-13295.
- Raposo F, De la Rubia MA, Fernandez-Cegri V and Borja R. Anaerobic digestion of solid organic substrates in batch mode: An overview relating to methane yields and experimental procedures. *Renewable and Sustainable Energy Reviews*. 2011; 16: 861-877.
- Reungsang A, Pattras S and Sittijunda S. Optimization of key factors affecting methane production from acidic effluent coming from the sugarcane juice hydrogen fermentation process. *Energies*. 2012; 5: 4746-4757.
- Selambo PA, Merrill MD and Logan BE. The use of stainless steel and nickel alloy as low-cost cathodes in microbial electrolysis cells. *Journal of Power Sources*. 2009; 190: 271-278.

- Shin HS, Youn JH and Kim SH. Hydrogen production from food waste in anaerobic mesophilic and thermophilic acidogenesis. *International Journal of Hydrogen Energy*. 2004; 29: 1355-1363.
- Show KY, Lee DJ, Tay JH, Lin CY and Chang JS. Biohydrogen production: Current perspectives and the way forward. *International Journal of Hydrogen Energy*. 2012; 37: 15616-15631.
- Sinha P and Pandey A. An evaluative report and challenges for fermentative biohydrogen production. *International Journal of Hydrogen Energy*. 2011; 36: 7460-7478.
- Sreela-or C, Imai T, Plangklang P and Reungsang A. Optimization of key factors affecting hydrogen production from food waste by anaerobic mixed cultures. *International Journal of Hydrogen Energy*. 2011a; 36: 14120-14133.
- Sreela-or C, Plangklang P, Imai T and Reungsang A. Co-digestion of food waste and sludge for hydrogen production by anaerobic mixed cultures: Statistical key factors optimization. *International Journal of Hydrogen Energy*. 2011b; 36: 14227-14237.
- Tawfik A, Salem A and El-Qelish M. Two stage anaerobic baffled reactors for biohydrogen production from municipal food waste. *Bioresource Technology*. 2011; 102: 8723-8726.
- Thanwiset P, Wirojanagud W and Reungsang A. Effect of hydraulic retention time on hydrogen production and chemical oxygen demand removal from tapioca wastewater using anaerobic mixed cultured in anaerobic baffled reactor (ABR). *International Journal of Hydrogen Energy*. 2012; 37: 15503-15510.
- Valdez-Vazquez I and Poggi-Varaldo HM. Hydrogen production by fermentative consortia. *Renewable and Sustainable Energy Reviews*. 2009; 13: 1000-1013.
- Wang J and Wan W. Combined effects of temperature and pH on biohydrogen production by anaerobic digestion sludge. *International Journal of Hydrogen Energy*. 2011; 35: 3896-3901.

- Wang J and Wan W. Effect of temperature on fermentative hydrogen production by mixed cultures. *International Journal of Hydrogen Energy*. 2008; 33: 5392-5397.
- Wang J and Wan W. Experimental design methods for fermentative hydrogen production: A review. *International Journal of Hydrogen Energy*. 2009a; 34: 235-244.
- Wang J and Wan W. Factors influencing fermentative hydrogen production: A review. *International Journal of Hydrogen Energy*. 2009b; 34: 799-811.
- Wang J and Wan W. Kinetic models for fermentative hydrogen production: A review. *International Journal of Hydrogen Energy*. 2009c; 34: 3313-3323.
- Wang X and Zhao Y. A bench scale study of fermentative hydrogen and methane production from food waste in integrated two-stage process. *International Journal of Hydrogen Energy*. 2009; 34: 245-254.
- Wang YY, Ai P, Hu CX and Zhang YL. Effects of various pretreatment methods of anaerobic mixed microflora on biohydrogen production and the fermentation pathway of glucose. *International Journal of Hydrogen Energy*. 2011; 36: 390-396.
- Wei J, Liu ZT and Zhang X. Biohydrogen production from starch wastewater and application in fuel cell. *International Journal of Hydrogen Energy*. 2010; 35: 2949-2952.
- Yang Z, Guo R, Xu X, Fan X and Luo S. Hydrogen and methane production from lipid-extracted microalgal biomass residues. *International Journal of Hydrogen Energy*. 2011; 36: 3465-3470.
- Zhou P, Elbeshbishy E and Nakhla G. Optimization of biological hydrogen production for anaerobic co-digestion of food waste and wastewater biosolids. *Bioresource Technology*. 2013; 130: 710-718.