List of Tables

Table	
2-1	Comparison between acute and chronic inflammation
2-2	Description of cell derived mediators
2-3	Functional features of molecular mediators of inflammation and oncogenesis
2-4	Biological activities of plants of Ridsiduangmahakan preparation.
2-5	Chemical constituents found in plants of Ridsiduangmahakan preparation
3-1	Plants and part of plants component in Ridsidaungmahakan preparation
3-2	Plants component in Ridsidaungmahakan preparation
4-1	Results of loss on drying, total ash, acid insoluble ash and extractive value of plant materials of Ridsriduangmahakan preparation
4-2	Results of the presence of heavy metals of Ridsriduangmahakan preparation and its ingredients
4-3	Inhibition of Nitric Oxide (NO) production of ethanolic extracts of Ridsriduangmahakan preparation and its plant ingredients at various concentration with calculation of IC ₅₀
4-4	Inhibition on LPS-induced TNF- α release of ethanolic extracts of Ridsiduangmahakan preparation and its plant ingredients at various concentration with calculation of IC ₅₀
4-5	Inhibition on LPS- stimulated PGE ₂ release of ethanolic extracts of Ridsiduangmahakan preparation and its plant ingredients at various concentration with calculation of IC ₅₀
4-6	Comparision of three type of anti-inflammatory activities on NO production, TNF-α release and COX-2 inhibition

List of Tables (continued)

Table		Page
4-7	Cytotoxicity activity at 50 $\mu g/ml$ concentration and IC ₅₀ ($\mu g/ml$) of	
	Ridsiduangmahakan preparation and its plant ingredients extract	
	against two type of colon cancer cell lines (LS174T and SW480)	
	and one type of normal cell line (MRC5)	93
4-8	IC ₅₀ (μg/ml) of the fractions from Ridsiduangmahakan preparation	
	separated by vacuum liquid chromatography on LPS-induced NO	
	release	96
4-9	Anti-inflammatory activities of piperine on LPS-induced from	
	RAW 264.7 cells	98
4-10	Chemical analysis using GC MS of Ridsiduangmahakan	
	preparation	100
4-11	Amount of piperine, IC ₅₀ (µg/ml) of Nitric Oxide (NO) production	
	and PGE ₂ release of the ehtanolic extract of Ridsiduangmahakan	
	preparation under accelerated condition (45 \pm 2°C with 75 \pm 5%	
	RH) for 120 days	102
4-12	Chemical analysis using GC MS of Ridsiduangmahakan	
	preparation under accelerated condition (45 \pm 2°C with 75 \pm 5%	
	RH) for 120 days	106