

เอกสารอ้างอิง

- Aarts MM, Tymianski M (2003) Novel treatment of excitotoxicity: targeted disruption of intracellular signalling from glutamate receptors. *Biochem Pharmacol* 66:877-86.
- Agarwal NB, Agarwal NK, Mediratta PK, Sharma KK (2011) Effect of lamotrigine, oxcarbazepine and topiramate on cognitive functions and oxidative stress in PTZ-kindled mice. *Seizure* 20:257-62.
- Andre VM, Flores-Hernandez J, Cepeda C, Starling AJ, Nguyen S, Lobo MK, et al. (2004) NMDA receptor alterations in neurons from pediatric cortical dysplasia tissue. *Cereb Cortex* 14:634-46.
- Benardo LS (2003) Prevention of epilepsy after head trauma: do we need new drugs or a new approach? *Epilepsia* 44:27-33.
- Brahmane RI, Wanmali VV, Pathak SS, Salwe KJ (2010) Role of cinnarizine and nifedipine on anticonvulsant effect of sodium valproate and carbamazepine in maximal electroshock and pentylenetetrazole model of seizures in mice. *J Pharmacol Pharmacother* 1:78-81.
- Calcagnotto ME, Paredes MF, Tihan T, Barbaro NM, Baraban SC (2005) Dysfunction of synaptic inhibition in epilepsy associated with focal cortical dysplasia. *J Neurosci* 25:9649-57.
- Chang BS, Lowenstein DH (2003) Epilepsy. *N Engl J Med* 349:1257-66.
- Cremer CM, Palomero-Gallagher N, Bidmon HJ, Schleicher A, Speckmann EJ, Zilles K (2009) Pentylenetetrazole-induced seizures affect binding site densities for GABA, glutamate and adenosine receptors in the rat brain. *Neuroscience* 163:490-9.
- de Lanerolle NC, Kim JH, Williamson A, Spencer SS, Zaveri HP, Eid T, et al. (2003) A retrospective analysis of hippocampal pathology in human temporal lobe epilepsy: evidence for distinctive patient subcategories. *Epilepsia* 44:677-87.
- Delorenzo RJ, Sun DA, Deshpande LS (2005) Cellular mechanisms underlying acquired epilepsy: the calcium hypothesis of the induction and maintenance of epilepsy. *Pharmacol Ther* 105:229-66.
- Demeter I, Nagy K, Gellért L, Vécsei L, Fülöp F, Toldi J (2012) A novel kynurenic acid analog (SZR104) inhibits pentylenetetrazole-induced epileptiform seizures. An electrophysiological study : special issue related to Kynurenine. *J Neural Transm* 119:151-4.

- D'Hooge R, Pei YQ, Raes A, Lebrun P, van Bogaert PP, de Deyn PP (1996) Anticonvulsant activity of piperine on seizures induced by excitatory amino acid receptor agonists. *Arzneimittelforschung* 46:557-60.
- Ekonomou A, Angelatou F (1999) Upregulation of NMDA receptors in hippocampus and cortex in the pentylenetetrazol-induced "kindling" model of epilepsy. *Neurochem Res* 24:1515-22.
- Errington AC, Stohr T, Lees G (2005) Voltage gated ion channels: targets for anticonvulsant drugs. *Curr Top Med Chem* 5:15-30.
- French JA, Faught E (2009) Rational polytherapy. *Epilepsia* 50 Suppl 8:63-8.
- Hosford DA, Crain BJ, Cao Z, Bonhaus DW, Friedman AH, Okazaki MM, et al. (1991) Increased AMPA-sensitive quisqualate receptor binding and reduced NMDA receptor binding in epileptic human hippocampus. *J Neurosci* 11:428-34.
- Isokawa M (1996) Decrement of GABA_A receptor-mediated inhibitory postsynaptic currents in dentate granule cells in epileptic hippocampus. *J Neurophysiol* 75:1901-8.
- Jentsch TJ, Poet M, Fuhrmann JC, Zdebik AA (2005) Physiological functions of CLC Cl⁻ channels gleaned from human genetic disease and mouse models. *Annu Rev Physiol* 67:779-807.
- Júnior JS, de Almeida AA, Tomé Ada R, Citó AM, Saffi J, de Freitas RM (2011) Evaluation of possible antioxidant and anticonvulsant effects of the ethyl acetate fraction from *Platonia insignis* Mart. (Bacuri) on epilepsy models. *Epilepsy Behav* 22:678-84.
- Kamphuis W, De Rijk TC, Lopes da Silva FH (1995) Expression of GABA_A receptor subunit mRNAs in hippocampal pyramidal and granular neurons in the kindling model of epileptogenesis: an in situ hybridization study. *Brain Res Mol Brain Res* 31:33-47.
- Koepp MJ, Labbe C, Richardson MP, Brooks DJ, Van Paesschen W, Cunningham VJ, et al. (1997) Regional hippocampal [¹¹C]flumazenil PET in temporal lobe epilepsy with unilateral and bilateral hippocampal sclerosis. *Brain* 120:1865-76.
- Kulak W, Sobaniec W, Wojtal K, Czuczwar SJ (2004) Calcium modulation in epilepsy. *Pol J Pharmacol* 56:29-41.
- Kuwert T, Stodieck SR, Puskas C, Diehl B, Puskas Z, Schuierer G, et al. (1996) Reduced GABA_A receptor density contralateral to a potentially epileptogenic MRI abnormality in a patient with complex partial seizures. *Eur J Nucl Med* 23:95-8.

- Lee KC, Finley PR, Alldredge BK (2003) Risk of seizures associated with psychotropic medications: emphasis on new drugs and new findings. *Expert Opin Drug Saf* 2:233-47.
- Lee KW, Kim HC, Lee SY, Jang CG (2011) Methamphetamine-sensitized mice are accompanied by memory impairment and reduction of N-methyl-D-aspartate receptor ligand binding in the prefrontal cortex and hippocampus. *Neuroscience* 178, 101-7.
- Lerche H, Weber YG, Jurkat-Rott K, Lehmann-Horn F (2005) Ion channel defects in idiopathic epilepsies. *Curr Pharm Des* 11:2737-52.
- Liu FY, Wang XF, Li MW, Li JM, Xi ZQ, Luan GM, et al. (2007) Upregulated expression of postsynaptic density-93 and N-methyl-D-aspartate receptors subunits 2B mRNA in temporal lobe tissue of epilepsy. *Biochem Biophys Res Commun* 358:825-30.
- Loup F, Wieser HG, Yonekawa Y, Aguzzi A, Fritschy JM (2000) Selective alterations in GABA_A receptor subtypes in human temporal lobe epilepsy. *J Neurosci* 20:5401-19.
- Makarska-Bialek K, Kaminski RM, Czuczwar SJ (2005) Influence of the antagonist of the glycine site of NMDA receptors, MRZ 2/576, on the anticonvulsant activity of conventional antiepileptic drugs in mice. *Pharmacol Rep* 57:458-66.
- Mathern GW, Pretorius JK, Leite JP, Kornblum HI, Mendoza D, Lozada A, et al. (1998a). Hippocampal AMPA and NMDA mRNA levels and subunit immunoreactivity in human temporal lobe epilepsy patients and a rodent model of chronic mesial limbic epilepsy. *Epilepsy Res* 32:154-71.
- Mathern GW, Pretorius JK, Mendoza D, Lozada A, Leite JP, Chimelli L, et al. (1998b). Increased hippocampal AMPA and NMDA receptor subunit immunoreactivity in temporal lobe epilepsy patients. *J Neuropathol Exp Neurol* 57:615-34.
- Mathern GW, Pretorius JK, Mendoza D, Leite JP, Chimelli L, Born DE, et al. (1999) Hippocampal N-methyl-D-aspartate receptor subunit mRNA levels in temporal lobe epilepsy patients. *Ann Neurol* 46:343-58.
- McCorry D, Chadwick D, Marson A (2004) Current drug treatment of epilepsy in adults. *Lancet Neurol* 3:729-35.
- Mori A, Kabuto H, Pei YQ (1985) Effects of piperine on convulsions and on brain serotonin and catecholamine levels in E1 mice. *Neurochem Res* 10:1269-75.
- Najm IM, Ying Z, Babb T, Mohamed A, Hadam J, LaPresto E, et al. (2000) Epileptogenicity correlated with increased N-methyl-D-aspartate receptor subunit NR2A/B in human focal cortical dysplasia. *Epilepsia* 41:971-6.

- Neder L, Valente V, Carlotti CG Jr, Leite JP, Assirati JA, Paco-Larson ML, et al. (2002) Glutamate NMDA receptor subunit R1 and GAD mRNA expression in human temporal lobe epilepsy. *Cell Mol Neurobiol* 22:689-98.
- Palizvan MR, Fathollahi Y, Semnanian S (2005) Epileptogenic insult causes a shift in the form of long-term potentiation expression. *Neuroscience* 134, 415-23.
- Palma E, Spinelli G, Torchia G, Martinez-Torres A, Ragozzino D, Miledi R, et al. (2005) Abnormal GABA_A receptors from the human epileptic hippocampal subiculum microtransplanted to *Xenopus oocytes*. *Proc Natl Acad Sci USA* 102:2514-8.
- Pavlova TV, Stepanichev MY, Gekht AB, Gulyaeva NV (2010) Acquisition of an active avoidance reaction in rats and morphological changes in the hippocampus in pentylenetetrazol kindling. *Neurosci Behav Physiol* 40:501-6.
- Porter RJ, Meldrum BS. Antiseizure drugs. In: Katzung BG, editor. *Basic & clinical pharmacology*. New York: McGraw-Hill; 2002. p. 395-418.
- Semah F, Picot MC, Adam C, Broglin D, Arzimanoglou A, Bazin B, Cavalcanti D, Baulac M (1998) Is the underlying cause of epilepsy a major prognostic factor for recurrence? *Neurology* 51, 1256-62.
- Semyanov A, Walker MC, Kullmann DM, Silver RA (2004) Tonically active GABA_A receptors: modulating gain and maintaining the tone. *Trends Neurosci* 27:262-9.
- Sherwin AL (1999). Neuroactive amino acids in focally epileptic human brain: a review. *Neurochem Res* 24:1387-95.
- Slamberova R, Rokyta R (2005) Occurrence of bicuculline-, NMDA- and kainic acid-induced seizures in prenatally methamphetamine-exposed adult male rats. *Naunyn Schmiedebergs Arch Pharmacol* 25:1-6.
- Smith TA (2001) Type A gamma-aminobutyric acid (GABA_A) receptor subunits and benzodiazepine binding: significance to clinical syndromes and their treatment. *Br J Biomed Sci* 58:111-21.
- Szyndler J, Piechal A, Blecharz-Klin K, Skórzewska A, Maciejak P, Walkowiak J, et al. (2006) Effect of kindled seizures on rat behavior in water Morris maze test and amino acid concentrations in brain structures. *Pharmacol Rep* 58, 75-82.
- Tronel S, Sara SJ (2003) Blockade of NMDA receptors in prelimbic cortex induces an enduring amnesia for odor-reward associative learning. *J Neurosci* 23, 5472-6.
- Varho T, Komu M, Sonninen P, Lahdetie J, Holopainen IE (2005) Quantitative HMRS and MRI volumetry indicate neuronal damage in the hippocampus of children with focal epilepsy and infrequent seizures. *Epilepsia* 46:696-703.
- Walker DG, Renwick AG, Hillier K, editors. *Medical pharmacology and therapeutics*. 2nd ed. Edinburgh (UK): McGraw-Hill; 2005.

- Zilles K, Qu MS, Kohling R, Speckmann EJ (1999) Ionotropic glutamate and GABA receptors in human epileptic neocortical tissue: quantitative *in vitro* receptor autoradiography. *Neuroscience* 94:1051-61.
- Zhu LJ, Chen Z, Zhang LS, Xu SJ, Xu AJ, Luo JH (2004) Spatiotemporal changes of the N-methyl-D-aspartate receptor subunit levels in rats with pentylenetetrazole-induced seizures. *Neurosci Lett* 356:53-6.